Ido Efrat · Andrei S. Rapinchuk · Yoav Segev

# **On graphs and valuations**

Received: 26 May 2014 / Accepted: 31 August 2014 Published online: 7 October 2014

**Abstract.** In the last two decades new techniques emerged to construct valuations on an infinite division ring D, given a normal subgroup  $N \subseteq D^{\times}$  of finite index. These techniques were based on the commuting graph of  $D^{\times}/N$  in the case where D is non-commutative, and on the Milnor K-graph on  $D^{\times}/N$ , in the case where D is commutative. In this paper we unify these two approaches and consider V-graphs on  $D^{\times}/N$  and how they lead to valuations. We furthermore generalize previous results to situations of finitely many valuations.

## 1. Introduction

Let D be a division ring, and  $D^{\times}$  be its multiplicative group. We recall that a *valuation* on D is a surjective homomorphism  $v: D^{\times} \to \Gamma$  to a *totally ordered* group  $\Gamma$  such that

$$v(a+b) \ge \min(v(a), v(b))$$
 for all  $a, b \in D^{\times}, b \ne -a$ .

It is well-known that the presence of a non-trivial valuation (or a suitable family of valuations) can be the key to understanding a wide range of properties of D. Therefore one is interested in conditions that guarantee the existence of a valuation on D with nice properties. In the case where D is a commutative field, this was the focus of an extensive study, and several methods to detect valuations were developed, notably the *rigidity* method (see Sect. 1.3.2).

In the non-commutative case a method to construct non-trivial valuations had emerged in [27,34] and [28]. There a valuation v on D was constructed from the assumption that  $D^{\times}$  possesses a finite index normal subgroup N with the quotient

I. Efrat (⊠) · Y. Segev: Department of Mathematics, Ben-Gurion University of the Negev, Beer-Sheva 84105, Israel e-mail: efrat@math.bgu.ac.il

Y. Segev: e-mail: yoavs@math.bgu.ac.il

A. S. Rapinchuk: Department of Mathematics, University of Virginia, Charlottesville, VA 22904, USA. e-mail: asr3x@virginia.edu

Mathematics Subject Classification: Primary 16W60 · 16K20 · 19D45 · 12J10 · Secondary 05C25

Ido Efrat: This research was supported by the Israel Science Foundation (Grant No. 152/13). Andrei S. Rapinchuk: Partially supported by NSF Grant DMS-1301800 and BSF Grant 201049.

 $D^{\times}/N$  having certain properties. In particular the diameter of the *commuting graph* of this quotient should not be too small (see Sect. 1.3.1). Then, in addition, the subgroup N turns out to be open in the topology on D defined by v. Results of this nature were shown to be very useful in the analysis of the normal subgroup structure of  $D^{\times}$ , when D is a finite-dimensional division algebra.

Next, in [12] it was shown that the method of [27,34] and [28] can be used also when *D* is a commutative field, once the above commuting graph (which is the complete graph in the commutative case) is replaced by a graph related to the Milnor *K*-ring of *D* modulo the subgroup *N* of  $D^{\times}$  (see Sect. 1.3.2).

The purpose of this paper is to unify and generalize these two constructions, by axiomatizing this new approach to the construction of valuations on division rings, whether commutative or not. This axiomatization leads us to the notion of a *valuation graph associated with* a finite index normal subgroup N of  $D^{\times}$  (see Sect. 1.1). This notion, in turn, leads to a uniform approach for constructing maps  $\varphi: N \to \Gamma$  having certain properties resembling those of a valuation on D, where  $\Gamma$  is a *partially ordered group* (see Theorem B below). Once such a map  $\varphi$  is obtained machinery from [12,27] can be used to construct a valuation (see Theorem A below). Further, using  $\varphi$  and certain additional hypotheses, and expanding on machinery from [25,28] leads us to new openness results with respect to a *finite set of valuations* (and not a single valuation) for N. This is done in Theorem 10.3 which is then applied to obtain Theorem C.

We hope that our notion of a valuation graph will facilitate future applications of our methods.

We refer to e.g., [11,13,14,22,31,32,39] for general facts and notions in valuation theory. See also Sect. 8.

## 1.1. Valuations via valuation graphs

In what follows let *D* be an infinite division ring and let *N* be a proper finite index normal subgroup of  $D^{\times}$  containing -1. Given  $a \in D^{\times}$  let  $a^* = aN$  be the corresponding coset in  $D^{\times}/N$ .

We consider undirected graphs  $\Delta$  whose vertices are the non-identity elements of the quotient  $D^{\times}/N$ . We denote the distance function on the vertices of  $\Delta$  by  $d(\cdot, \cdot)$ . We set  $d(a^*, b^*) = \infty$  if the vertices  $a^*, b^*$  are not on the same connected component of  $\Delta$ . Also, let diam( $\Delta$ ) be the *diameter* of the graph  $\Delta$ , i.e., diam( $\Delta$ ) = sup  $d(a^*, b^*)$ , with  $a^*, b^*$  ranging over all vertices.

**Definition 1.** We say that  $\Delta$  as above is a V-graph (or a valuation graph) for D if for every  $a, b, c \in D^{\times} \setminus N$  the following three conditions hold:

- (V1) if  $a b \in N$  then  $d(a^*, b^*) \leq 1$ ,
- (V2) if  $d(a^*, b^*) \leq 1$  then  $d((a^{-1})^*, b^*) \leq 1$ ,
- (V3) if  $ab \notin N$  and both  $d(a^*, c^*) \leq 1$  and  $d(a^*b^*, c^*) \leq 1$ , then also  $d(b^*, c^*) \leq 1$ .

We then say that the V-graph  $\Delta$  is *associated* with the normal subgroup N, or that the quotient  $D^{\times}/N$  supports the V-graph  $\Delta$ .

*Remarks.* (1) Axiom (V1) is equivalent to the representative-free condition (V1') if  $1 \in a^* + b^*$  then  $d(a^*, b^*) \leq 1$ .

Here  $a^* + b^*$  is the set of all sums of an element of  $a^*$  and an element of  $b^*$ . The condition  $1 \in a^* + b^*$  is precisely the *Steinberg relation* in the relative version of Milnor's *K*-theory (which will be described in Sect. 1.3.2).

- (2) In Sects. 4, 5 and 6 we actually use the following weaker
  - (V3') if  $d(a^*, (ab)^*) \le 2$  or  $d(a^*, (ba)^*) \le 2$ , then  $d(a^*, b^*) \le 2$ , in place of (V3) (see Remarks 4.2).
- (3) Axiom (V3) is equivalent to the following axiom:
- (V3") For all  $a, b, c \in D^{\times} \setminus N$ , with  $a^* \neq b^*$ , if  $d(a^*, c^*) \leq 1$  and  $d(b^*, c^*) \leq 1$ , then  $d((a^*)^{-1}b^*, c^*) \leq 1$ .

Therefore axioms (V2) and (V3) just mean that, for every  $c \in D^{\times} \setminus N$ , the set

$$\{a^* \in D^{\times} / N \smallsetminus \{1^*\} \mid d(a^*, c^*) \le 1\} \cup \{1^*\}.$$

is a subgroup of N.

One of our results is the following.

**Theorem A.** Let D be an infinite division ring and let  $N \subseteq D^{\times}$  be a finite index normal subgroup containing -1. Assume that  $D^{\times}/N$  supports a V-graph  $\Delta$  of diameter  $\geq 4$ . Then in each of the following situations:

- (i) D is commutative,
- (ii) D is a finite-dimensional division algebra over a field of finite transcendence degree over its prime field,

there exists a non-trivial valuation v of D such that N is open in D with respect to the topology defined by v.

Theorem A is proved at the very end of Sect. 10. One can also think about the openness of the subgroup N of Theorem A as a *congruence subgroup property*, turning Theorem A into a *congruence subgroup theorem* for finite index normal subgroups N of  $D^{\times}$  such that the quotient  $D^{\times}/N$  supports a V-graph of diameter  $\geq 4$ .

It is important to point out that the lower bound of  $\geq 4$  on diam( $\Delta$ ) in Theorem A is optimal. Namely, there are examples where the quotient  $D^{\times}/N$  supports a V-graph of diameter 3, but N is not open with respect to any non-trivial valuation (cf. [27, Example 8.4], [12, Example 7.2], and Examples 11.4 and 11.7).

As noted above, expanding on techniques described earlier in [25,28] we prove a "congruence subgroup property" also in the case where  $D^{\times}/N$  supports a V-graph of diameter 3, but then we require an additional hypothesis (see Theorem C and the paragraphs following it). We do not know if this hypothesis could be removed and we ask:

**Question 2.** Let *D* be an infinite division ring and let  $N \subseteq D^{\times}$  be finite index normal subgroup containing -1. Assume that  $D^{\times}/N$  supports a V-graph  $\Delta$  of diameter  $\geq 3$ , and that one of the following holds:

- (i) D is commutative.
- (ii) D is a finite-dimensional division algebra over a field of finite transcendence degree over its prime field.

Does there exist a non-empty finite set  $\tilde{T}$  of non-trivial valuations of D such that N is open in D with respect to the topology defined by  $\tilde{T}$ ?

A positive answer to Question 2 will have various applications: it will enable one to deduce the existence of valuations in more general situations; it would restrict, in some cases, the structure of  $D^{\times}/N$ , and in particular, would enable one, to complete the proof of the main result of [28] (see Sect. 1.3.1 below) using only the fact that the diameter of minimal non-solvable groups is  $\geq 3$  (see [33]), which is much easier to establish than Property  $(3\frac{1}{2})$  employed in the argument given in [28]). Question 2 was asked in [28, Question 1, p. 932], in the case where  $\Delta$  is the commuting graph of  $D^{\times}/N$ .

## 1.2. Valuation-like and leveled maps

The construction of a valuation from a V-graph  $\Delta$  whose vertices are the nonidentity elements of  $D^{\times}/N$ , is a *two-step procedure*:

**Step 1.** Use the axioms of a V-graph in conjunction with additional hypotheses, in particular assumptions on its diameter, to produce a surjective group homomorphism  $\varphi: N \to \Gamma$  to a partially ordered group  $\Gamma$ , with special properties making it a *valuation-like* or (*strongly*) *leveled* map (see the definitions below).

**Step 2.** Use the maps obtained in *step 1* to construct certain subrings of D with properties analogous to those of valuation rings, and eventually to produce a desired valuation.

We would like to give some indications of the first step as it most directly relies on the formalism of V-graphs, and does not require any additional assumptions on D.

So let *D* be an arbitrary infinite division ring and let  $N \subseteq D^{\times}$  be a finiteindex normal subgroup containing -1. Given a partially ordered group  $\Gamma$  (written additively, but not necessarily commutative) and a homomorphism  $\varphi \colon N \to \Gamma$ , we will frequently use the following (and similar) notation: for  $\alpha \in \Gamma$  we set

$$\Gamma_{<\alpha} = \{\beta \in \Gamma \mid \beta < \alpha\} \text{ and } N_{<\alpha} = \{x \in N \mid \varphi(x) < \alpha\}$$

A homomorphism  $\varphi \colon N \to \Gamma$  to a partially ordered group is said to be a *leveled* map if there exists a non-negative  $\alpha \in \Gamma$  (called a *level* of  $\varphi$ ) such that  $N_{<-\alpha} \neq \emptyset$ and

$$N_{<-\alpha} + 1 \subseteq N_{<-\alpha}.\tag{L}$$

A leveled map to a *totally ordered* group  $\Gamma$  is called a *valuation-like map*. Next, we say that a homomorphism  $\varphi \colon N \to \Gamma$  to a partially ordered group is a *strongly leveled map*, if there exists a non-negative  $\alpha \in \Gamma$  (called a *s-level* of  $\varphi$ ) such that  $N_{>\alpha} \neq \emptyset$  and

$$1 \pm N_{>\alpha} \subseteq N_{\leqslant 0} \tag{SL}$$

(note that while  $-1 \in N$  by our assumption, we are *not* assuming that  $\varphi(-1) = 0$ , which explains the presence of  $\pm$ ). A strongly leveled map to a *totally ordered* group  $\Gamma$  is called a *strong valuation-like map*. We note that by Lemma 9.2(2), *a strongly leveled map of s-level*  $\alpha$  *is a leveled map of level*  $\alpha$ .

One of our main results is:

**Theorem B.** Let D be an infinite division ring and let  $N \subseteq D^{\times}$  be a finite index normal subgroup containing -1. Assume that  $D^{\times}/N$  supports a V-graph  $\Delta$ . Then

(1) *if* diam( $\Delta$ )  $\geq$  3 *then* N *admits a strongly leveled map;* 

(2) *if* diam( $\Delta$ )  $\geq$  4 *then* N *admits a strong valuation-like map;* 

(3) If diam( $\Delta$ )  $\geq$  5 then N admits a strong valuation-like map of s-level 0.

Furthermore, if D is finite-dimensional over an infinite subfield  $k \subseteq F = Z(D)$ , then in all three cases above  $N_{\geq 0}$  contains a basis of D over k.

Parts (1), (2) and (3) of Theorem B are Theorems 4.1, 5.1 and 6.1, respectively. The last part of Theorem B is Corollary 4.6(3).

Using part (1) of Theorem B we prove:

**Theorem C.** Let *D* be a finite-dimensional separable<sup>1</sup> (but not necessarily central) division algebra over an infinite field k of finite transcendence degree over its prime field, and let  $N \subseteq D^{\times}$  be a normal subgroup of finite index containing -1. Assume that  $D^{\times}/N$  supports a V-graph of diameter  $\geq 3$ , and let  $\varphi \colon N \to \Gamma$  be the strongly leveled map obtained in Theorem B(1).

Suppose in addition that the subgroup  $\varphi(N \cap k^{\times}) \subseteq \Gamma$  is totally ordered. Then

- (1) the restriction  $\varphi_k = \varphi \upharpoonright_{(N \cap k^{\times})}$  is a strong valuation-like map;
- (2) there exists a height one valuation v of k such that N ∩ k<sup>×</sup> is open in the v-adic topology on k<sup>×</sup>;
- (3) there exists a non-empty finite set T of valuations of the center F = Z(D)extending v such that  $|T| \leq [F:k]$ , and such that each  $w \in T$  uniquely extends to a valuation  $\tilde{w}$  of D, and N is open in  $D^{\times}$  in the  $\tilde{T}$ -adic topology, where  $\tilde{T} = \{\tilde{w} \mid w \in T\}$ .

We mention that the hypothesis in Theorem C that  $\varphi(N \cap k^{\times}) \subseteq \Gamma$  is totally ordered is used to obtain part C(1). Then, C(2) follows from Theorem 9.5. In Theorem 9.5 there is no use of the notion of V-graphs. Next, Theorem B(1), the hypothesis that  $\varphi(N \cap k^{\times})$  is totally ordered and Proposition 10.4, enable one to use Theorem 10.3, which yields C(3). In Theorem 10.3 there is no use of the notion of V-graphs.

Theorem C is proved at the end of Sect. 10. As noted above, we do not know whether the hypothesis in Theorem C, that  $\varphi(N \cap k^{\times})$  is totally ordered, can be removed. The examples in Sect. 11 show that Theorem C considers situations which are more general than those considered by Theorem A. Finally, we draw the attention of the reader to Theorems 7.3 and 7.5.

Going back to V-graphs, notice that for every D and N as above there is a canonical (minimal) V-graph supported by  $D^{\times}/N$ , namely, the intersection of all

<sup>&</sup>lt;sup>1</sup> This means that the center F of D is a separable extension of k, cf. [23].

V-graphs supported by  $D^{\times}/N$ . In Theorems A–C, the assumptions that  $D^{\times}/N$  supports a V-graph with sufficiently large diameter can therefore be replaced by the assumption that this canonical V-graph has such a diameter.

## 1.3. The origin and examples of V-graphs

The notion of a V-graph has two prototypes: the commuting graph and the Milnor K-graph—the axioms in Definition 1 simply postulate the properties that were used to produce valuations in these two cases. It is quite remarkable that the essential properties turned out to be identical in these two quite different situations, which we will now review to put our results in perspective.

1.3.1. Commuting graphs. Let G be a finite group. The commuting graph  $\Delta_G$  is the undirected graph whose vertex set consists of the non-identity elements of G, and in which two vertices are connected by an edge if and only if the corresponding elements commute in G. Given an infinite division ring D and a normal subgroup  $N \subseteq D^{\times}$  (not necessarily of finite index), it is an easy exercise to check that  $\Delta_{D^{\times}/N}$ is a V-graph associated with the normal subgroup N. Indeed see [34, Remark 2.2] for (V1), (V2) is trivial and (V3) is straightforward.

Let *D* be any *finite dimensional* division algebra and *N* a *finite index* normal subgroup of  $D^{\times}$ . Set  $\Delta := \Delta_{D^{\times}/N}$ . As mentioned in Sect. 1.2, producing a valuation on *D* using the commuting graph  $\Delta$  requires *two steps*.

The basic machinery for implementing *step 1* was developed in [34]. Subsequently, it was further developed and improved in [27] and [28]. Cumulatively, the results obtained in [27,34] and [28] yield a proof of Theorem B in the case where  $\Delta = \Delta_{D^{\times}/N}$ .

In [35] it was shown (using the classification of finite simple groups (CFSG)) that if L is а non-abelian finite simple group, then either diam( $\Delta_L$ )  $\geq$  5, or  $\Delta_L$  is *balanced* (see [34] for the definition of a balanced commuting graph). This result, together with [34, Theorem A] proved [35, Theorem 3, p. 126], which states that for D and N as above,  $D^{\times}/N$  is not a non-abelian finite simple group. This last result was conjectured in [26], and in view of the reduction obtained therein, concluded the proof of the Margulis-Platonov conjecture (MP) for inner forms of anisotropic groups of type  $A_n$ , i.e. groups of the form  $SL_{1,D}$ where D is finite-dimensional division algebra over a global field K [see [24, Ch. 9] and Appendix A in [27] for a discussion of (MP)].

A systematic use of valuations in this context was introduced in [27], although some features of valuations can already be seen in [34] (like the local ring constructed in 10 of [34]—see Appendix B in [27] for a discussion of this ring in the context of valuations). Indeed valuation theory together with the machinery developed in [27] supplies the tools adequate for handling *step 2*. This, together with improved results for *step 1* in [27], enabled the second and third-named authors to construct, under the hypothesis that diam $(\Delta) \ge 4$ , a valuation v on D such that Nis v-adically open.

The next major development was the result proved in [28], stating that *for any finite-dimensional division algebra D over an arbitrary field, every finite quotient of* 

the multiplicative group  $D^{\times}$  is solvable. This was based on upgrading the techniques both in step 1 and step 2 in the context of a new property  $(3\frac{1}{2})$  for  $\Delta_{D^{\times}/N}$  (see Sect. 7) and establishing this property for minimal non-solvable groups.

In [25], improving and expanding on machinery for *step 2* and using the results on *step 1* from [28], it was shown that if the center of *D* is a global field, and if diam( $\Delta$ )  $\geq$  3, then there is a *finite set* of valuations on *D* so that *N* is open in the topology on *D* defined by this set. This also enabled the proof of (MP) for inner forms of anisotropic groups of type  $A_n$ , using the fact (which relies on CFSG) that all finite simple groups are generated by two elements.

*1.3.2. Milnor K-graphs.* To describe the second prototype of a V-graph, recall from [11, Ch. 24] the definition of the Milnor *K*-groups of a field *F* relative to a subgroup *N* of  $F^{\times}$ . For a non-negative integer *r* let  $K_r^M(F)/N$  be the quotient of the *r*th tensor power

$$(F^{\times}/N)^{\otimes r} = (F^{\times}/N) \otimes_{\mathbb{Z}} \cdots \otimes_{\mathbb{Z}} (F^{\times}/N)$$

by the subgroup generated by all *Steinberg elements*, i.e., elementary tensors  $a_1N \otimes \cdots \otimes a_rN$  such that  $1 \in a_iN + a_jN$  for some  $1 \leq i < j \leq r$  (compare with axiom (V1') above). The tensor product induces on

$$K^M_*(F)/N := \bigoplus_{r=0}^{\infty} K^M_r(F)/N$$

the structure of a graded ring. It is called the *Milnor K-ring of F modulo* N. Equivalently,  $K_*^M(F)/N$  is the quotient of the (classical) Milnor K-ring  $K_*^M(F) (=K_*^M(F)/\{1\})$  by the graded ideal generated by N, considered as a subgroup of  $F^* = K_1^M(F)$ . Following traditional notation, the image of  $a_1N \otimes \cdots \otimes a_rN$  in  $K_r^M(F)/N$  (where  $a_1, \ldots, a_r \in F^*$ ) will be denoted by  $\{a_1, \ldots, a_r\}_N$ .

Now one defines the *Milnor K-graph of F modulo N* to be the undirected graph whose vertices are the non-identity elements of  $F^{\times}/N$ , and where vertices aN and bN are connected by an edge if and only if  $\{a, b\}_N = 0$  in  $K_2^M(F)/N$ . It follows from [12, Lemma 2.1] that this is indeed a V-graph. The main result of [12] is just Theorem A for this V-graph on D = F.

We point out that connections between existence of arithmetically interesting valuations on a (commutative) field F and Milnor K-theory were noted before. Notably, a series of works by Ware [40], Arason, Elman, Jacob, and Hwang [1,15–17] developed a method to produce valuations on F using so-called *rigid subgroups* of  $F^{\times}$ . In [9] it was shown that this method can be naturally interpreted in terms of relative Milnor K-theory. In fact, this was one of the main motivations for introducing the relative Milnor K-ring functor  $K_*^M(F)/N$ . This new perspective opened the way to further strengthening of the rigidity method for producing valuations in [10] and [11, Ch. 26], and recently in [36] and [37]. Another powerful approach for the detection of valuations on fields related to Milnor K-theory was developed by Bogomolov and Tschinkel (see e.g., [4–6]).

For some other approaches for the construction of valuations on fields see [19] and [2].

As we see, the situations where the commuting graphs and the Milnor K-graphs were used to construct valuations are indeed quite different (in fact, disjoint), while the results and the techniques involved in their proofs are very much parallel. This observation led us to generalize and axiomatize these considerations which resulted in the notion of a V-graph.

## 2. Partially preordered and ordered groups

The goal of this paper is to construct *valuations* on an infinite, finite dimensional division algebra D, given a finite index normal subgroup  $N \subseteq D^{\times}$  such that the quotients  $D^{\times}/N$  supports a V-graph. As indicated in Sect. 1.2 of the introduction, this process is carried out in two main steps. The purpose of this section is to give more details about *step 1*. Since Step 1 leads to partially preordered and ordered groups, we discuss in this section such groups in more detail.

So for  $x \in D^{\times}$ , let  $x^*$  denote its image in  $D^{\times}/N$ . Now *step 1* is achieved using the following further steps:

**Step 1a.** Given  $y \in D^{\times}$  we define an *invariant binary relation*  $\mathfrak{P}_{y^*}$  on N such that  $(N, \mathfrak{P}_{y^*})$  is a *partially preordered group*. The relation  $\mathfrak{P}_{y^*}$  will only depend on the coset  $y^* = yN$  and not on the coset representative y. This step *does not* require  $D^{\times}/N$  to support a V-graph. Thus below we define and discuss all notions relevant to step 1a.

**Step 1b.** Given the partially preordered group  $(N, \mathfrak{P}_{v^*})$  of step 1a we define

$$U_{\mathcal{V}^*} := \{ n \in N \mid 1 \mathfrak{P}_{\mathcal{V}^*} n \text{ and } n \mathfrak{P}_{\mathcal{V}^*} 1 \}.$$

We show that  $U_{y^*} \leq N$  and that  $\Gamma_{y^*} := N/U_{y^*}$  is a partially ordered group. The order relation  $\leq_{y^*}$  on  $\Gamma_{y^*}$  is given by  $mU_{y^*} \leq_{y^*} nU_{y^*}$  iff  $m \mathfrak{P}_{y^*} n$ , where  $m, n \in N$ . We let

$$\varphi_{y^*}: N \to \Gamma_{y^*},$$

be the canonical homomorphism. This step as well *does not* require  $D^{\times}/N$  to support a V-graph. Thus below we also discuss all notions relevant to step 1b. **Step 1c.** We show that if  $-1 \in N$  and  $D^{\times}/N$  supports a V-graph  $\Delta$ , then the assertions of Theorem B of the introduction hold, where the asserted map in parts (1)–(3) of Theorem B is  $\varphi_{\gamma^*}$ , for an appropriate  $\gamma^*$ .

*Remark 2.1.* Step 1, and all its parts above, *do not* require that *D* be finite dimensional. It is only in Step 2, when we construct valuations on *D*, that we assume that *D* is finite-dimensional.

## Partially preordered and ordered groups

Let  $\Gamma$  be a group and let  $\leq$  be a partial order on  $\Gamma$ . We say that  $(\Gamma, \leq)$  is a *partially ordered group* if

$$\alpha \leq \gamma \text{ and } \beta \leq \delta \implies \alpha + \beta \leq \gamma + \delta,$$

for all  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta \in \Gamma$ . Notice that we always use *additive notation* for a partially ordered group ( $\Gamma$ ,  $\leq$ ), even though  $\Gamma$  need not be commutative. As before for  $\alpha \in \Gamma$  we write  $\Gamma_{>\alpha} = \{\beta \in \Gamma \mid \beta > \alpha\}$ .

A partial order  $\leq$  on  $\Gamma$  is called *trivial* if  $\alpha \leq \beta \iff \alpha = \beta$ , for all  $\alpha, \beta \in \Gamma$ . Next let N be an arbitrary group. We study pullbacks of partial orderings on

groups to N.

We say that a binary relation  $\mathfrak{P}$  on *N* is *invariant* if

$$m \mathfrak{P} n \implies sm \mathfrak{P} sn \text{ and } ms \mathfrak{P} ns,$$
 (2.1)

for all  $m, n, s \in N$ .

**Lemma 2.2.** Let  $\mathfrak{P}$  be a reflexive and transitive binary relation on N. Then  $\mathfrak{P}$  is invariant if and only if

$$m \mathfrak{P} n \text{ and } s \mathfrak{P} t \implies ms \mathfrak{P} nt, \quad \forall m, n, s, t \in N.$$

Proof. The "if" part is immediate.

For the "only if" part let  $m, n, s, t \in N$  and suppose that  $m \mathfrak{P} n$  and  $s \mathfrak{P} t$ . By the invariance,  $ms \mathfrak{P} ns \mathfrak{P} nt$ , so by the transitivity,  $ms \mathfrak{P} nt$ .

When the conditions of Lemma 2.2 are satisfied we will say that  $(N, \mathfrak{P})$  is a *partially preordered group*.

**Lemma 2.3.** The following conditions on a binary relation  $\mathfrak{P}$  on N are equivalent:

- (1)  $\mathfrak{P}$  is reflexive, transitive and invariant (i.e.,  $(N, \mathfrak{P})$  is a partially preordered group).
- (2) There exist a partially ordered group  $(\Gamma, \leq)$  and a group-epimorphism  $\varphi \colon N \to \Gamma$  such that  $m \mathfrak{P} n \iff \varphi(m) \leq \varphi(n)$  for all  $m, n \in N$ .

Moreover, when these conditions are satisfied, the kernel of  $\varphi$  is

$$U = \{n \in N \mid 1 \mathfrak{P} n \text{ and } n \mathfrak{P} 1\}.$$

*Proof.* (1) $\Rightarrow$ (2): We first show that *U* is a subgroup of *N*. By the reflexivity,  $1 \in U$ . If  $m, n \in U$ , then by Lemma 2.2,  $mn \mathfrak{P} 1 \cdot 1$  and  $1 \cdot 1 \mathfrak{P} mn$ , so  $mn \in U$ . Also, the invariance gives  $m^{-1}m \mathfrak{P} m^{-1} \cdot 1$  and  $1 \cdot m^{-1} \mathfrak{P} mm^{-1}$ , so  $m^{-1} \in U$ , as desired.

Next we observe that U is normal in N. Indeed, let  $m \in U$  and  $n \in N$ . Then, by the invariance,  $n^{-1}mn \mathfrak{P} n^{-1} \cdot 1 \cdot n$  and  $n^{-1} \cdot 1 \cdot n \mathfrak{P} n^{-1}mn$ , so  $n^{-1}mn \in U$ .

We further notice that the relation  $m \mathfrak{P} n$  depends only on the cosets of m and n modulo U. Indeed let  $u, v \in U$ . Then  $u \mathfrak{P} 1$  and  $1 \mathfrak{P} v$ . Therefore  $m \mathfrak{P} n$  implies (by the invariance) that  $mu \cdot 1 \mathfrak{P} n \cdot 1 \cdot v$ .

Now set  $\Gamma = N/U$  and let  $\varphi \colon N \to \Gamma$  be the canonical epimorphism. By what we have just seen, we may define a binary relation  $\leq$  on  $\Gamma$  by

$$\varphi(m) \le \varphi(n) \iff m \mathfrak{P} n \text{ for } m, n \in \mathbb{N}.$$
 (2.2)

Since  $\mathfrak{P}$  is reflexive and transitive, so is  $\leq$ . Also, if  $m, n \in N$  and  $\varphi(m) \leq \varphi(n) \leq \varphi(m)$ , then  $m \mathfrak{P} n \mathfrak{P} m$ . Multiplying by  $m^{-1}$  on the left, we see that  $1 \mathfrak{P} m^{-1} n \mathfrak{P} 1$ , so  $m^{-1} n \in U$ , whence  $\varphi(m) = \varphi(n)$ . Thus  $\leq$  is a partial order.

Finally, the fact that  $(\Gamma, \leq)$  is a partially ordered *group* follows from Lemma 2.2. (2) $\Rightarrow$ (1): Straightforward. *Remark 2.4.* Notice that in Lemma 2.3, and throughout this article, we use multiplicative notation for N and additive notation for  $\Gamma$ .

## **3.** The group $\Gamma_{y^*}$ and the map $\varphi_{y^*}: N \to \Gamma_{y^*}$

In this section D is an arbitrary infinite division algebra (not necessarily finite dimensional over its center), and N is a normal subgroup of  $D^{\times}$  of finite index. Note that in this section we make *no additional hypotheses*. In particular, we do not assume that  $D^{\times}/N$  supports a V-graph. As before, for  $a \in D^{\times}$  denote by  $a^*$  the image of a in  $D^{\times}/N$  under the canonical homomorphism.

Our goal in this section is, using only the above information, to construct for any  $y \in D^{\times} \setminus N$  a map

$$\varphi_{y^*}: N \to \Gamma_{y^*},$$

where here  $\Gamma_{y^*}$  is a partially ordered group.

We start with defining a binary relation  $\mathfrak{P}_{y^*}$  on N by

$$m \mathfrak{P}_{y^*} n \iff N(my) \subseteq N(ny)$$

A crucial role is played by the sets N(y): for  $y \in D^{\times}$  we let

$$N(y) := \{ n \in N \mid y + n \in N \} = N \cap (N - y).$$

Lemma 3.1 below gives some basic properties of the sets N(y). Then Corollary 3.2(2) shows that  $\mathfrak{P}_{y^*}$  depends only on the coset  $y^* = Ny$  and not on the coset representative *y*. Furthermore, Corollary 3.2(1) shows that  $\mathfrak{P}_{y^*}$  does not depend on the "side", i.e., the relation defined by  $N(ym) \subseteq N(yn)$  coincides with  $\mathfrak{P}_{y^*}$ . In Lemma 3.3 we see that  $(N, \mathfrak{P}_{y^*})$  is a partially preordered group, and then we use Lemma 2.3 to define the partially ordered group  $(\Gamma_{y^*}, \leq_{y^*})$  and the map  $\varphi_{y^*}$ .

**Lemma 3.1.** [27, Lemma 6.3]. Let  $y \in D^{\times} \setminus N$  and  $n \in N$ . Then

- (1) N(ny) = nN(y) and N(yn) = N(y)n;
- (2)  $N(y^x) = x^{-1}N(y)x$ , for all  $x \in D^{\times}$ ;
- (3)  $N(y) \neq \emptyset$ ;
- (4) if  $n \in N(y^{-1})$ , then  $y + n^{-1} \in Ny$ . Consequently,  $n^{-1} \notin N(y)$ . In particular  $\emptyset \subseteq N(y) \subseteq N$ .

*Proof.* (1):  $N(ny) = N \cap (N - ny) = nN \cap (nN - ny) = n(N \cap (N - y)) = nN(y)$ , and similarly for N(yn).

(2):  $N(y^x) = N \cap (N - y^x) = N^x \cap (N^x - y^x) = N(y)^x$ .

- (3): This is an immediate consequence of the fact that D = N N (cf. [3,38]).
- (4): The first part of (4) follows from the definition of N(y) and the rest of (4) is a consequence of the first part, and of (3).

**Corollary 3.2.** Let  $y \in D^{\times} \setminus N$  and  $m, n \in N$ . Then

(1)  $N(my) \subseteq N(ny)$  if and only if  $N(ym) \subseteq N(yn)$ ;

- (2) if  $N(my) \subseteq N(ny)$ , then  $N(my') \subseteq N(ny')$  for all  $y' \in yN = Ny$ .
- *Proof.* (1): The first inclusion is equivalent to  $yN(my)y^{-1} \subseteq yN(ny)y^{-1}$ , which by Lemma 3.1(2) is just the second inclusion.
- (2): Multiply the inclusion  $N(my) \subseteq N(ny)$  on the right by elements of N and use Lemma 3.1(1).

We now show that  $(N, \mathfrak{P}_{v^*})$  is a partially preordered group, for any  $y \in D^{\times} \setminus N$ .

**Lemma 3.3.** [27, Lemma 6.4]. For any  $y \in D^{\times} \setminus N$ , the relation  $\mathfrak{P} := \mathfrak{P}_{y^*}$  has the following properties

(1)  $\mathfrak{P}$  is reflexive and transitive;

(2)  $\mathfrak{P}$  is invariant.

*Proof.* (1): This is immediate from the definition of  $\mathfrak{P}$ .

(2): Recall from Eq. (2.1) the notion of an invariant relation. Let  $m, n, s \in N$ , and assume that  $m \mathfrak{P} n$ . By the definition of  $\mathfrak{P}$  and Lemma 3.1(1),  $sm \mathfrak{P} sn$ . Next we have  $N(my') \subseteq N(ny')$ , for all  $y' \in Ny$ , by Corollary 3.2(2). Taking y' = sy, we see that  $N(msy) \subseteq N(nsy)$ , i.e.,  $ms \mathfrak{P} ns$ .

Since  $(N, \mathfrak{P}_{y^*})$  is a partially preordered group, Lemma 2.3 yields a partially ordered group  $(\Gamma_{y^*}, \leq_{y^*})$  and a group epimorphism  $\varphi_{y^*} \colon N \to \Gamma_{y^*}$  such that

$$\varphi_{y^*}(m) \leq_{y^*} \varphi_{y^*}(n) \iff m \mathfrak{P}_{y^*} n \iff N(my) \subseteq N(ny),$$
(3.1)

for all  $m, n \in N$ . More concretely,

$$\Gamma_{y^*} = N/U_{y^*},$$

where  $U_{y^*}$  is the normal subgroup

$$U_{y^*} := \{n \in N \mid n \mathfrak{P}_{y^*} \mid 1 \text{ and } 1 \mathfrak{P}_{y^*} n\} = \{n \in N \mid N(ny) = N(y)\}$$

of N, and

$$\varphi_{y^*}: N \to N/U_{y^*},$$

is the canonical homomorphism.

Next, we let

$$\mathbb{P}_{\mathbf{v}^*} = \{ b \in N\mathbf{v} \mid 1 \in N(b) \}.$$

Note that it follows from Lemma 3.1(3) and Lemma 3.1(1) that  $\mathbb{P}_{v^*} \neq \emptyset$ .

One has  $b \in \mathbb{P}_{y^*}$  if and only if  $yb^{-1} \in N$  and  $yb^{-1} \in Nyb^{-1} - y = N - y$ , or equivalently,  $yb^{-1} \in N(y)$ . Therefore

$$\mathbb{P}_{y^*} = N(y)^{-1}y,$$

and similarly,

$$\mathbb{P}_{y^*} = yN(y)^{-1}.$$

**Lemma 3.4.** Let  $y \in D^{\times} \setminus N$ . The following conditions on  $m, n \in N$  are equivalent:

(1)  $m \mathfrak{P}_{y^*} n$ . (2)  $\varphi_{y^*}(m) \leq_{y^*} \varphi_{y^*}(n)$ . (3)  $N(my') \subseteq N(ny')$  for all  $y' \in Ny$ . (4)  $m \in N(nb)$  for all  $b \in \mathbb{P}_{y^*}$ . (5)  $nb + m \in N$  for all  $b \in \mathbb{P}_{y^*}$ . (6)  $n\mathbb{P}_{y^*} \subseteq m\mathbb{P}_{y^*}$ . (7) For all  $y' \in Ny$ , if  $n \in N(y')$ , then  $m \in N(y')$ .

*Proof.* To simplify notation we denote  $\mathbb{P} := \mathbb{P}_{v^*}$ .

(1) $\Leftrightarrow$ (2): This is immediate from the definition of  $\varphi_{y^*}$  and of  $\leq_{y^*}$ . (1) $\Rightarrow$ (3): This is Corollary 3.2(2). (3) $\Rightarrow$ (4): Let  $b \in \mathbb{P}$ . As  $\mathbb{P} \subseteq Ny$  we have  $N(mb) \subseteq N(nb)$ , by (3). But  $m \in N(mb)$ , so  $m \in N(nb)$ (4) $\Rightarrow$ (1): We need to show that (4) implies  $mN(y) \subseteq N(ny)$ . So take  $s \in N(y)$ . Then  $b := ys^{-1} \in \mathbb{P}$ . By (4),  $m \in N(nb) = N(nys^{-1})$  whence  $ms \in N(ny)$ . (4) $\Leftrightarrow$ (5): As  $m \in N$ , this is immediate from the definition of N(nb). (4) $\Leftrightarrow$ (6): Condition (4) is equivalent to  $1 \in N(m^{-1}nb)$  for all  $b \in \mathbb{P}$ . As  $m^{-1}n\mathbb{P} \subseteq Ny$ , this means that  $m^{-1}n\mathbb{P} \subseteq \mathbb{P}$ , as desired. (3) $\Rightarrow$ (7): Let  $y' \in Ny$ . By (3) (with y' replaced by  $n^{-1}y'$ ),  $N(mn^{-1}y') \subseteq N(y')$ . Now if  $n \in N(y')$ , then  $m \in N(mn^{-1}y')$ , and it follows that  $m \in N(y')$ . (7) $\Rightarrow$ (1): Let  $s \in N(my)$ . Then  $1 \in N(mys^{-1})$ , so  $n \in N(y')$ , where  $y' := mys^{-1}n \in Ny$ . By (7),  $m \in N(y')$ . Thus  $1 \in N(ys^{-1}n)$  and therefore  $n^{-1}s \in N(y)$ . We conclude that  $s \in nN(y) = N(ny)$ .

To continue the discussion we recall some notation. Given  $y \in D^{\times} \setminus N$ , we let

 $N_{\leq_{y^*}\gamma} = \{m \in N \mid \varphi_{y^*}(m) \leq \gamma\}$  for  $\gamma \in \Gamma_{y^*}$ ,

and the sets  $N_{<_{v^*}\gamma}$ ,  $N_{>_{v^*}\gamma}$  etc. are defined similarly.

**Corollary 3.5.** Let  $y \in D^{\times} \setminus N$ . Then

(1)  $N_{\leq_{v^*} 0} = \{m \in N \mid m \in N(b), \text{ for all } b \in \mathbb{P}_{v^*}\};$ 

(2) for all  $y' \in Ny$ , if  $n \in N(y')$ ,  $m \in N$  and  $\varphi_{y^*}(m) \leq_{y^*} \varphi_{y^*}(n)$ , then  $m \in N(y')$ .

*Proof.* (1): This follows from the equivalence of (2) and (4) in Lemma 3.4, with n = 1.

(2): This follows from the implication  $(2) \Rightarrow (7)$  in Lemma 3.4.

The following proposition gives an important and surprising property of the sets N(y),  $y \in D^{\times} \setminus N$ .

**Proposition 3.6.** Assume that D is a finite-dimensional (but not necessarily central) division algebra over an infinite field k, and let  $N \subseteq D^{\times}$  be a subgroup of finite index. Then for any  $a \in D^{\times} \setminus N$ , each of the sets N(a) and  $N(a)^{-1} := \{n^{-1} \mid n \in N(a)\}$  contains a basis of D over k.

*Proof.* We first recall the following Proposition due to Turnwald.

**Proposition 3.7.** [38, Proposition 1.3]. Let *D* be an infinite division ring and let  $N \subseteq D^{\times}$  be a subgroup of finite index. Then for any  $x_1, \ldots, x_m \in D^{\times}$  there exists  $c \in D^{\times}$  such that  $1 + cx_j \in N$  for all  $j = 1, \ldots, m$ .

We now continue with the proof of Proposition 3.6. Let  $x_1, \ldots, x_\ell$  be a left transversal for *N* in  $D^{\times}$ .

We claim that there is a basis  $y_1, \ldots, y_m$  of D over k contained in N. Indeed, let  $H = \operatorname{GL}_{1,D}$  be the algebraic group associated with D. Then  $H = \bigcup x_i \overline{N}$ , where  $\overline{N}$  denotes the Zariski-closure of N in H. Since H is connected, we conclude that  $H = \overline{N}$ . Then N is also Zariski-dense in D. The existence of a required basis now follows from the fact that the *m*-tuples  $(y_1, \ldots, y_m) \in D^m$  that constitute a basis of D over k form a Zariski-open subset.

Now consider the finite set of elements

$$x_i y_j^{-1}$$
 where  $i = 1, ..., \ell; j = 1, ..., m.$  (3.2)

By Proposition 3.7, there exists  $c \in D^{\times}$  such that

$$1 + cx_i y_i^{-1} \in N$$
 for all  $i, j$ .

Since  $x_1, \ldots, x_\ell$  is a left transversal for N in  $D^{\times}$ , there exists  $i_0 \in \{1, \ldots, \ell\}$  and  $s \in N$ , such that  $c^{-1}a = x_{i_0}s^{-1}$ , that is  $cx_{i_0} = as$ . Then

$$1 + asy_i^{-1} \in N$$
 for all  $j = 1, ..., m$ .

It follows that the elements  $y_1s^{-1}, \ldots, y_ms^{-1}$ , which form a basis of *D* over *k*, are all contained in N(a). To show that  $N(a)^{-1}$  contains a basis, we apply the same argument to the family

$$x_i y_j$$
 where  $i = 1, ..., \ell; j = 1, ..., m$ 

in place of the family in Eq. (3.2), to obtain that  $sy_1, \ldots, sy_m$  is a basis of *D* over *k* contained in  $N(a)^{-1}$ .

## 4. The case where diam( $\Delta$ ) $\geq$ 3

In this section we continue the notation and hypotheses of Sect. 3. In addition we assume that  $-1 \in N$  and that  $D^{\times}/N$  supports a V-graph  $\Delta$ . We write  $\Delta$  also for the vertex set of  $\Delta$ , i.e., for the non-identity elements of  $D^{\times}/N$ . The purpose of this section is to prove Theorem B(1) of the Introduction:

**Theorem 4.1.** Assume that there are elements  $x^*$ ,  $y^* \in \Delta$  such that  $d(x^*, y^*) \ge 3$ . Then the map  $\varphi_{y^*} \colon N \to \Gamma_{y^*}$  is a strongly leveled map.

## We start with

- *Remark 4.2.* (1) Note that in Theorem B of the introduction we are assuming that  $-1 \in N$ . Hence for all  $a \in D^{\times}$ ,  $(-a)^* = a^*$ . Also, by property (V1), for every  $x, y \in D^{\times} \setminus N$  and  $n \in N$ , if  $x + y \in N$  or  $x y \in N$ , then  $d(x^*, y^*) \leq 1$ . Similarly, if  $n \notin N(y)$ , then  $d(y^*, (y + n)^*) \leq 1$ . We use these facts without further reference.
- (2) Axiom (V3), in conjunction with axiom (V2), also implies:
  (\*) If d(a\*, c\*) ≤ 1 and d(b\*a\*, c\*) ≤ 1, then d(b\*, c\*) ≤ 1, for all a, b, c ∈ D<sup>×</sup> \ N such that ba ∉ N. Indeed, d((a<sup>-1</sup>)\*, c\*) ≤ 1 and d((a<sup>-1</sup>)\*(b<sup>-1</sup>)\*, c\*) ≤ 1, so by (V3), d((b<sup>-1</sup>)\*, c\*) ≤ 1, whence d(b\*, c\*) ≤ 1.
- (3) Axiom (V3) and (\*) have the following immediate consequence:
- (V3') If  $d(a^*, (ab)^*) \le 2$  or  $d(a^*, (ba)^*) \le 2$ , then  $d(a^*, b^*) \le 2$ , for all  $a, b \in D^{\times} \setminus N$  such that  $ab \notin N$ . In fact, in this section, as well as in Sects. 5 and 6 (which are based on the results of this section) we will need axiom (V3) only in this weaker form (V3'). The full strength of axiom (V3) will be needed only in Sect. 7.
- (4) Notice that for  $a, b \in D \setminus N$ , if  $a^* \neq b^* \neq (a^*)^{-1}$ , then  $d(a^*, b^*) = d((a^*)^{-1}, b^*)$ . Indeed this follows from axiom (V2).

The next two lemmas list some advanced and useful properties of the sets N(x). We mention that Lemma 4.4(1) will be used only in the next section.

**Lemma 4.3.** [27, Lemma 6.8]. *Let*  $x, y \in D^{\times} \setminus N$ , and assume that  $d(x^*, y^*) \ge 3$ , *then*,

(1) 
$$x + y \notin N$$
;  
(2)  $N(x + y) = N(x) \cap N(y)$ ;  
(3) if  $d((x + y)^*, x^*) \ge 3$ , then  $N(x + y) = N(y) \subseteq N(x) \cap N(-x)$ .

*Proof.* The proof of this lemma uses only property (V1) of  $\Delta$ . Part (1) follows from Remark 4.2(1).

(2): Let z := x + y and let  $n \in N(z)$ . Suppose, say, that  $n \notin N(x)$ . We have

$$(x+n) + y = z + n \in N.$$

By Remark 4.2(1),  $d(x^*, (x + n)^*) \le 1 \ge d((x + n)^*, y^*)$ ; thus  $d(x^*, y^*) \le 2$ , a contradiction. This shows that  $N(z) \subseteq N(x)$ . Similarly,  $N(z) \subseteq N(y)$ .

It remains to show that  $N(x) \cap N(y) \subseteq N(z)$ . Assume false and let  $n \in (N(x) \cap N(y)) \setminus N(z)$ . Then  $(z + n) - x = (x + y + n) - x = y + n \in N$ . Similarly,  $(z + n) - y \in N$ , so by Remark 4.2(1),  $x^*$ ,  $(z + n)^*$ ,  $y^*$  is a path in  $\Delta$ , a contradiction.

(3): This follows from (2), since the latter also implies that

$$N(y) = N(x + y - x) = N(x + y) \cap N(-x).$$

**Lemma 4.4.** [27, Lemma 6.9]. Let  $a, b \in D^{\times} \setminus N$  and  $\epsilon \in \{1, -1\}$ . Then

(1) if  $d(a^*, b^*) \ge 4$  and  $\epsilon \notin N(b)$ , then  $N(ab) \cup N(ba) \subseteq N(a) \cap N(-a)$ ; (2) if  $d(a^*, b^*) \ge 3$  and  $\epsilon \in N(b^{-1})$ , then  $N(ab) \cup N(ba) \subseteq N(a) \cap N(-a)$ ; (3) if  $d(a^*, b^*) \ge 3$  and  $\epsilon \in N(a)$ , then  $N(b) \subseteq N(ab) \cap N(ba)$ .

*Proof.* (1) and (2): We first claim that under the hypothesis of either (1) or (2) we have  $b + \epsilon \notin N$  and  $d(a^*, (b + \epsilon)^*) \ge 3$ .

Indeed, in (1) we assume  $d(a^*, b^*) \ge 4$  and  $b + \epsilon \notin N$ . Since  $d(b^*, (b+\epsilon)^*) \le 1$  we get  $d(a^*, (b+\epsilon)^*) \ge 3$ .

In (2) we assume  $d(a^*, b^*) \ge 3$  and  $\epsilon \in N(b^{-1})$ . By Lemma 3.1(4),  $b + \epsilon \notin N$  and  $(b + \epsilon)^* = b^*$ , so again  $d(a^*, (b + \epsilon)^*) \ge 3$ , proving our claim.

From the claim and (V3') we deduce that  $ab + \epsilon a \notin N$  and  $d(a^*, (ab + \epsilon a)^*) \ge$ 3. Also,  $d(a^*, (ab)^*) \ge 3$ , by (V3') again. By Lemma 4.3(3) (with  $\epsilon a, ab$  in place of x, y, respectively),  $N(ab) \subseteq N(\epsilon a) \cap N(-\epsilon a) = N(a) \cap N(-a)$ . For the other inclusion conjugate by a using Lemma 3.1(2).

(3): Since  $d(a^*, b^*) \ge 3$ , (V2) and (V3') imply that  $d((ab)^*, (a^{-1})^*) \ge 3$ , and it follows from (2) (taking  $ab, a^{-1}$  in place of a, b, respectively), that  $N(a^{-1}ab) \subseteq N(ab)$ , that is  $N(b) \subseteq N(ab)$ . The other inclusion is obtained by conjugating by b.

The next two results are based on [28, Propositions 5.2 and 5.3].

**Proposition 4.5.** Let  $x, y \in D^{\times} \setminus N$  satisfy  $d(x^*, y^*) \geq 3$ . Let  $a \in \mathbb{P}_{x^*}$  and  $b \in \mathbb{P}_{y^*}$ . Then:

(1)  $a + 1 \in U_{y^*}$ , and hence  $a + 1 \in N(b)$ ; (2)  $N(a^{-1}) \subseteq N(b)$ ; (3)  $N(a^{-1}b^{-1}) \subseteq N(\epsilon a^{-1}) \cap N(\epsilon b^{-1})$  for  $\epsilon = \pm 1$ .

Further, let  $n \in N$  satisfy  $n^{-1} \in N(a^{-1}b^{-1})$ . Then

(4)  $N(nb) \neq N(b);$ (5)  $(N(a) \cap N(b)) \pm n \subseteq N(a) \cap N(b);$ (6)  $1 \pm n \in N(c)$  for every  $c \in \mathbb{P}_{(x^{-1}y^{-1})^*}.$ 

*Proof.* (1): First,  $a + 1 \in N$ , because  $1 \in N(a)$ . By definition, to show that  $a + 1 \in U_{y^*}$ , it remains to show that N(ay + y) = N(y).

By (V3') and Remark 4.2(2),  $d((ay)^*, y^*) = d(x^*y^*, y^*) \ge 3$ . Now  $(ay + y)^* = y^*$ , since  $a + 1 \in N$ , so  $d((ay + y)^*, (ay)^*) = d(y^*, (xy)^*) \ge 3$ . By Lemma 4.3(3) (with ay, y in place of x, y, respectively), N(ay + y) = N(y).

But now,  $\varphi_{y^*}(a+1) = 0$  since  $a+1 \in U_{y^*}$ , so by Corollary 3.5(1),  $a+1 \in N(b)$ . (2): By (V2),  $d(y^*, (x^{-1})^*) \ge 3$ . By Lemma 4.4(3),(2) (taking  $b, a^{-1}$  in place of a, b, respectively, and  $\epsilon = 1$ ),

$$N(a^{-1}) \subseteq N(a^{-1}b) \subseteq N(b).$$

(3): Since  $d((x^{-1})^*, (y^{-1})^*) \ge 3$ , we may use Lemma 4.4(2), with  $\epsilon a^{-1}, \epsilon b^{-1}$  in place of *a*, *b*, and then with  $\epsilon b^{-1}, \epsilon a^{-1}$  in place of *a*, *b*.

(4): As  $1 \in N(b)$  we have  $n \in N(nb)$ . On the other hand,  $n^{-1} \in N(b^{-1})$ , by (3). By Lemma 3.1(4),  $n \notin N(b)$ . Therefore  $N(nb) \neq N(b)$ .

(5): By (3),  $\pm n^{-1} \in N(a^{-1}) \cap N(b^{-1})$ , so  $a \pm n \in aN$  and  $b \pm n \in bN$ . By Lemma 4.3(2) (with a+n, b-n in place of x, y),  $N(a+b) = N(a+n) \cap N(b-n)$ . Similarly  $N(a+b) = N(a-n) \cap N(b+n)$ . Further, by Lemma 4.3(2),  $N(a) \cap N(b) = N(a+b)$ , so we obtain

$$N(a) \cap N(b) \subseteq N(a + \epsilon n) \cap N(b + \epsilon n),$$

for  $\epsilon = \pm 1$ .

Let now  $m \in N(a) \cap N(b)$ . We obtain that  $a + \epsilon n + m, b + \epsilon n + m \in N$ . Therefore, if  $\epsilon n + m \notin N$ , then by Remark 4.2(1),  $a^*$ ,  $(\epsilon n + m)^*$ ,  $b^*$  is a path in  $\Delta$ , contradicting  $d(a^*, b^*) \geq 3$ . Consequently,  $\epsilon n + m \in N$ , whence  $m + \epsilon n \in N(a) \cap N(b)$ .

(6): By (5),  $1 \pm n \in N$ . By (1) (with  $x^{-1}y^{-1}$ , x, c, a in place of x, y, a, b) we have  $c + 1 \in N(a)$ . Similarly,  $c + 1 \in N(b)$ . It follows from (5) that  $c + 1 \pm n \in N(a) \cap N(b) \subseteq N$ . Hence  $1 \pm n \in N(c)$ .

**Corollary 4.6.** Let  $x, y \in D^{\times} \setminus N$  satisfy  $d(x^*, y^*) \ge 3$ . Then

- (1)  $N(a^{-1}) \subseteq N_{\leq_{y^*}0}$  for every  $a \in \mathbb{P}_{x^*}$ ;
- (2)  $N_{>_{y}*\alpha} \neq \emptyset$ , for all  $\alpha \in \Gamma_{y^*}$ ;
- (3) if D is a finite-dimensional algebra over an infinite field k, then for every  $\alpha \in \Gamma_{v^*}$ , the set  $N_{\geq_{v^*}\alpha}$  contains a basis of D over k.

*Proof.* (1): For every  $b \in \mathbb{P}_{y^*}$  Proposition 4.5(2) gives  $N(a^{-1}) \subseteq N(b)$ . Now use Corollary 3.5(1).

(2): Let *a*, *b* and *n* be as in Proposition 4.5. By Proposition 4.5(3),  $n^{-1} \in N(a^{-1})$ , so by (1),  $n^{-1} \in N_{\leq_{v}*0}$ . It follows that  $n \in N_{\geq_{v}*0}$ .

Also, by Proposition 4.5(4),  $N(nb) \neq N(b)$ , so, by definition,  $n \notin U_{y^*}$ . Hence  $\varphi_{y^*}(n) \neq 0$ . It follows that  $n \in N_{>_{y^*}0}$ . Since  $N_{>_{y^*}0} \neq \emptyset$ , the assertion now follows from the surjectivity of  $\varphi_{y^*}$ .

(3): Pick  $a \in Nx$  so that  $1 \in N(a)$ . Then according to (1) we have  $N(a^{-1}) \subseteq N_{\leq_y * 0}$ , so  $N(a^{-1})^{-1} \subseteq N_{\geq_y * 0}$ . Proposition 3.6 gives a basis of *D* over *k* inside  $N_{\geq_y * 0}$ .

Now take  $z \in N$  with  $\alpha = \varphi_{y^*}(z)$ . We multiply the above basis by z to obtain a basis of D over k in  $zN_{\geq_y*0} = N_{\geq_y*\alpha}$ .

*Proof of Theorem 4.1.* Let  $a \in \mathbb{P}_{x^*}$ ,  $b \in \mathbb{P}_{y^*}$  and  $n \in N$  with  $n^{-1} \in N(a^{-1}b^{-1})$ . Set

$$z := x^{-1}y^{-1}.$$

We claim that  $\varphi_{z^*}$  is a strongly leveled map with level  $\alpha := \varphi_{z^*}(n)$ . We recall that this means that

- (i)  $\alpha \geq_{z^*} 0$  and  $N_{\geq_{z^*}\alpha} \neq \emptyset$ .
- (ii) If  $m \in N$  and  $\varphi_{z^*}(m) >_{z^*} \alpha$ , then  $\varphi_{z^*}(1 \pm m) \leq_{z^*} 0$ .

For (i) we use Corollary 4.6 (but with *x*, *z* in place of *x*, *y*). Indeed by Proposition 4.5(3),  $n^{-1} \in N(a^{-1})$ , so by Corollary 4.6(1),  $\varphi_{z^*}(n^{-1}) \leq_{z^*} 0$ , i.e.,  $\alpha \geq_{z^*} 0$ . By Corollary 4.6(2),  $N_{z^*\alpha} \neq \emptyset$ .

For *m* as in (ii) we have  $\varphi_{z^*}(m^{-1}) < \varphi_{z^*}(n^{-1})$ . By Corollary 3.5(2) (with  $z, a^{-1}b^{-1}, m^{-1}, n^{-1}$  in place of y, y', m, n),  $m^{-1} \in N(a^{-1}b^{-1})$ . Hence, by Proposition 4.5(6),  $1 \pm m \in N(c)$  for every  $c \in \mathbb{P}_{z^*}$ . We now use Corollary 3.5(1).

We may now take  $x^{-1}$ ,  $y^{-1}x$ , y in place of x, y, z to conclude that  $\varphi_{y^*}$  is also a strongly leveled map.

## 5. The case where diam( $\Delta$ ) $\geq 4$

In this section we continue with the notation and hypotheses of Sect. 4. The purpose of this section is to prove Theorem B(2) of the Introduction:

**Theorem 5.1.** Assume that there are elements  $x^*$ ,  $y^* \in \Delta$  such that  $d(x^*, y^*) \ge 4$ . Then (after perhaps interchanging x and y) the map  $\varphi_{y^*} \colon N \to \Gamma_{y^*}$  is a strong valuation-like map.

The following notation should be compared with [28, Notation 5.5, p. 944].

**Notation 5.2.** Let  $M \subseteq N$  be a subgroup, and let  $r, s \in D \setminus N$ .

- (1) We denote  $\dot{N}_M(r) = N(r) \cap M$ . When M is clear from the context we will omit the subscript M and write  $\dot{N}(r)$  in place of  $\dot{N}_M(r)$ . Note that while N(r) is always non-empty,  $\dot{N}_M(r)$  may well be empty. If  $\dot{N}_M(r) = \emptyset$  and  $m \in M$ , then our convention is that  $m\dot{N}_M(r) = \emptyset$ .
- (2) We denote by  $\text{In}_M(r^*, s^*)$  the following relation: for any  $a \in Nr$  and  $b \in Ns$ we have  $\dot{N}_M(a) \subseteq \dot{N}_M(b)$  or  $\dot{N}_M(b) \subseteq \dot{N}_M(a)$ .
- (3) We denote by  $\operatorname{Inc}_M(s^*, r^*)$  the following relation:  $\operatorname{In}_M(r^*, s^*)$  and for any  $b \in \mathbb{P}_{s^*}$  there exists  $a \in \mathbb{P}_{r^*}$  such that  $\dot{N}_M(b) \supseteq \dot{N}_M(a)$ .

*Of course, when* M = N *we have*  $N(r) = \dot{N}_N(r)$ *. In this case we abbreviate* 

 $In(r^*, s^*) = In_N(r^*, s^*), \quad Inc(s^*, r^*) = Inc_N(s^*, r^*)$ 

Note that in Notation 5.2(2) we allow  $r^* = s^*$ . Furthermore we have (compare with [28, Proposition 5.7(2)]):

**Lemma 5.3.** Let  $M \subseteq N$  be a subgroup and let  $s \in D^{\times} \setminus N$ . Assume that  $\operatorname{In}_{M}(s^{*}, s^{*})$  holds. Then the subgroup  $\varphi_{s^{*}}(M) \subseteq \Gamma_{s^{*}}$  is totally ordered. In particular, if M = N, then  $(\Gamma_{s^{*}}, \leq_{s^{*}})$  is a totally ordered group.

*Proof.* Let  $m, m' \in M$ , and suppose that  $\varphi_{s^*}(m) \leq \varphi_{s^*}(m')$  does not hold. By the equivalence (2) $\Leftrightarrow$ (4) in Lemma 3.4, there exists  $b \in \mathbb{P}_{s^*}$  such that  $m \notin N(m'b)$ . Let  $c \in \mathbb{P}_{s^*}$ . Then  $m \in \dot{N}_M(mc) \setminus \dot{N}_M(m'b)$ . As  $mc, m'b \in Ns$ , our assumption  $\ln_M(s^*, s^*)$  implies that  $\dot{N}_M(m'b) \subseteq \dot{N}_M(mc)$ . Therefore  $m' \in \dot{N}_M(m'b) \subseteq \dot{N}_M(mc) \subseteq N(mc)$ . Using again the equivalence (2) $\Leftrightarrow$ (4) in Lemma 3.4, we conclude that  $\varphi_{s^*}(m') \leq \varphi_{s^*}(m)$ . Next we have

**Proposition 5.4.** [27, Proposition 6.11]. *Let*  $x, y \in D^{\times} \setminus N$  *with*  $d(x^*, y^*) \ge 4$ . *Then*  $In(x^*, y^*)$ .

*Proof.* Let  $a \in Nx$  and  $b \in Ny$ . We need to show that  $N(a) \subseteq N(b)$  or  $N(b) \subseteq N(a)$ . So assume  $N(b) \nsubseteq N(a)$ , and let  $n \in N(b) \smallsetminus N(a)$ . Let  $c := n^{-1}a$  and  $d := n^{-1}b$ . Then  $1 \in N(d) \smallsetminus N(c)$ . By Lemma 4.4(3),(1), respectively,  $N(c) \subseteq N(cd) \subseteq N(d)$ . Hence also  $N(a) \subseteq N(b)$ .

The next two lemmas list some properties of the relations  $In_M$  and  $Inc_M$ . They follow [34, 3.10] and [27, Lemma 6.12].

**Lemma 5.5.** Let  $r, s, t \in D^{\times} \setminus N$  and let  $M \subseteq N$  be a subgroup. Then,

- (1) if  $\text{In}_M(r^*, s^*)$ , then  $\text{Inc}_M(r^*, s^*)$  or  $\text{Inc}_M(s^*, r^*)$ ;
- (2)  $\operatorname{Inc}_{M}(s^{*}, r^{*})$  and  $\operatorname{In}_{M}(r^{*}, t^{*})$  imply  $\operatorname{In}_{M}(s^{*}, t^{*})$ ;
- (3)  $\operatorname{Inc}_{M}(s^{*}, r^{*})$  and  $\operatorname{Inc}_{M}(r^{*}, t^{*})$  imply  $\operatorname{Inc}_{M}(s^{*}, t^{*})$ ;
- (4) *if*  $Inc_M(s^*, r^*)$ , *then*  $In_M(s^*, s^*)$ .
- *Proof.* (1): Suppose  $\text{Inc}_M(r^*, s^*)$  does not hold. Then there is  $a \in \mathbb{P}_{r^*}$  such that  $\dot{N}(a) \subseteq \dot{N}(b)$ , for all  $b \in \mathbb{P}_{s^*}$ , so  $\text{Inc}_M(s^*, r^*)$  holds.
- (2): Let  $b \in Ns$  and  $c \in Nt$ . Suppose that  $\dot{N}(b) \nsubseteq \dot{N}(c)$ . We need to show that  $\dot{N}(b) \supseteq \dot{N}(c)$ . Let  $m \in \dot{N}(b) \smallsetminus \dot{N}(c)$ . Then replacing b by  $m^{-1}b$  and c by  $m^{-1}c$ , we may assume that  $1 \in \dot{N}(b) \smallsetminus \dot{N}(c)$  (we note that  $m \in M$ , so  $\dot{N}(m^{-1}d) = m^{-1}\dot{N}(d)$ , for any  $d \in D^{\times} \smallsetminus N$ ). By  $\operatorname{Inc}_{M}(s^{*}, r^{*})$ , we can pick  $a \in \mathbb{P}_{r^{*}}$ , with  $\dot{N}(b) \supseteq \dot{N}(a)$ . As  $1 \in \dot{N}(a) \smallsetminus \dot{N}(c)$ ,  $\operatorname{In}_{M}(r^{*}, t^{*})$  implies that  $\dot{N}(a) \supseteq \dot{N}(c)$ , so  $\dot{N}(b) \supseteq \dot{N}(a) \supseteq \dot{N}(c)$ , as asserted.
- (3): First, by (2) we have  $\operatorname{In}_M(s^*, t^*)$ . Let  $b \in \mathbb{P}_{s^*}$ . By  $\operatorname{Inc}_M(s^*, r^*)$ , there is  $a \in \mathbb{P}_{r^*}$ , with  $\dot{N}(b) \supseteq \dot{N}(a)$ . By  $\operatorname{Inc}_M(r^*, t^*)$ , there is  $c \in \mathbb{P}_{t^*}$ , with  $\dot{N}(a) \supseteq \dot{N}(c)$ . Thus  $\dot{N}(b) \supseteq \dot{N}(a) \supseteq \dot{N}(c)$ , and we get  $\operatorname{Inc}_M(s^*, t^*)$ .
- (4): Since  $\text{Inc}_M(s^*, r^*)$  we have  $\text{In}_M(r^*, s^*)$ . Hence (2) (with t = s) gives  $\text{In}_M(s^*, s^*)$ .

**Lemma 5.6.** Let  $r, s \in D^{\times} \setminus N$  and suppose that  $Inc(s^*, r^*)$ . Then

- (1)  $\leq_{s^*}$  is a total order relation;
- (2)  $N_{\leq_{r^*} 0} \supseteq N_{\leq_{r^*} 0}$ ;
- (3) there is an epimorphism  $\psi \colon \Gamma_{r^*} \to \Gamma_{s^*}$  of partially ordered groups such that  $\varphi_{s^*} = \psi \circ \varphi_{r^*}$ .
- *Proof.* (1): As  $Inc(s^*, r^*)$ , Lemma 5.5(4) gives  $In(s^*, s^*)$ . Now use Lemma 5.3.
- (2): This follows from Corollary 3.5(1). Indeed, let  $b \in \mathbb{P}_{s^*}$ , and (using  $Inc(s^*, r^*)$ ) pick  $a \in \mathbb{P}_{r^*}$  with  $N(b) \supseteq N(a)$ . Then  $N(b) \supseteq N(a) \supseteq N_{\leq_{r^*} 0}$ . As this holds for all  $b \in \mathbb{P}_{s^*}$ ,  $N_{\leq_{s^*} 0} \supseteq N_{\leq_{r^*} 0}$ .
- (3): Recall that for z = r, s, the kernel of the map  $\varphi_{z^*} : N \to \Gamma_{z^*}$  is  $U_{z^*}$ . By (2),  $U_{r^*} \subseteq U_{s^*}$ , and hence there exists an epimorphism  $\psi : \Gamma_{r^*} \to \Gamma_{s^*}$ , such that  $\varphi_{s^*} = \psi \circ \varphi_{r^*}$ , as required. Moreover, by (2) again,  $\psi$  is a homomorphism of partially ordered groups.

*Proof of Theorem 5.1.* Let  $x, y \in D^{\times} \setminus N$  such that  $d(y^*, x^*) \ge 4$ . By Theorem 4.1, both  $\varphi_{x^*}: N \to \Gamma_{x^*}$  and  $\varphi_{y^*}: N \to \Gamma_{y^*}$  are strongly leveled maps. Thus it remains to show that one of the groups  $\Gamma_{x^*}, \Gamma_{y^*}$  is totally ordered.

Now by Proposition 5.4,  $In(x^*, y^*)$ . By Lemma 5.5(1), after perhaps interchanging x and y, we may assume that  $Inc(y^*, x^*)$ . Then by Lemma 5.6(1),  $\Gamma_{y^*}$  is a totally ordered group.

## 6. The case where diam( $\Delta$ ) $\geq$ 5

In this section we continue the notation and hypotheses of Sect. 4. The purpose of this section is to prove Theorem B(3) of the Introduction. We thus assume that  $x^*, y^* \in \Delta$  are such that  $d(x^*, y^*) \ge 5$ . By Proposition 5.4 and Lemma 5.5(1), we may assume without loss of generality that  $Inc(y^*, x^*)$  holds. We will prove

**Theorem 6.1.** The map  $\varphi_{v^*} \colon N \to \Gamma_{v^*}$  is a strong valuation like map of s-level 0.

By Lemma 5.5(4),  $In(y^*, y^*)$  holds, and this together with Lemma 3.1(1) and the fact that  $-1 \in N$  implies that

$$N(g) = N(-g), \text{ for all } g \in Ny.$$
(6.1)

**Lemma 6.2.** Let  $a \in xN$ ,  $b \in yN$  such that  $N(a) \subseteq N(b)$  and let  $n \in N \setminus N(b)$ . Then

$$N(a) \subseteq N(a+n) \cap N(b+n).$$

*Proof.* Since N(b) = N(-b) we have N(b) = -N(b) and  $N(-a) \subseteq N(b)$ .

If  $a-n \in N$ , then  $-n \in N(a) \subseteq N(b) = -N(b)$ , a contradiction. Thus  $a-n \notin N$ , so by Remark 4.2(1),  $d(a^*, (a-n)^*) \leq 1$ . Also,  $b+n \notin N$  and we similarly have  $d((b+n)^*, b^*) \leq 1$ . As  $d(a^*, b^*) \geq 5$  we conclude that  $d((a-n)^*, (b+n)^*) \geq 3$ . Hence, by Lemma 4.3(2),  $N(a+b) = N(a-n) \cap N(b+n)$ . In addition, Lemma 4.3(2) implies that  $N(a+b) = N(a) \cap N(b) = N(a)$ . Therefore

$$N(a) = N(a - n) \cap N(b + n).$$
 (6.2)

The same argument, with -a in place of a, shows that

$$N(-a) = N(-a - n) \cap N(b + n).$$
(6.3)

Altogether Eqs. (6.2) and (6.3) imply the assertion.

*Proof of Theorem 6.1.* We denote  $\leq = \leq_{y^*}$  and  $\varphi = \varphi_{y^*}$ . By Theorem 5.1 it is enough to show that  $\varphi$  is a strongly leveled map having s-level 0, i.e.

$$1 + N_{>0} \subseteq N_{<0} \tag{6.4}$$

(we notice that since  $\Gamma_{y^*}$  is totally ordered, we have  $\pm N_{>0} = N_{>0}$ , hence equation (SL) in subsection 1.2 of the introduction simplifies to Eq. (6.4)).

We first show that

$$1 + N_{>0} \subseteq N. \tag{6.5}$$

To this end let  $n \in N_{>0}$ . It follows from Corollary 3.5(1) that there exists  $b \in \mathbb{P}_{y^*}$  such that  $n \notin N(b)$ . Using  $Inc(y^*, x^*)$  we pick  $a \in \mathbb{P}_{x^*}$  such that  $N(b) \supseteq N(a)$ . Since  $1 \in N(a)$ , Lemma 6.2 shows that both a + n + 1 and b + n + 1 belong to N. If  $n + 1 \notin N$ , then  $a^*$ ,  $(n + 1)^*$ ,  $b^*$  would be a path in  $\Delta$ , contrary to  $d(a^*, b^*) \ge 5$ . Thus  $n + 1 \in N$  proving Eq. (6.5).

Next we note that  $1 - (1 + a) = -a \notin N$ . By eq. (6.5),  $-(1 + a) \in N_{\leq 0}$ , whence also  $1 + a \in N_{\leq 0}$  (see Eq. (6.1)).

Corollary 3.5(1) now implies that  $1 + a \in N(z)$  for every  $z \in \mathbb{P}_{y^*}$ . Equivalently,  $z + 1 \in N(a)$  for every such z. By Lemma 6.2,

$$a + (z + 1 + n), b + (z + 1 + n) \in N.$$

The assumption  $z + 1 + n \notin N$  would lead us again to the false conclusion that  $d(a^*, b^*) \leq 2$ . Thus,  $z + (1 + n) \in N$ . Since this is true for all  $z \in \mathbb{P}_{y^*}$ , we see that  $1 + n \in N_{\leq 0}$ , by Corollary 3.5(1), proving Eq. (6.4).

## 7. Property $(3\frac{1}{2})$

In this section we continue the notation and hypotheses of Sect. 4. Recall that we denote by *F* the center of *D*. Throughout this section  $\Sigma \leq \text{Sym}(D^{\times})$  is a permutation group on  $D^{\times}$ . For  $x \in D^{\times}$  and  $\sigma \in \Sigma$ , let  $\sigma(xN) := \{\sigma(xn) \mid n \in N\}$ . We assume that  $\sigma(N) = N$ , for all  $\sigma \in \Sigma$ , and

$$\sigma(xN) = \sigma(x)N$$
, for all  $x \in D^{\times}$  and  $\sigma \in \Sigma$ ,

that is  $\sigma(x^*) = \sigma(x)^*$ , for all  $x \in D^{\times}$  and  $\sigma \in \Sigma$ . We use the letter  $\Sigma$  to also denote the group of permutations of  $\Delta$  induced by  $\Sigma$ .

*Remark 7.1.* Note that if  $\Sigma \leq \operatorname{Aut}(D^{\times})$  is a subgroup that normalizes N, then of course  $\Sigma$  satisfies our hypothesis.

**Definition 7.2.** We say that  $\Delta$  satisfies *Property*  $(3\frac{1}{2})$  with respect to  $x^*$ ,  $y^* \in \Delta$ , the subgroup  $\Sigma \leq \text{Sym}(D^{\times})$ , and the subgroup  $M \subseteq N$ , if  $-1 \in M$  and

- (1)  $\sigma(a+k)^* = (\sigma(a)+k)^*$ , for every  $\sigma \in \Sigma$ ,  $a \in D^{\times} \setminus N$  and  $k \in M$ .
- (2)  $d(x^*, y^*) \ge 3$ .
- (3) For *every*  $r^*$ ,  $s^* \in \Delta$  such that  $x^*$ ,  $r^*$ ,  $s^*$ ,  $y^*$  is a path in  $\Delta$ , there exists  $\sigma \in \Sigma$  such that  $d(\sigma(x^*), y^*) \ge 3$ , and  $\sigma(x)^*$ ,  $\sigma(r)^*$ ,  $s^*$ ,  $y^*$ , is *not* a path in  $\Delta$ .

The purpose of this section is to prove:

**Theorem 7.3.** Assume that  $\Delta$  satisfies property  $(3\frac{1}{2})$  with respect to  $x^*, y^* \in \Delta$ , the subgroup  $\Sigma \leq \text{Sym}(D^{\times})$ , and the subgroup  $M \subseteq N$ . Then, after perhaps interchanging  $x^*$  and  $y^*$ , the map  $\varphi_{y^*} \colon N \to \Gamma_{y^*}$  is a strongly leveled map such that the subgroup  $\varphi_{y^*}(M)$  of  $\Gamma_{y^*}$  is totally ordered. **Lemma 7.4.** Suppose that  $\Delta$  satisfies property  $(3\frac{1}{2})$  with respect to  $x^*$ ,  $y^* \in \Delta$ ,  $\Sigma$  and M. Then  $\ln_M(x^*, y^*)$  holds.

*Proof.* Suppose that  $\ln_M(x^*, y^*)$  does not hold. Then there exist  $a \in xN$  and  $b \in yN$  such that  $\dot{N}(a) \not\subseteq \dot{N}(b)$  and  $\dot{N}(b) \not\subseteq \dot{N}(a)$ . Pick  $m \in \dot{N}(a) \setminus \dot{N}(b)$ . We may replace a, b by  $-m^{-1}a, -m^{-1}b$  respectively to assume that

$$-1 \in \dot{N}(a) \smallsetminus \dot{N}(b).$$

Also pick

$$k \in \dot{N}(b) \smallsetminus \dot{N}(a),$$

and set r = a + k and s = b - 1. Then  $r, s \notin N$ .

Take  $\sigma \in \Sigma$  such that  $d(\sigma(x)^*, y^*) \ge 3$ . We show that

$$\sigma(x)^*, \sigma(r)^*, s^*, y^* \tag{P}$$

is a path in  $\Delta$ . In particular, applying this for  $\sigma = \text{id}$ , we obtain from (2) of Definition 7.2 that  $x^*, r^*, s^*, y^*$  is a path in  $\Delta$ , which is a contradiction to (3) of Definition 7.2.

Now since  $d(\sigma(x)^*, y^*) \ge 3$ , it is enough to show that in each step in  $\mathcal{P}$  the distance is *at most* 1.

Indeed, by (1) of Definition 7.2,  $\sigma(r)^* = \sigma(a+k)^* = (\sigma(a)+k)^*$ . Since  $k \in M \subseteq N$ , (V1) implies that

$$d(\sigma(x)^*, \sigma(r)^*) = d(\sigma(a)^*, (\sigma(a) + k)^*) \le 1.$$

Next, we have  $a - 1 \in N$ , which by (1) of Definition 7.2 implies that  $(\sigma(a) - 1)^* = \sigma(a - 1)^* = 1^*$ . Therefore  $-1 \in N(\sigma(a))$ . Since  $d(\sigma(a)^*, b^*) = d(\sigma(x)^*, y^*) \ge 3$ , we deduce from Lemma 4.4(3) that  $k \in N(b) \subseteq N(b\sigma(a))$ , and hence  $b\sigma(a) + k \in N$ . Therefore, by (V1),

$$d(b^*(\sigma(a)+k)^*, (b-1)^*) = d((b\sigma(a)+k) + (b-1)k)^*, ((b-1)k)^*) \le 1.$$

In addition  $d(b^*, (b-1)^*) \le 1$ , so by (V3),

$$d(\sigma(r)^*, s^*) = d((\sigma(a) + k)^*, (b-1)^*) \le 1.$$

Finally, (V1) implies that

$$d(s^*, y^*) = d((b-1)^*, b^*) \le 1,$$

as required.

Proof of Theorem 7.3. Since  $d(x^*, y^*) \ge 3$ , Theorem 4.1 implies that both  $\varphi_{y^*}$  and  $\varphi_{x^*}$  are strongly leveled maps. By Lemma 7.4,  $\ln_M(x^*, y^*)$  holds. In view of Lemma 5.5(1), we may assume (after perhaps interchanging  $x^*$  and  $y^*$ ) that  $\operatorname{Inc}_M(y^*, x^*)$  holds. Then Lemma 5.5(4) shows that  $\operatorname{In}_M(y^*, y^*)$  also holds. By Lemma 5.3, the subgroup  $\varphi_{y^*}(M)$  of  $\Gamma_{y^*}$  is totally ordered.

Using Theorem 7.3 we will prove at the end of Sect. 10 (compare with [28, Theorem 1, p. 931]):

**Theorem 7.5.** Let D be a finite-dimensional separable (but not necessarily central) division algebra over an infinite field k of finite transcendence degree over its prime field, and let  $N \subseteq D^{\times}$  be a normal subgroup of finite index containing -1. Assume that  $D^{\times}/N$  supports a V-graph  $\Delta$  that satisfies property  $(3\frac{1}{2})$  with respect to  $x^*$ ,  $y^* \in \Delta$ , the subgroup  $\Sigma \leq \text{Sym}(D^{\times})$ , and the subgroup  $M = N \cap k^{\times}$ .

Then there exists a non-empty finite set  $\tilde{T}$  of non-trivial valuations of D such that N is open in  $D^{\times}$  with respect to the  $\tilde{T}$ -adic topology.

#### 8. Valuations on division algebras

In this section we recall some notions and facts on valuations on division algebras, which will be needed in the next two sections.

We recall that a valuation on a division ring D is a *surjective* group homomorphism  $v: D^{\times} \to \Gamma_v$  onto a totally ordered group  $\Gamma_v$  (the *value group*) such that

$$v(x + y) \ge \min\{v(x), v(y)\}$$
 whenever  $x + y \ne 0$ .

We let

$$\mathcal{O}_{D,v} = \{x \in D^{\times} \mid v(x) \ge 0\} \cup \{0\}$$

denote the corresponding valuation ring, and  $\mathfrak{m}_{D,v}$  its maximal ideal. More generally, for  $\delta \in (\Gamma_v)_{\geq 0}$ , we define the following two-sided ideal of  $\mathcal{O}_{D,v}$ :

$$\mathfrak{m}_{D,v}(\delta) = \{ x \in D^{\times} \mid v(x) > \delta \} \cup \{ 0 \}$$

(so that  $\mathfrak{m}_{D,v} = \mathfrak{m}_{D,v}(0)$ ).

Recall that the *v*-adic topology on *D* is the ring topology which has the ideals  $\mathfrak{m}_{D,v}(\delta)$  for  $\delta \in (\Gamma_v)_{\geq 0}$  as a fundamental system of neighborhoods of zero (see Sect. 5, no. 1 in [7, Ch. 6]). This topology turns  $D^{\times}$  into a topological group, and the openness of a subgroup  $N \subseteq D^{\times}$  in the *v*-adic topology is equivalent to the existence of  $\delta \in (\Gamma_v)_{\geq 0}$  such that  $1 + \mathfrak{m}_{D,v}(\delta) \subseteq N$ .

More generally, given a finite set  $T = \{v_1, \ldots, v_r\}$  of valuations of D, and  $\delta_i \in (\Gamma_{v_i})_{\geq 0}, i = 1, \ldots, r$ , we define

$$\mathfrak{m}_{D,T}(\delta_1,\ldots,\delta_r)=\bigcap_{i=1}^r\mathfrak{m}_{D,v_i}(\delta_i).$$

Clearly,  $\mathfrak{m}_{D,T}(\delta_1, \ldots, \delta_r)$  is a two-sided ideal of  $\mathcal{O}_{D,T} = \bigcap_{v \in T} \mathcal{O}_{D,v}$ . The ideals  $\mathfrak{m}_{D,T}(\delta_1, \ldots, \delta_r)$  form a fundamental system of neighborhoods of zero for a topology on D compatible with the ring structure; this topology will be called T-adic. Thus, a subgroup  $N \subseteq D^{\times}$  (resp. a subring  $R \subseteq D$ ) is T-adically open iff it contains the congruence subgroup  $1 + \mathfrak{m}_{D,T}(\delta_1, \ldots, \delta_r)$  (resp. the ideal  $\mathfrak{m}_{D,T}(\delta_1, \ldots, \delta_r)$ ) for some  $\delta_i \in (\Gamma_{v_i})_{\geq 0}, i = 1, \ldots, r$ .

Next let  $\Gamma = (\Gamma, \leq)$  be a totally ordered commutative group. We recall that the *height* (also called *rank*) of  $\Gamma$  is the supremum of all non-negative integers *r* such that there exist epimorphisms

$$\Gamma = \Gamma_0 \xrightarrow{\mu_1} \Gamma_1 \xrightarrow{\mu_2} \cdots \rightarrow \Gamma_{r-1} \xrightarrow{\mu_r} \Gamma_r = \{0\}$$

of totally ordered groups, where  $\mu_1, \ldots, \mu_r$  have non-trivial kernels. The group  $\Gamma$  has height  $\leq 1$  if and only if it embeds in the ordered additive group of  $\mathbb{R}$  [11, Th. 2.5.2]. We note that if  $\Gamma$  is commutative and has finite height, then there is an epimorphism  $\mu: \Gamma \to \overline{\Gamma}$ , with  $\overline{\Gamma}$  of height one.

The *height* of a valuation v is defined to be the height of its value group  $\Gamma_v$  (cf. [7, ch. 6, Sect. 4, no. 4] or [11, Sect. 2.2] for a discussion on the height of a totally ordered group/valuation).

The following definition describes a useful connection between leveled maps and valuations.

**Definition 8.1.** Let *N* be a subgroup of  $D^{\times}$ ,  $\Gamma$  a partially ordered group,  $\varphi \colon N \to \Gamma$  a group homomorphism, and  $v \colon D^{\times} \to \Gamma_v$  a valuation. We say that *v* is *associated* with  $\varphi$  if  $\varphi(n) \ge 0$  implies  $v(n) \ge 0$ , for all  $n \in N$ .

Remark 8.2. [27, Remarks 2.5].

(1) Given a non-trivial homomorphism  $\varphi \colon N \to \Gamma$ , the non-trivial valuation  $v \colon D^{\times} \to \Gamma_{v}$  is associated with  $\varphi$  if and only if there exists a non-trivial homomorphism  $\theta \colon \varphi(N) \to \Gamma_{v}$  of ordered groups such that the square



in which  $\iota$  is the inclusion map, commutes. In fact, this was the original definition used in [25,27,28].

(2) If v: D<sup>×</sup> → Γ<sub>v</sub> is a valuation and μ: Γ<sub>v</sub> → Γ is a surjective homomorphism of totally ordered groups, then v
 = μ ∘ v: D<sup>×</sup> → Γ is also a valuation. Also, if v is associated with φ: N → Γ with respect to θ, then v
 is also associated with φ, with respect to μ ∘ θ. Further, every v-adically open subgroup of D<sup>×</sup> is also v
 is also v
 adically open.

Given valuations u, u' on D with value groups  $\Gamma_u, \Gamma_{u'}$ , respectively, we say that u' is *coarser* than u if there is an epimorphism of totally ordered groups  $\mu \colon \Gamma_u \to \Gamma_{u'}$  such that  $u' = u \circ \mu$ .

**Lemma 8.3.** Let D be a division ring which is finite-dimensional over its center. Let  $r \ge 2$  and let  $w_1, \ldots, w_r$  be distinct valuations of height 1 on D. Let  $N_1, \ldots, N_r$  be proper subgroups of  $D^{\times}$  which are open in the  $w_i$ -adic topologies,  $i = 1, 2, \ldots, r$ , respectively. Let  $N = N_1 \cap \cdots \cap N_r$ . Then N is not u-adically open for any nontrivial valuation u on D.

**Proof.** Assume the contrary. The assumption on D implies that  $\Gamma_u$  is commutative (see [18, p. 628]). The set of valuations on D which are coarser than u is linearly ordered with respect to the coarsening relation (see e.g., [11, Prop. 2.1.3 and Prop. 2.2.1]). Since  $w_1, \ldots, w_r$  have height 1 and are distinct, none of them is coarser than the other. Therefore at most one of them can be coarser than u. Without loss of generality  $w_1$  is not coarser than u. Since it has height 1, there is no nontrivial common coarsening of  $w_1$  and u, i.e., they are independent valuations.

As  $N_1$  contains N, it is open in both the  $w_1$ -adic topology and the *u*-adic topology. Furthermore, for every  $d \in D^{\times}$  the coset  $dN_1$  is *u*-adically open in  $D^{\times}$ . By the weak approximation theorem for independent valuations [21],  $N_1 \cap dN_1 \neq \emptyset$ , so  $d \in N_1$ . This contradicts the assumption that  $N_1 \neq D^{\times}$ .

## 9. Valuations from strongly leveled maps

The goal of this section is to show that strongly leveled maps, under some additional assumptions, give rise to valuations. The main result of this section is the following theorem. We refer the reader to Sect. 1.2 of the Introduction for the notion of a *leveled*, *strongly leveled*, and *strong valuation-like* map.

**Theorem 9.1.** Let *D* be a finite-dimensional (but not necessarily central) division algebra over a field k of finite transcendence degree over its prime field, and let  $N \subseteq D^{\times}$  be a normal subgroup containing -1 of finite index. Let  $\varphi \colon N \to \Gamma$  be a strongly leveled map, where  $(\Gamma, \leq)$  is a partially ordered group, such that the subgroup  $\varphi(N \cap k^{\times})$  of  $\Gamma$  is totally ordered. Then:

- (a) the restriction  $\varphi_k = \varphi \upharpoonright_{N \cap k^{\times}}$  is a strong valuation-like map;
- (b) there exists a height one valuation v of k associated with φ<sub>k</sub> such that N ∩ k<sup>×</sup> is open in k<sup>×</sup> in the v-adic topology.

Let us first make a connection between a strongly leveled and a leveled map.

**Lemma 9.2.** Let D be an infinite division ring, let  $N \subseteq D^{\times}$  be a finite index normal subgroup containing -1, and let  $\varphi \colon N \to \Gamma$  be a strongly leveled map of s-level  $\alpha \in \Gamma_{\geq 0}$ . Then

(1) if  $\beta \in \Gamma_{\geq \alpha}$  is such that  $N_{>\beta} \neq \emptyset$ , then  $\beta$  is an s-level of  $\varphi$ ; (2)  $\varphi$  is a leveled map of level  $\alpha$ .

*Proof.* Part (1) follows immediately from the definitions. For part (2), note that  $N_{<-\alpha} \neq \emptyset$  since  $N_{>\alpha} \neq \emptyset$ . Let  $n \in N_{<-\alpha}$ . Then  $1 + n^{-1} \in 1 + N_{>\alpha} \subseteq N_{\leq 0}$ , and therefore

$$\varphi(1+n) = \varphi(n(1+n^{-1})) \le \varphi(n) < -\alpha,$$

i.e.,  $1 + n \in N_{<-\alpha}$ . Thus,  $\varphi$  is a leveled map of level  $\alpha$ .

We record the following two results from [25] and [28], respectively. Here D is a division ring and N a normal subgroup of  $D^{\times}$  containing -1.

**Proposition 9.3.** [25, Proposition 3]. Let  $\varphi: N \to \Gamma$  be a leveled map, and T a finite set of valuations of D associated with  $\varphi$ . Assume that there is a T-adically open subring  $\mathcal{M}$  of D such that

$$\mathcal{M} \cap N \subseteq N_{>-\beta}$$

for some  $\beta \in \Gamma_{>0}$ . Then N is T-adically open in  $D^{\times}$ .

**Lemma 9.4.** [28, Proposition 4.2]. Let  $\varphi: N \to \Gamma$  be a strongly leveled map with *s*-level  $\alpha$ . Let  $\mathcal{A}$  (resp.,  $\mathcal{R}$ ) be the subring of D generated by  $N_{>\alpha}$  (resp.,  $N_{\geq 0}$ ). Then  $-1 \notin \mathcal{A}$ , and for every  $m \in \mathcal{A} \cap N_{>0}$ , the element  $\varphi(m)$  is an *s*-level of  $\varphi$ .

Note that  $\mathcal{R}$  (resp.,  $\mathcal{A}$ ) coincides with the set of all elements of the form  $\epsilon_1 a_1 + \cdots + \epsilon_l a_l$  with  $\epsilon_i = \pm 1$  and  $a_i \in N_{\geq 0}$  (resp.,  $a_i \in N_{>\alpha}$ ), and that  $\mathcal{A}$  is in fact a ring without identity.

The following result establishes the existence of a valuation associated to a given strongly leveled map in the simplest case where the division algebra is assumed to be commutative and the map  $\varphi$  to be strong valuation-like.

**Theorem 9.5.** (Commutative case; see [27, Theorem 4.1]). Let *K* be a field, and let  $N \subseteq K^{\times}$  be a subgroup of finite index containing -1. Assume that  $\varphi: N \to \Gamma$  is a strong valuation-like map. Then

- (1) there exists a non-trivial valuation v on K associated with  $\varphi$  such that N is open in the v-adic topology;
- (2) additionally, if *K* has finite transcendence degree over its prime field, then *v* can be taken to have height one;
- (3) the subring  $\mathcal{R}$  of K generated by  $N_{\geq 0}$  is v-adically open.

*Proof.* Let  $\alpha$  be an s-level of  $\varphi$ , and let  $\mathcal{R}$  and  $\mathcal{A}$  be the subrings introduced in Lemma 9.4 for K in place of D. Let  $\widetilde{R}$  be the integral closure of  $\mathcal{R}$  in K.

We claim that  $\widehat{\mathcal{R}}$  is a valuation ring. Indeed, given any  $x \in K^{\times}$ , for  $m = [K^{\times} : N]$ , we have  $x^m \in N$ , hence either  $x^m$  or  $x^{-m}$  is in  $N_{\geq 0} \subseteq \mathcal{R}$ . Then, respectively, either x or  $x^{-1}$  is in  $\widehat{\mathcal{R}}$ .

Next, we show that

$$\mathcal{R} \cap N_{<-\alpha} = \emptyset. \tag{9.1}$$

Since  $N_{<-\alpha} \neq \emptyset$ , this will also show that the valuation ring  $\widetilde{\mathcal{R}}$  is a proper subring of *K*. Suppose that  $z \in \widetilde{\mathcal{R}} \cap N_{<-\alpha}$ . Then *z* satisfies a polynomial equation

$$z^{d} + a_{1}z^{d-1} + \dots + a_{d} = 0$$

with  $a_i \in \mathcal{R}$ . Since  $z^{-1} \in N_{>\alpha}$ , it follows that

$$-z = a_1 + \dots + a_d z^{-(d-1)} \in \mathcal{R}.$$

On the other hand, the inclusion  $N_{\geq 0}N_{>\alpha} \subseteq N_{>\alpha}$  implies that  $\mathcal{RA} \subseteq \mathcal{A}$ , i.e.  $\mathcal{A}$  is an ideal of  $\mathcal{R}$ . Since  $z^{-1} \in N_{>\alpha} \subseteq \mathcal{A}$ , we obtain  $-1 = -z \cdot z^{-1} \in \mathcal{A}$ , contrary to Lemma 9.4.

Let v be the valuation of K corresponding to the valuation ring  $\widetilde{\mathcal{R}}$ , i.e.,  $x \in \widetilde{\mathcal{R}}$  if and only if  $v(x) \ge 0$ , for  $x \in K^{\times}$  (cf. [7,11]). Since  $N_{\ge 0} \subseteq \mathcal{R} \subseteq \widetilde{\mathcal{R}}$ , for

any  $x \in N_{\geq 0}$  we have  $v(x) \geq 0$ , which implies that v is associated with  $\varphi$ . By construction,  $\tilde{\mathcal{R}}$  is *v*-adically open in *K*. Equation (9.1) implies that for any  $\beta > \alpha$  we have

$$\widetilde{\mathcal{R}} \cap N \subseteq N_{>-\beta},$$

so the *v*-adic openness of N in  $K^{\times}$  follows from Proposition 9.3. This proves (1).

For (2), suppose that *K* has finite transcendence degree over it prime field. It follows from Cor. 1 in no. 3 and Prop. 3 in no. 2 of [7, Ch. 6, Sect. 10], that the (commutative) value group  $\Gamma_v = v(K^{\times})$  has finite height. As we have observed in Sect. 8, there is an epimorphism  $\mu \colon \Gamma_v \to \overline{\Gamma}$  of totally ordered groups, with  $\overline{\Gamma}$  of height one. By Remark 8.2(2),  $\overline{v} = \mu \circ v \colon K^{\times} \to \overline{\Gamma}$  is a valuation of height one which is associated with  $\varphi$ , and since *N* is *v*-adically open, it is also  $\overline{v}$ -adically open.

Finally, we prove (3). By (1), there exists a non-negative  $\delta \in \Gamma_v = v(K^{\times})$  such that  $1 + \mathfrak{m}_{K,v}(\delta) \subseteq N$ . Pick any  $c \in N$  with v(c) > 0. Then for any

$$x \in c(1 + \mathfrak{m}_{K,v}(\delta))$$

we have v(x) > 0. Since  $\varphi(N)$  is totally ordered and v is associated with  $\varphi$ , this implies that  $\varphi(x) > 0$ . Thus

$$c(1 + \mathfrak{m}_{K,v}(\delta)) \subseteq N_{>0} \subseteq \mathcal{R}.$$

Setting  $\delta' = v(c) + \delta$  we obtain from this and from  $c \in N_{>0} \subseteq \mathcal{R}$  that

$$\mathfrak{m}_{K,v}(\delta') = c\mathfrak{m}_{K,v}(\delta) \subseteq \mathcal{R},$$

proving that the ring  $\mathcal{R}$  is *v*-adically open in *K*.

**Lemma 9.6.** Let D be a division ring and let N be a normal subgroup of  $D^{\times}$  of finite index. Let  $\Gamma$  be a totally ordered group and  $\varphi \colon N \to \Gamma$  a homomorphism. Let  $\mathcal{R}$  be the subring of D generated by  $N_{\geq 0}$ . Then any  $x \in D$  can be written in the form  $x = ab^{-1}$  with  $a \in \mathcal{R}$  and  $b \in N_{\geq 0}$ .

*Proof.* By [38, Theorem 1, p. 377], we can write  $x = n_1 - n_2$  with  $n_1, n_2 \in N$ . Suppose first that  $\varphi(n_1) \ge \varphi(n_2)$ . When  $\varphi(n_2) \ge 0$ , we take a = x and b = 1.

When  $\varphi(n_2) < 0$  we take  $a = n_1 n_2^{-1} - 1$  and  $b = n_2^{-1}$ .

The case  $\varphi(n_1) \leq \varphi(n_2)$  is proved similarly.

*Proof of Theorem 9.1.* In view of Theorem 9.5 (with  $k, N \cap k^{\times}$  in place of K, N), it suffices to prove (a), that is, that  $\varphi_k$  is a strongly leveled map. For this, we let  $\alpha \in \Gamma_{\geq 0}$  be an s-level for  $\varphi$ . Let  $\mathcal{A}$  denote the subring of D generated by  $N_{>\alpha}$ , and let  $\mathcal{R}_0$  be the subring of k generated by  $(N \cap k^{\times})_{\geq 0}$ . Pick an arbitrary  $s \in N_{>\alpha}$ , and let  $p(t) = c_{\ell} x^{\ell} + c_{\ell-1} x^{\ell-1} + \cdots + c_0$  be a minimal polynomial of s over k. By Lemma 9.6 (for the subgroup  $N \cap k^{\times}$  of  $k^{\times}$ ), we can write  $c_i = a_i b_i^{-1}$  with  $a_i \in \mathcal{R}_0$  and  $b_i \in (N \cap k^{\times})_{\geq 0}$ ,  $i = 0, 1, \ldots, \ell$ . Multiplying by  $b_0 \cdots b_{\ell}$ , we may therefore assume that  $c_i \in \mathcal{R}_0$  for every i. Of course,  $c_0 \neq 0$ , and we have

$$c_0 = -(c_\ell s^\ell + \dots + c_1 s) \in \mathcal{A}.$$

Then for  $d = [D^{\times} : N]$  we have

$$b := (c_0)^d \in \mathcal{A} \cap N \cap k^{\times}.$$

We claim that  $b \in N_{>0}$ . Indeed, since the group  $\varphi(N \cap k^{\times})$  is totally ordered, we would otherwise have  $b^{-1} \in N_{\geq 0}$ . As before, the inclusion  $N_{\geq 0}N_{>\alpha} \subseteq N_{>\alpha}$ implies that  $\mathcal{R}_0\mathcal{A} \subseteq \mathcal{A}$ . We obtain that  $-1 = b^{-1}(-b) \in \mathcal{A}$ , which contradicts the first part of Lemma 9.4.

Thus,  $b \in \mathcal{A} \cap N_{>0}$ . By the second part of Lemma 9.4,  $\beta = \varphi(b)$  is an s-level for  $\varphi$ . It is therefore also an s-level for  $\varphi_k$ .

#### 10. Openness with respect to finitely many valuations

Throughout this section we consider a finite-dimensional division algebra D over a field k. We assume that D is *separable* over k, i.e., the center F = Z(D) of Dis a separable algebraic extension of k. We further assume that k is equipped with a height one valuation v. Since D is finite-dimensional over k, the field extension F/k is finite, and there are at most [F : k] extensions of v to F (see [7, Ch. 6, Sect. 8, no. 3, Th. 1]). Each such extension is also of height one [11, Cor. 14.2.3(c)].

Let  $| v_v|_v$  denote the absolute value on *k* associated with *v*. Given a basis  $\omega_1, \ldots, \omega_m$  of *D* over *k*, we define a norm on *D* by

$$||a_1\omega_1 + \dots + a_m\omega_m||_{\nu} := \max_{i=1,\dots,m} |a_i|_{\nu}.$$
 (10.1)

This norm turns *D* into a normed vector space over  $(k, ||_v)$ . Let  $\tau_v$  denote the induced topology on *D*. One easily verifies that  $\tau_v$  and the notion of boundedness on *D* associated with  $|| ||_v$  do not depend on the choice of the basis.

Next let *S* be a non-empty finite set of height one valuations on F = Z(D). Let  $\kappa_1, \ldots, \kappa_{n^2}$  be a fixed basis of *D* over *F*. For  $w \in S$  we may define a norm  $|| ||_w$  on *D* with respect to this basis in a way similar to (10.1). Then

$$D_w = D \otimes_F F_w,$$

is a finite-dimensional algebra over the completion  $F_w$  of F with respect to  $||_w$ . The norm  $|| ||_w$  extends to  $D_w$ , it is defined exactly as in (10.1) using the basis  $\omega_1 \otimes 1, \ldots, \omega_m \otimes 1$  of  $D_w$  over  $F_w$ . We endow  $D_w$  with the corresponding topology and the notion of boundedness. Now set

$$D_S = \prod_{w \in S} D_w,$$

and endow it with the product topology  $\tau_S$ . We have a diagonal embedding  $\iota_S : D \to D_S$ . The topology on D induced from  $\tau_S$  via  $\iota_S$  is then the  $\tau_S$ -topology on D. It restricts to the S-adic topology on F. Let  $\operatorname{pr}_w : D_S \to D_w$ , for  $w \in S$ , and  $\operatorname{pr}_T : D_S \to D_T$ , for  $T \subseteq S$ , be the projection maps (with the usual convention that  $D_T$  is a singleton if T is empty).

We will need the following generalization of [25, Lemma 2]:

**Lemma 10.1.** In the above setup, let  $\mathcal{B}$  be a  $\tau_S$ -open subring of  $D_S$ . Let  $S_0$  be a subset of S such that  $\operatorname{pr}_w(\mathcal{B})$  is unbounded for every  $w \in S_0$ , and set  $T := S \setminus S_0$ . Then

$$\mathcal{B} = \mathrm{pr}_T(\mathcal{B}) \times D_{S_0}.$$

In particular, if  $T = \emptyset$  then  $\mathcal{B} = D_S = D_{S_0}$ .

*Proof.* When  $|S_0| = 1$  this is proved (in an equivalent form) in [25, Lemma 2]. The general case follows by induction.

We will also need the following fact from [28]:

**Proposition 10.2.** [28, Theorem 2.4]. Let *D* be a finite-dimensional central division algebra over a field *F*, and let *w* be a height one valuation of *F*. Assume that there exists a subring  $\mathcal{B} \subsetneq D$  such that

- (i)  $\mathcal{B}$  is open in D with respect to the topology defined by the norm  $|| ||_w$ ;
- (ii) there exists a positive integer  $\ell$  such that  $d\mathcal{B}d^{-1} \subseteq \mathcal{B}$  for all  $d \in (D^{\times})^{\ell} := \{x^{\ell} \mid x \in D^{\times}\}.$

Then w extends uniquely to a height one valuation  $\widetilde{w}$  of D such that  $\mathcal{B} \subseteq \mathcal{O}_{D,\widetilde{w}}$ .

**Theorem 10.3.** Let *D* be a finite-dimensional separable division algebra over a field *k* of finite transcendence degree over its prime field, and let F = Z(D) be its center. Let  $N \subseteq D^{\times}$  be a normal subgroup containing -1 and of finite index. Assume that  $\varphi \colon N \to \Gamma$  is a strongly leveled map such that

- (i)  $\varphi(N \cap k^{\times})$  is totally ordered.
- (ii)  $N_{\geq 0}$  contains a basis of D over k.
- (iii) for the subring  $\mathcal{R}$  of D generated by  $N_{\geq 0}$ , there exists  $\gamma \in \Gamma_{>0}$  such that  $\mathcal{R} \cap N \subseteq N_{>-\gamma}$ .

Then

- (1) the restriction  $\varphi_k = \varphi \upharpoonright_{N \cap k^{\times}}$  is a strong valuation-like map;
- (2) there exists a height one valuation v of k associated with φ<sub>k</sub> such that N ∩ k<sup>×</sup> is open in k<sup>×</sup> in the v-adic topology;
- (3) there exists a finite non-empty set T of valuations on F extending v such that  $|T| \leq [F:k]$ , and such that
  - (a) each  $w \in T$  uniquely extends to a valuation  $\tilde{w}$  of D associated with  $\varphi$ ;
  - (b) N is open in  $D^{\times}$  in the  $\tilde{T}$ -adic topology, where  $\tilde{T} = {\tilde{w} \mid w \in T}$ .

*Proof.* Parts (1) and (2) hold by Theorem 9.1, using hypothesis (i).

Let *S* be the set of all extensions of *v* to *F*. Then *S* is non-empty, and as already noted,  $|S| \leq [F:k] < \infty$ . Let  $w_1, \ldots, w_r$  be the distinct valuations in *S*. Let  $k_v$  be the completion of *k* with respect to *v*. Since F/k is separable, we have

$$F \otimes_k k_v \cong \prod_{i=1}^r F_{w_i}$$

(see [7, Ch. 6, Sect. 8, no. 2, Cor. 2]), and therefore

$$D \otimes_k k_v = D \otimes_F (F \otimes_k k_v) \cong D \otimes_F \left(\prod_{i=1}^r F_{w_i}\right) = \prod_{i=1}^r D_{w_i} = D_S.$$

This isomorphism commutes with the natural map  $D \to D \otimes_k k_v$ ,  $d \mapsto d \otimes 1$ , and the diagonal map  $\iota_S \colon D \to D_S$ . We extend the norm  $|| ||_v$  on D to  $D \otimes_k k_v$ . Since any two norms on a finite-dimensional vector space over a complete normed field are equivalent (see e.g., [20, p. 470, Prop. 2.2]), the  $|| ||_v$ -topology on  $D \otimes_k k_v$ coincides under this isomorphism with the product topology  $\tau_S$  on  $D_S$ . Thus  $\tau_S$ restricts to the topology  $\tau_v$  on D.

Take a basis  $v_1, \ldots, v_m$  of D over k which is contained in  $N_{\geq 0}$ . Then  $D = kv_1 + \cdots + kv_m$ , with the  $\tau_v$ -topology, is homeomorphic to the direct product of m copies of k, with the v-adic topology, via the coordinate map. Let  $\mathcal{R}$  (resp.,  $\mathcal{R}_0$ ) denote the subring of D (resp., k) generated by  $N_{\geq 0}$  (resp.,  $(N \cap k^{\times})_{\geq 0}$ ). By (1) and (2), Theorem 9.5(3) applies and we conclude that the subring  $\mathcal{R}_0$  of k is v-adically open. We have

$$\mathcal{R}_0 \nu_1 + \cdots + \mathcal{R}_0 \nu_m \subseteq \mathcal{R},$$

so  $\mathcal{R}$  is open in D with respect to  $\tau_v$ .

Further, let  $\overline{\mathcal{R}}$  be the  $\tau_S$ -closure of  $\mathcal{R}$  in  $D_S$ . Then  $\iota_S^{-1}(\overline{\mathcal{R}})$  is the  $\tau_v$ -closure of  $\mathcal{R}$ in D. Being a  $\tau_v$ -open ring,  $\mathcal{R}$  is also  $\tau_v$ -closed in D, and therefore  $\mathcal{R} = \iota_S^{-1}(\overline{\mathcal{R}})$ . Also, since  $\mathcal{R}$  is  $\tau_v$ -open in D, the closure  $\overline{\mathcal{R}}$  is  $\tau_S$ -open in  $D_S$  (indeed, for an open ball  $B_D(0, \epsilon)$  in  $\mathcal{R}$  one has  $B_{D\otimes_k k_v}(0, \epsilon) \subseteq \overline{B_D(0, \epsilon)} \subseteq \overline{\mathcal{R}}$ ).

Next let  $S_0$  denote the set of all  $w \in S$  such that  $\operatorname{pr}_w(\overline{\mathcal{R}})$  is unbounded, and set  $T := S \setminus S_0$ . By Lemma 10.1,  $\overline{\mathcal{R}} = \operatorname{pr}_T(\overline{\mathcal{R}}) \times D_{S_0}$ . Consequently

$$\mathcal{R} = \iota_S^{-1}(\overline{\mathcal{R}}) = \iota_S^{-1}(\mathrm{pr}_T(\overline{\mathcal{R}}) \times D_{S_0}) = \iota_T^{-1}(\mathrm{pr}_T(\overline{\mathcal{R}})).$$
(10.2)

In case  $T = \emptyset$ , this would mean  $\mathcal{R} = D$ , which contradicts [RSS], Prop. 4.3(1). Thus, T is not empty. Furthermore, since  $\operatorname{pr}_T(\overline{\mathcal{R}})$  is open in  $D_T$ , we obtain from Eq. (10.2) that  $\mathcal{R}$  is open in D with respect to  $\tau_T$ , as required.

Next we show that each  $w \in T$  extends uniquely to D. Indeed, since  $\overline{\mathcal{R}}$  is  $\tau_S$ -open in  $D_S$ , the subring

$$\mathcal{R}(w) := \mathrm{pr}_w(\mathcal{R}) \cap D$$

is  $\tau_v$ -open in D. Since  $\operatorname{pr}_w(\overline{\mathcal{R}})$  is bounded, while D is obviously unbounded,  $\mathcal{R}(w) \neq D$ . Furthermore, being generated by  $N_{\geq 0}$ , the subring  $\mathcal{R}$  is invariant under conjugation by any element of N, so the same is true for  $\overline{\mathcal{R}}$  and consequently for  $\mathcal{R}(w)$ . Since  $(D^{\times})^{\ell} \subseteq N$ , for  $\ell = [D^{\times} : N]$ , we can apply Proposition 10.2 with  $\mathcal{B} = \mathcal{R}(w)$  and deduce that w extends uniquely to a height one valuation  $\widetilde{w}$  of D such that  $\mathcal{R}(w) \subseteq \mathcal{O}_{D,\widetilde{w}}$ . In particular,  $N_{\geq 0} \subseteq \mathcal{O}_{D,\widetilde{w}}$ , so  $\widetilde{w}$  is associated with  $\varphi$ .

The fact that w extends to D implies that  $D_w = D \otimes_F F_w$  is a division algebra (see [8] or [39, Th. 2.3]). Further,  $D_w$  can be identified with the completion of D with respect to  $\tilde{w}$ . Since  $D_w$  is finite-dimensional over the complete field  $F_w$ , the norm  $|| ||_w$  and the norm corresponding to  $\tilde{w}$  are equivalent, and hence induce

the same topology on  $D_w$ . It follows that the  $\tau_T$ -topology on D coincides with its  $\tilde{T}$ -adic topology. Therefore,  $\mathcal{R}$  is also  $\tilde{T}$ -adically open.

Next, by (iii), there exists  $\gamma \in \Gamma_{>0}$  such that  $\mathcal{R} \cap N \subseteq N_{>-\gamma}$ . Hence, Proposition 9.3 applies to  $\mathcal{M} = \mathcal{R}$  and gives the openness of N in the  $\tilde{T}$ -adic topology.

Assumption (iii) of Theorem 10.3 is satisfied in the following important situation:

**Lemma 10.4.** [28, Theorem 5.8(3)]. Let  $x, y \in D^{\times} \setminus N$  satisfy  $d(x^*, y^*) \ge 3$  and let  $\Gamma = \Gamma_{y^*}$  and  $\ge \ge_{y^*}$ . For the subring  $\mathcal{R}$  of D generated by  $N_{\ge_0}$ , there exists  $\gamma \in \Gamma_{>0}$  such that  $\mathcal{R} \cap N \subseteq N_{>-\gamma}$ .

We are now in a position to prove Theorems A and C of the Introduction and Theorem 7.5.

*Proof of Theorem C.* Take  $x, y \in D^{\times} \setminus N$  with  $d(x^*, y^*) \ge 3$  and set  $\ge = \ge_{y^*}$ . By Theorem 4.1,  $\varphi = \varphi_{y^*} \colon N \to \Gamma$  is strongly leveled map. Now Theorem 9.1 gives (1) and (2). Moreover, the valuation v in (2) is associated with  $\varphi \upharpoonright_{N \cap k^{\times}}$ . By Corollary 4.6(3),  $N_{\ge 0}$  contains a basis of D over k. By Lemma 10.4, there exists  $\gamma \in \Gamma_{>0}$  such that  $\mathcal{R} \cap N \subset N_{>-\gamma}$ , where  $\mathcal{R}$  is the subring of D generated by  $N_{\ge_0}$ . Therefore (3) follows from Theorem 10.3.

*Proof of Theorem* 7.5. Since diam( $\Delta$ )  $\geq$  3, and by Theorem 7.3, we can pick a nonidentity element  $y^* \in D^{\times}/N$  such that  $\varphi = \varphi_{y^*} \colon N \to \Gamma$  is a strongly leveled map as in Theorem B(1), and such that the subgroup  $\varphi_{y^*}(N \cap k^{\times})$  of  $\Gamma_{y^*}$  is totally ordered. Hence Theorem 7.5 follows from Theorem C.

We conclude this section with the proof of Theorem A.

*Proof of Theorem A.* By Theorem 5.1, we can pick a nonidentity element  $y^* \in D^{\times}/N$  such that  $\varphi = \varphi_{y^*} \colon N \to \Gamma$  is a strong valuation-like map.

Suppose first that D = K is abelian. Then, by Theorem 9.5(1), the assertion of Theorem A holds.

Next assume that *D* is a finite-dimensional division algebra over a field *K* of finite transcendence degree over its prime field. Of course *k* is infinite since *D* is. We may assume that k = F = Z(D). Since  $\varphi$  is a strong valuation-like map, the subgroup  $\varphi(N \cap F^{\times}) \subseteq \Gamma$  is totally ordered. Hence the assertion of Theorem A holds by Theorem C(3) (with k = F, and note that  $|T| = |\tilde{T}| = 1$ , since k = F.

## 11. Examples

#### 11.1. Constructions of V-graphs

*Example 11.1.* One can extend the notion of the relative Milnor *K*-rings from the case of commutative fields, as in [11, Part IV], to our general non-commutative context as follows. Let *D* be a division ring and let *N* be a normal subgroup of  $D^{\times}$ .

We stress however that, when N = 1, this definition is not the more common one, as in e.g. [29,30]. For  $r \ge 1$  we define the degree r Milnor K-group  $K_r^M(D)/N$ of D relative to N as the abelian group generated by all r-tuples  $\langle a_1^*, \ldots, a_r^* \rangle$  in  $(D^{\times}/N)^r$ , subject to the following defining relations:

(a) multi-linearity:

$$\langle a_1^*, \dots, a_i^*(a_i')^*, \dots, a_r^* \rangle = \langle a_1^*, \dots, a_i^*, \dots, a_r^* \rangle + \langle a_1^*, \dots, (a_i')^*, \dots, a_r^* \rangle$$

(b) Steinberg relations: ⟨a<sub>1</sub><sup>\*</sup>,..., a<sub>r</sub><sup>\*</sup>⟩ = 0 whenever 1 ∈ a<sub>i</sub><sup>\*</sup> + a<sub>j</sub><sup>\*</sup> for some distinct i, j.

In particular,  $K_1^M(D)/N$  is the maximal abelian quotient of  $D^{\times}/N$ . We also define  $K_0^M(D)/N = \mathbb{Z}$ .

We equip  $K_*^M(D)/N = \bigoplus_{r=0}^{\infty} K_r^M(D)/N$  with a graded ring structure, where the product is induced by concatenation. Denote the natural multi-linear map  $(D^{\times}/N)^r \to K_r^M(D^{\times})/N$  by  $(a_1^*, \ldots, a_r^*) \mapsto \{a_1, \ldots, a_r\}_N$ . Note that it is welldefined.

Now let  $a \in D^{\times}$ . We show that  $\{a, -a\}_N = 0$ . This is trivial for a = 1. For  $a \neq 1$  the identity

$$-a = (1-a)(a^{-1}(a-1))^{-1} = (1-a)(1-a^{-1})^{-1}$$

implies that

$$\{a, -a\}_N = \{a, 1-a\}_N + \{a, (1-a^{-1})^{-1}\}_N = \{a, 1-a\}_N + \{a^{-1}, 1-a^{-1}\}_N = 0,$$

as claimed. In the terminology of [11, Sect. 23.1] this shows that  $K_*^M(D)/N$  is a  $\kappa$ -structure with the distinguished element  $(-1)^*$ . It is now standard to see that  $K_*^M(D)/N$  is anti-commutative (see [11, Lemma 23.1.2]); indeed, for  $a, b \in D^{\times}$  we have

$$\{a, b\}_N + \{b, a\}_N = \{ab, ab\}_N - \{a, a\}_N - \{b, b\}_N = \{ab, -1\}_N - \{a, -1\}_N - \{b, -1\}_N = 0.$$

Now assume that  $(D^{\times} : N) < \infty$  and  $-1 \in N$ . Extending the construction in [12], we define the *Milnor K-graph of D relative to N* as the graph whose vertices are the non-trivial cosets in  $D^{\times}/N$ , and where vertices  $a^*$  and  $b^*$  are connected by an edge if and only if  $\{a, b\}_N = 0$  in  $K_2^M(D)/N$  (this relation is symmetric by the anti-commutativity). By construction, this graph satisfies condition (V1'). Exactly as in [12, Lemma 2.1] one shows using the bilinearity of  $\{\cdot, \cdot\}_N$  that it satisfies (V2) and (V3).

*Example 11.2.* Let *H* be a group and let  $\Theta$  be a graph on  $H \setminus \{1\}$  with distance function *d*. For  $h \in H \setminus \{1\}$  denote

$$\Theta_h := \{g \in H \setminus \{1\} \mid d(h, g) \le 1\} \cup \{1\}.$$

We say that  $\Theta$  is a *centralizer-graph* if  $\Theta_h$  is a subgroup of H contained in the centralizer  $C_H(h)$ , for all  $h \in H$ .

Notice that if we are given subgroups  $C_h \leq C_H(h)$ ,  $h \in H \setminus \{1\}$ , we can construct the minimal centralizer-graph  $\Theta$  on  $H \setminus \{1\}$  such that  $C_h \leq \Theta_h$ , for all  $h \in H \setminus \{1\}$ . This is just the intersection of all centralizer-graphs  $\Theta'$  such that  $C_h \leq \Theta'_h$ ,  $h \in H \setminus \{1\}$ .

For each non-identity  $a^* \in D^{\times}/N$  (notation as in subsection 1.1 of the Introduction), let  $C_{a^*} := \langle (a+n)^*, (a^{-1}+n)^* | n \in N \rangle$ . Let  $\Delta$  be the minimal centralizer-graph such that  $C_{a^*} \leq \Delta_{a^*}$ , for all  $a^* \in D^{\times}/N$ . Then  $\Delta$  is a V-graph, and it is the minimal centralizer-graph which is a V-graph.

#### 11.2. The bound diam $\geq 3$ is sharp in Theorem C(3)

In Theorem C(3), the bound 3 on the diameter is sharp. An example showing this for a commuting graph of the quaternion algebra  $(-1, -1/\mathbb{Q})$  was given in [25, Sect. 5]. The following analogous example shows this for a Milnor *K*-graph over  $\mathbb{Q}$ .

*Example 11.3.* Fix an odd integer *m* and a prime number *l* such that gcd(l-1, m) > 1. Thus  $\mathbb{F}_l^{\times} \neq (\mathbb{F}_l^{\times})^m$ . Similarly to [25, Sect. 5] and [27, p. 582] we consider the homomorphism

$$h: \mathbb{Q}^{\times} \to \mathbb{Z}, \quad \pm \prod_{i} p_{i}^{d_{i}} \mapsto \sum_{i} d_{i},$$

where the  $p_i$  denote the distinct primes,  $d_i \in \mathbb{Z}$ , and  $d_i = 0$  almost always. Let  $H = h^{-1}(m\mathbb{Z})$  and let  $U = (\mathbb{Q}^{\times})^m (1 + \mathfrak{m}_{v_l})$ , where  $v_l$  is the *l*-adic valuation on  $\mathbb{Q}$ . One has a split exact sequence

$$1 \to \mathbb{F}_l^{\times} / (\mathbb{F}_l^{\times})^m \to \mathbb{Q}^{\times} / U \xrightarrow{v_l} \mathbb{Z} / m \to 0,$$

(see e.g. [11, (3.2.7)]). Hence  $(\mathbb{Q}^{\times} : U) < \infty$ . Setting  $N = H \cap U$ , we obtain that

$$-1 \in (\mathbb{Q}^{\times})^m \le N \le \mathbb{Q}^{\times}, \quad (\mathbb{Q}^{\times}:N) < \infty.$$

We show that N cannot be open with respect to any finite set of non-trivial valuations on  $\mathbb{Q}$ . To this end let  $q_1, \ldots, q_t$  be distinct prime numbers, let  $r_1, \ldots, r_t$  be positive integers, and consider the  $q_i$ -adic valuations  $v_{q_i}, i = 1, \ldots, t$ . We need to show that N does not contain the congruence subgroup

$$W = 1 + q_1^{r_1} \mathcal{O}_{v_{q_1}} \cap \cdots \cap q_t^{r_t} \mathcal{O}_{v_{q_t}}.$$

Indeed, Dirichlet's theorem on primes in arithmetic progressions yields a positive integer k with  $p = 1 + kq_1^{r_1} \cdots q_t^{r_t}$  prime. Thus h(p) = 1, so  $p \notin H$ . Also note that  $p \in W$ . Consequently,  $W \not\subseteq H$ , whence  $W \not\subseteq N$ , as desired.

Finally we compute the diameter of the graph of  $K_*^M(\mathbb{Q})/N$ . In view of Theorem C (with  $D = F = k = \mathbb{Q}$ ) and what we have just seen, the diameter is at most 2. To show that it is exactly 2 we need to verify that  $K_2^M(\mathbb{Q})/N \neq 0$ . Now there is a canonical isomorphism of  $\kappa$ -structures

$$K^M_*(\mathbb{Q})/U \cong (K^M_*(\mathbb{F}_l)/(\mathbb{F}_l^{\times})^m)[\mathbb{Z}/m]$$

[11, Th. 26.1.2]. It follows that  $K_2^M(\mathbb{Q})/U \neq 0$  (see [11, Example 23.2.5]). Applying the canonical epimorphism Res:  $K_*^M(\mathbb{Q})/N \to K_*^M(\mathbb{Q})/U$  we get that  $K_2^M(\mathbb{Q})/N \neq 0$  as well. More concretely, take an *l*-adic unit *a* in  $\mathbb{Z}$  whose residue  $\bar{a} \in \mathbb{F}_l^{\times}$  is not an *m*th power. Then  $aU \otimes lU \notin St_{\mathbb{Q},2}(U)$ , and therefore also  $aN \otimes lN \notin St_{\mathbb{Q},2}(N)$ , i.e.,  $\{a, l\}_N \neq 0$  in  $K_2^M(\mathbb{Q})/N$ .

We remark that, by contrast, the diameter of  $K_*^M(\mathbb{Q})/H$  is 1. Indeed,  $\mathbb{Q}^{\times}/H$  is cyclic and *H* is not contained in any ordering (since  $-1 \in H$ ). Hence, by [11, Th. 25.2.1],  $K_2^M(\mathbb{Q})/H = 0$ , and the assertion follows.

## 11.3. Examples for Theorem C and property $(3\frac{1}{2})$

Examples 11.4 and 11.7 below show that Theorem C covers a more general situation than the previous works [12,25,27,28]. These examples also illustrate property  $(3\frac{1}{2})$ , and shows that Theorem 7.3 covers a more general situation than [28, Theorem 5.8].

*Example 11.4.* First we study the following local situation. Let D be a division algebra over a field F and  $\tilde{w}$  a discrete valuation on D with uniformizer  $\pi$ . Let  $U = O_{D,\tilde{w}}^{\times}$  be the group of  $\tilde{w}$ -units in D, let  $U^{(1)} = U_{\tilde{w}}^{(1)} = 1 + \mathfrak{m}_{D,\tilde{w}}$  be its first congruence subgroup, and let  $\overline{D} = \overline{D}_{\tilde{w}}$  be the residue field. Suppose that e is a positive integer with  $\pi^e \in Z(D)$ . Then

$$N = \langle \pi^e, U^{(1)} \rangle = \langle \pi^e \rangle \times U^{(1)}$$

is a normal subgroup of  $D^{\times}$ . One has an isomorphism

$$U/(U \cap N) = U/U^{(1)} \cong \overline{D}^{\times}$$

Therefore  $\widetilde{w}$  induces an exact sequence

$$1 \to \overline{D}^{\times} \to D^{\times}/N \to \mathbb{Z}/e\mathbb{Z} \to 0.$$
(11.1)

Assume further that the residue characteristic of  $\widetilde{w}$  is 2. Then  $-1 \in U^{(1)} \subseteq N$ .

Let N(y) be as in Sect. 3.

**Lemma 11.5.** Let  $y \in D^{\times} \setminus N$ . Then  $N(y) = \{n \in N \mid \widetilde{w}(n) < \widetilde{w}(y)\}$ .

*Proof.* Set  $s = \widetilde{w}(y)$ . Consider an arbitrary  $n \in N$  and write it in the form  $n = \pi^t u$ , with  $t = \widetilde{w}(n) \in e\mathbb{Z}$  and  $u \in U^{(1)}$ .

Case 1. t < s. Then  $\widetilde{w}(\pi^{-t}y) > 0$ , so  $\pi^{-t}y + u \in U^{(1)}$ , implying that  $y + n = \pi^t(\pi^{-t}y + u) \in N$  and  $n \in N(y)$ .

Case 2. t = s. Then  $-\pi^{-t}y \in U \setminus U^{(1)}$ , so  $\pi^{-t}y + u \in U \setminus U^{(1)}$ , and therefore  $y + n = \pi^t (\pi^{-t}y + u) \notin N$ , i.e.,  $n \notin N(y)$ .

*Case 3.* t > s. Then  $\widetilde{w}(\pi^{t-s}u) > 0$ . Since  $\pi^{-s}y \in U \setminus U^{(1)}$ , we have  $\pi^{-s}y + \pi^{t-s}u \in U \setminus U^{(1)}$ , and consequently  $y + n \notin N$  and  $n \notin N(y)$ .  $\Box$ 

Next we consider the following semi-local situation. Let  $\widetilde{w}_1, \ldots, \widetilde{w}_r$  be distinct discrete valuations on the division algebra D with residue characteristic 2. Suppose that  $\pi \in D$  is a common uniformizer for  $\widetilde{w}_1, \ldots, \widetilde{w}_r$ , and that  $\pi^e \in Z(D)$  with  $e \ge 1$ . Set  $N_i = \langle \pi^e \rangle \times U_{\widetilde{w}_i}^{(1)}$  and  $N = N_1 \cap \cdots \cap N_r$ . By the weak approximation theorem,  $D^{\times}/N \cong \prod_{i=1}^r D^{\times}/N_i$  via the diagonal map. In the terminology of Sect. 3, we set  $U = U_{\pi^*}$  and  $\Gamma = N/U$ .

Lemma 11.6. There is an isomorphism of partially ordered groups

$$\Gamma = N/U \xrightarrow{\sim} \mathbb{Z}^r, \quad nU \mapsto \left(\frac{1}{e}\widetilde{w}_1(n), \ldots, \frac{1}{e}\widetilde{w}_r(n)\right),$$

where  $\mathbb{Z}^r$  is endowed with the product partial order relation. Further,

$$U = U_{\widetilde{w}_1}^{(1)} \cap \dots \cap U_{\widetilde{w}_r}^{(1)}.$$

*Proof.* By Lemma 11.5 with  $y = \pi$ ,

$$N(\pi) = N_1(\pi) \cap \dots \cap N_r(\pi) = \{t \in N \mid \widetilde{w}_1(t), \dots, \widetilde{w}_r(t) \le 0\}.$$
 (11.2)

Hence  $1 \in N(\pi)$ , and every  $n \in U_{\widetilde{w}_1}^{(1)} \cap \cdots \cap U_{\widetilde{w}_r}^{(1)}$  satisfies  $nN(\pi) = N(\pi)$ . Conversely, if  $n \in N$  satisfies  $nN(\pi) = N(\pi)$ , then  $n, n^{-1} \in N(\pi)$ , so  $\widetilde{w}_i(n) = 0$  for every *i*. It follows that  $U = U_{\widetilde{w}_1}^{(1)} \cap \cdots \cap U_{\widetilde{w}_r}^{(1)}$ .

By the weak approximation,

$$\Gamma = N/U \cong N_1/U_{\widetilde{w}_1}^{(1)} \times \dots \times N_r/U_{\widetilde{w}_r}^{(1)} \cong \mathbb{Z}^r,$$
(11.3)

where the right isomorphism is induced by  $(\frac{1}{e}\widetilde{w}_1, \ldots, \frac{1}{e}\widetilde{w}_r)$ .

Moreover, if  $m, n \in N$  and  $N(m\pi) \subseteq N(n\pi)$ , then  $mN(\pi) \subseteq nN(\pi)$ , and (11.2) implies that  $\widetilde{w}_i(m) \leq \widetilde{w}_i(n)$  for i = 1, 2, ..., r. Therefore (11.3) is an isomorphism of partially ordered groups.

As a concrete example, take  $d \in \mathbb{Q}$  such that  $\sqrt{d} \in \mathbb{Q}_2 \setminus \mathbb{Q}$ ; e.g., we may take d = 17 (see [11, Prop. 18.2.1]). Let k be  $\mathbb{Q}$ , or more generally, a subfield of  $\mathbb{Q}_2$  which does not contain  $\sqrt{d}$ . Let  $F = k(\sqrt{d})$ , and let  $\sigma$  be the nontrivial automorphism in Gal(F/k). Let  $v, w_1$  be the restrictions of the 2-adic valuation of  $\mathbb{Q}_2$  to k, F, respectively. Also let  $w_2 = w_1 \circ \sigma$ . Then  $w_1, w_2$  are unramified over v and have residue field  $\mathbb{F}_2$ . Further, they are distinct, and are the only extensions of v to F. Consider the quaternion algebra D = (-1, -1/F) over F with its standard basis  $\mathbf{1}, \mathbf{i}, \mathbf{j}, \mathbf{k}$ , and take e = 2. Then  $w_1, w_2$  uniquely extend to valuations  $\widetilde{w}_1, \widetilde{w}_2$ , respectively, on D with  $(\widetilde{w}_i(D^{\times}) : w_i(F^{\times})) = 2$  and  $\overline{D}_{\widetilde{w}_i} = \mathbb{F}_4, i = 1, 2$ . Specifically, let

$$\pi = \mathbf{i} + \mathbf{j}, \quad a = \frac{1}{2}(-1 + \mathbf{i} + \mathbf{j} + \mathbf{k}) = -\frac{1}{2}(1 - \mathbf{j})(1 - \mathbf{i}).$$

Note that  $\pi^2 = -2$  and  $a^2 + a + 1 = 0$ . Then  $\pi$  is a uniformizer for  $\widetilde{w}_i$  and a is a  $\widetilde{w}_i$ -unit with residue not in  $\mathbb{F}_2$ , i = 1, 2. Let

$$N_i = \langle \pi^2 \rangle \times U_{\widetilde{w}_i}^{(1)} = \langle -2 \rangle \times U_{\widetilde{w}_i}^{(1)}, \quad N = N_1 \cap N_2.$$

By the exact sequence (11.1), the group  $D^{\times}/N_i$  has order 6 (where i = 1, 2). We show that it is noncommutative. Indeed,

$$\pi a \pi^{-1} a^{-1} = \frac{1}{2} (\mathbf{1} - \mathbf{i} + \mathbf{j} + \mathbf{k}) = \frac{1}{2} (\mathbf{1} + \mathbf{j})(\mathbf{1} - \mathbf{i}),$$
  
$$\pi a \pi^{-1} a^{-1} - 1 = \frac{1}{2} (-\mathbf{1} - \mathbf{i} + \mathbf{j} + \mathbf{k}) = -\frac{1}{2} (\mathbf{1} + \mathbf{i})(\mathbf{1} - \mathbf{j})$$

are in  $U_{\widetilde{w}_i}$ . Thus  $\pi a \pi^{-1} a^{-1} \in U_{\widetilde{w}_i} \setminus U_{\widetilde{w}_i}^{(1)}$ , so  $aN_i$  and  $\pi N_i$  do not commute in  $D^{\times}/N_i$  (this also follows from the general fact that, if  $\mathcal{D}$  is a finite-dimensional central division algebra over a complete discretely valued field  $\mathcal{K}$  with a perfect residue field  $\overline{\mathcal{K}}$ , then the center  $Z(\overline{\mathcal{D}})$  is a cyclic Galois extension of  $\overline{\mathcal{K}}$ , and a uniformizer  $\Pi$  in  $\mathcal{D}$  induces a generator of  $\operatorname{Gal}(Z(\overline{\mathcal{D}})/\overline{\mathcal{K}})$ , cf. [39, Prop. 2.5]. In our situation,  $\overline{a} \in Z(\overline{\mathcal{D}}) \setminus \overline{\mathcal{K}} = \mathbb{F}_4 \setminus \mathbb{F}_2$ , so this generator must act non-trivially on  $\overline{a}$ ).

Consequently,  $D^{\times}/N_1 \cong D^{\times}/N_2 \cong S_3$  and

$$D^{\times}/N \cong S_3 \times S_3. \tag{11.4}$$

Observe that the diameter of the commuting graph of  $S_3 \times S_3$  is 3. Indeed, any  $(\sigma_1, \sigma_2), (\tau_1, \tau_2) \in S_3 \times S_3$  are connected in the commuting graph by the following path of length 3:

$$(\sigma_1, \sigma_2)$$
,  $(1, \sigma_2)$ ,  $(\tau_1, 1)$ ,  $(\tau_1, \tau_2)$ 

provided that  $\sigma_2$ ,  $\tau_1 \neq 1$ ; other cases are considered similarly. On the other hand, if  $\sigma_1, \sigma_2 \in S_3$  are transpositions and  $\tau_1, \tau_2 \in S_3$  are 3-cycles, then  $(\sigma_1, \sigma_2)$  and  $(\tau_1, \tau_2)$  are not at distance  $\leq 2$ .

Now  $\pi N_i$  has order 2 and  $aN_i$  has order 3 in  $D^{\times}/N_i \cong S_3$ . Hence  $\pi N, aN \in D^{\times}/N$  correspond under (11.4) to elements of the form ( $\sigma_1, \sigma_2$ ), ( $\tau_1, \tau_2$ ), respectively, where  $\sigma_1, \sigma_2$  are transpositions and  $\tau_1, \tau_2$  are 3-cycles. It follows that  $d(aN, \pi N) = 3$ .

Therefore the assumptions of Theorem C are satisfied with *k* a field of finite transcendence degree as above,  $y = \pi$ , and with  $\varphi = \varphi_{y^*} \colon N \to \Gamma = N/(U_{\widetilde{w}_1} \cap U_{\widetilde{w}_2}) \cong \mathbb{Z}^2$  (see Lemma 11.6). Then  $\varphi(nU) = (\frac{1}{2}v(n), \frac{1}{2}v(n))$  for  $n \in N \cap k^{\times}$ , so  $\varphi(N \cap k^{\times}) \cong \mathbb{Z}$  is indeed totally ordered. This also shows that  $\varphi|_{N \cap k^{\times}}$  is a strong valuation-like map, and  $N \cap k^{\times}$  is open in the *v*-adic topology on *k*, in accordance with (1) and (2) of Theorem C. By construction, (3) of Theorem C holds with  $T = \{w_1, w_2\}$  and  $\widetilde{T} = \{\widetilde{w}_1, \widetilde{w}_2\}$ .

On the other hand, the results of [27,28] do not apply to characterize *N*. Namely, by Lemma 8.3, *N* is not  $\tilde{u}$ -adically open for any nontrivial valuation  $\tilde{u}$  on *D*. Technically, the results of [25] may not apply either as *k* and *F* do not have to be number fields (although some methods of [25] were instrumental in proving Theorem C).

This construction also provides an example of a situation where property  $(3\frac{1}{2})$  of Sect. 7 applies. More precisely, the nontrivial automorphism  $\sigma$  of Gal(F/k) extends to an automorphism  $\sigma_D$  of order 2 of the algebra D. This automorphism

switches the valuations  $w_1$  and  $w_2$  of F, the valuations  $\tilde{w}_1$  and  $\tilde{w}_2$  of D, and the subgroups  $N_1$  and  $N_2$  of  $D^{\times}$ . The subgroup N of  $D^{\times}$  is invariant under  $\sigma_D$ , and  $\sigma_D$  switches the factors in  $D^{\times}/N = S_3 \times S_3$ . Consider elements  $x^* = (\sigma_1, \sigma_2)$  and  $y^* = (\tau_1, \tau_2)$  as above. We claim that  $D^{\times}/N$  satisfies  $(3\frac{1}{2})$  with respect to  $x^*$ ,  $y^*$ , the group  $\Sigma = \{id, \sigma_D\}$ , and  $M = N \cap k^{\times}$ . Indeed, essentially the only path of length 3 between  $x^*$  and  $y^*$  is

$$x^*$$
,  $(\sigma_1, 1)$ ,  $(1, \tau_2)$ ,  $y^*$ .

But  $d(\sigma_D(x^*), y^*) = 3$ , and  $\sigma_D(\sigma_1, 1) = (1, \sigma_1)$  and  $(1, \tau_2)$  do not commute.

*Example 11.7.* We give a similar construction for a Milnor *K*-graph. Let *p* be a prime number such that  $p \equiv 1 \pmod{4}$ . Let F/k be a nontrivial Galois field extension such that  $F \subseteq \mathbb{Q}_p$ . Let v, w be the restrictions of the *p*-adic valuation on  $\mathbb{Q}_p$  to *k*, *F*, respectively. The extension (F, w)/(k, v) is immediate, i.e., has the same value group  $\mathbb{Z}$  and residue field  $\mathbb{F}_p$ . Since *v* is discrete, this implies that there are exactly [F : k] extensions of *v* to *F*, namely  $w \circ \sigma$  with  $\sigma \in \text{Gal}(F/k)$  (see [11, Cor. 17.4.4]). Set  $w_1 = w$  and  $w_2 = w \circ \sigma$  with  $\sigma \neq \text{id}$ .

Now set

$$N_1 = (F^{\times})^2 (1 + \mathfrak{m}_{w_1}), \quad N_2 = (F^{\times})^2 (1 + \mathfrak{m}_{w_2}), \quad N = N_1 \cap N_2$$

Since  $p \equiv 1 \pmod{4}$  we have  $-1 \in (\mathbb{F}_p^{\times})^2 = (\overline{F}_{w_i}^{\times})^2$ , i = 1, 2, and therefore  $-1 \in N$ .

We compute the relative Milnor K-ring  $K_*^M(F)/N$ . We use the terminology and results of [11, Part IV]. First, the graded ring  $K_*^M(\mathbb{F}_p)/(\mathbb{F}_p^{\times})^2$  is  $\mathbb{F}_p^{\times}/(\mathbb{F}_p^{\times})^2 \cong \mathbb{Z}/2\mathbb{Z}$ in degree 1, and is trivial in degrees  $\geq 2$  [11, Cor. 25.2.4]. Thus  $K_*^M(\mathbb{F}_p)/(\mathbb{F}_p^{\times})^2$  is the extension  $\mathbf{0}[\mathbb{Z}/2\mathbb{Z}]$  of the trivial  $\kappa$ -structure  $\mathbf{0}$  by the group  $\mathbb{Z}/2\mathbb{Z}$  [11, Example 23.2.4]. Since  $w(N_1) = 2\mathbb{Z}$ , [11, Th. 26.1.2 and Example 26.1.1(1)], says that  $K_*^M(F)/N_1$  is the extension  $(K_*^M(\mathbb{F}_2)/(\mathbb{F}_p^{\times})^2)[\mathbb{Z}/2\mathbb{Z}]$ . It follows that

$$K^M_*(F)/N_1 \cong (\mathbf{0}[\mathbb{Z}/2\mathbb{Z}])[\mathbb{Z}/2\mathbb{Z}] \cong \mathbf{0}[(\mathbb{Z}/2\mathbb{Z})^2]$$

(see [11, Lemma 23.2.3]). Similarly,  $K_*^M(F)/N_2 \cong \mathbf{0}[(\mathbb{Z}/2\mathbb{Z})^2]$ .

Since w, w' are distinct and discrete, they are independent. Therefore [11, Cor. 28.2.4], shows that

$$K^{M}_{*}(F)/N \cong \left(K^{M}_{*}(F)/N_{1}\right) \times \left(K^{M}_{*}(F)/N_{2}\right) \cong \left(\mathbf{0}[(\mathbb{Z}/2\mathbb{Z})^{2}]\right) \times \left(\mathbf{0}[(\mathbb{Z}/2\mathbb{Z})^{2}]\right).$$

The Milnor K-graph of F relative to N was computed in [12, Sect. 7], and it has diameter 3.

We may further take k to be finitely generated over  $\mathbb{Q}$ . By construction, N is open in the T-adic topology, where  $T = \{w_1, w_2\}$ . This is in accordance with Theorem C, when we take D = F and  $T = \widetilde{T}$ .

Finally, by Lemma 8.3, N is not *u*-adically open with respect to any single non-trivial valuation u on F.

In this example as well property  $(3\frac{1}{2})$  holds, when we take F/k to be an extension of degree 2,  $\Sigma = \text{Gal}(F/k)$ , and  $M = N \cap k^{\times}$ . The generator  $\sigma$  of Gal(F/k)

switches  $N_1$ ,  $N_2$ , and therefore fixes N. For i = 1, 2, the group  $F^{\times}/N_i$  is the degree 1 component of  $K_*^M(F)/N_i \cong \mathbf{0}[(\mathbb{Z}/2\mathbb{Z})^2]$ , and is isomorphic to  $(\mathbb{Z}/2\mathbb{Z})^2$ . Let  $\alpha, \beta, \gamma, \delta$  be generators of  $F^{\times}/N$  such that  $\alpha, \beta$  project to generators  $F^{\times}/N_1$  and  $\gamma, \delta$  project to generators of  $F^{\times}/N_2$ . We may assume that  $\sigma$  switches  $\alpha, \gamma$  and  $\beta, \delta$ . As was shown in [12, Sect. 7] (in a multiplicative notation), the vertices  $\alpha + \gamma, \beta + \delta$  of the Milnor *K*-graph of *F* relative to *N* have distance 3, and there are exactly two paths of length 3 connecting them, namely

$$\alpha + \gamma, \alpha, \delta, \beta + \delta$$
 and  $\alpha + \gamma, \gamma, \beta, \beta + \delta$ .

Further,  $\sigma(\alpha + \gamma) = \alpha + \gamma$ , but the vertices  $\gamma = \sigma(\alpha)$  and  $\delta$  are not connected by an edge, and neither are the vertices  $\alpha = \sigma(\gamma)$  and  $\beta$ . Therefore property  $(3\frac{1}{2})$  holds for  $x^* = \alpha + \gamma$  and  $y^* = \beta + \delta$ .

Acknowledgements. We warmly thank the referee for the thorough, quick, and extremely useful report, which helped us to correct certain flaws, as well as to improve the exposition of the paper.

## References

- Arason, J.K., Elman, R., Jacob, B.: Rigid elements, valuations, and realization of Witt rings. J. Algebra 110, 449–467 (1987)
- [2] Becker, E.: Summen *n*-ter Potenzen in Körpern. J. Reine Angew. Math. 307(308), 8– 30 (1979)
- [3] Bergelson, V., Shapiro, D.B.: Multiplicative subgroups of finite index in a ring. Proc. Am. Math. Soc. 116, 885–896 (1992)
- [4] Bogomolov, F., Tschinkel, Y.: Reconstruction of function fields. GAFA 18, 400– 462 (2008)
- [5] Bogomolov, F., Tschinkel, Y.: Introduction to birational anabelian geometry. In: Caporaso, L., McKernan, J., Mustata, M., Popa, M. (eds.) Current Developments in Algebraic Geometry, vol. 59, pp. 17–63. Cambridge University Press, Cambridge (2012)
- [6] Bogomolov, F., Tschinkel, Y.: Galois theory and projective geometry. Commun. Pure Appl. Math. 66, 1335–1359 (2013)
- [7] Bourbaki, N.: Commutative Algebra, ch. 1–7. Springer, Berlin (1989)
- [8] Cohn, P.M.: On extending valuations in division algebras. Stud. Sci. Math. Hungar. 16, 65–70 (1981)
- [9] Efrat, I.: Construction of valuations from K-theory. Math. Res. Lett. 6, 335–343 (1999)
- [10] Efrat, I.: Compatible valuations and generalized Milnor *K*-theory. Trans. Am. Math. Soc. 359, 4695–4709 (2007)
- [11] Efrat, I.: Valuations, Orderings, and Milnor K-Theory. Mathematical Surveys and Monographs, vol. 124. American Mathematical Society, Providence (2006)
- [12] Efrat, I.: Valuations and diameters of Milnor K-rings. Isr. J. Math. 172, 75–92 (2009)
- [13] Endler, O.: Valuation Theory. Springer, Berlin (1972)
- [14] Engler, A.J., Prestel, A.: Valued Fields. Springer, Berlin (2005)
- [15] Hwang, Y.S., Jacob, B.: Brauer group analogues of results relating the Witt ring to valuations and Galois theory. Can. J. Math. 47, 527–543 (1995)
- [16] Jacob, B.: On the structure of Pythagorean fields. J. Algebra 68, 247–267 (1981)
- [17] Jacob, B.: Fans, real valuations, and hereditarily-pythagorean fields. Pac. J. Math. 93, 95–105 (1981)

- [18] Jacob, B., Wadsworth, A.R.: A new construction of noncrossed division algebras. Tran. Am. Math. Soc. 293, 693–721 (1986)
- [19] Koenigsmann, J.: From *p*-rigid elements to valuations (with a Galois-characterization of *p*-adic fields). J. Reine Angew. Math. 465, 165–182 (1995)
- [20] Lang, S.: Algebra, 3rd edn, GTM 211. Springer, Berlin (2002)
- [21] Marshall, M.A.: An approximation theorem for coarse V-topologies on rings. Can. Math. Bull. 37, 527–533 (1994)
- [22] Marubayashi, H., Miyamoto, H., Ueda, A.: Non-Commutative Valuation Rings and Semi-Hereditary Orders. Kluwer, Dordrecht (1997)
- [23] Pierce, R.S.: Associative Algebras, GTM 88. Springer, Berlin (1982)
- [24] Platonov, V.P., Rapinchuk, A.S.: Algebraic Groups and Number Theory. Academic Press, New York (1993)
- [25] Rapinchuk, A.S.: The Margulis–Platonov conjecture for SL<sub>1,D</sub> and 2-generation of finite simple groups. Math. Z. 252, 295–313 (2006)
- [26] Rapinchuk, A., Potapchik, A.: Normal subgroups of SL<sub>1,D</sub> and the classification of finite simple groups. Proc. Indian Acad. Sci. 106, 329–336 (1996)
- [27] Rapinchuk, A.S., Segev, Y.: Valuation-like maps and the congruence subgroup property. Invent. Math. 144, 571–607 (2001)
- [28] Rapinchuk, A.S., Segev, Y., Seitz, G.: Finite quotients of the multiplicative group of a finite dimensional division algebra are solvable. J. Am. Math. Soc. 15, 929–978 (2002)
- [29] Rehmann, U.: Zentrale Erweiterungen der speziellen linearen Gruppe eines Schiefkörpers. J. Reine Angew. Math. 301, 77–104 (1978)
- [30] Rehmann, U., Stuhler, U.: On K<sub>2</sub> of finite-dimensional division algebras over arithmetical fields. Invent. Math. 50, 75–90 (1978/79)
- [31] Ribenboim, P.: The Theory of Classical Valuations. Springer, Berlin (1999)
- [32] Schilling, O.F.G.: The Theory of Valuations. American Mathematical Society, Providence (1950)
- [33] Segev, Y.: The commuting graph of minimal nonsolvable groups. Geom. Dedic. **88** (1–3), 55–66 (2001)
- [34] Segev, Y.: On finite homomorphic images of the multiplicative group of a division algebra. Ann. Math. 149, 219–251 (1999)
- [35] Segev, Y., Seitz, G.M.: Anisotropic groups of type  $A_n$  and the commuting graph of finite simple groups. Pac. J. Math. **202**, 125–226 (2002)
- [36] Topaz, A.: Almost-commuting-liftable subgroups of Galois groups. Preprint, available at arXiv:1202.1786
- [37] Topaz, A.: Commuting-liftable subgroups of Galois groups II. Preprint, available at arXiv:1208.0583
- [38] Turnwald, G.: Multiplicative subgroups of finite index in rings. Proc. Am. Math. Soc. 120, 377–381 (1994)
- [39] Wadsworth, A.R.: (2002) Valuation theory on finite dimensional division algebras. In: Saskatoon, S.K. (eds) Valuation Theory and Its Applications, vol. I. Fields Inst. Commun., vol. 32, pp. 385–449. American Mathematical Society, Providence, RI (1999)
- [40] Ware, R.: Valuation rings and rigid elements in fields. Can. J. Math. 33, 1338– 1355 (1981)