

Effective Action and Cluster Properties of the Abelian Higgs Model

Tadeusz Bałaban^{1,*}, John Z. Imbrie^{2,**}, and Arthur Jaffe^{2,***}

¹ Department of Mathematics, Northeastern University, Boston, MA 02115, USA

² Harvard University, Cambridge, MA 02138, USA

Abstract. We continue our program to establish the Higgs mechanism and mass gap for the abelian Higgs model in two and three dimensions. We develop a multiscale cluster expansion for the high frequency modes of the theory, within a framework of iterated renormalization group transformations. The expansions yield decoupling properties needed for a proof of exponential decay of correlations. The result of this analysis is a gauge invariant unit lattice theory with a deep Higgs potential of the shape required to exhibit the Higgs mechanism.

Table of Contents

1. Introduction	258
2. Localized Kernels	260
3. The First Renormalization Step	265
4. The Inductive Hypothesis	273
5. Renormalization and Decoupling in the General Step	277
5.1. Renormalization Transformation	277
5.2. Restrictions on the Fields	278
5.3. First Gauge Field Translation	280
5.4. Gauge Transformation	281
5.5. Second Gauge Field Translation	283
5.6. Expansion with Respect to the Fluctuation Field	285
5.7. The Gaussian Normalization Factors	289
5.8. Scalar Field Translation	295
5.9. Bounds on Fluctuation and Block Fields	296
5.10. The Interaction for the Fluctuation Fields	298
5.11. Mayer Expansion I	299
5.12. Conditional Integration	300
5.13. Decoupling of the Small Field Region	303
5.14. Resummation and Extraction of the Perturbation Expansion	307
5.15. Second Mayer Expansion and Scaling	312
References	314

* Research partially supported by the National Science Foundation under Grant DMS-8602207 and by the Air Force Office of Scientific Research under Grant AFOSR-86-0229

** Alfred P. Sloan Research Fellow. Research partially supported by the National Science Foundation under Grants PHY-84-13285 and PHY-85-13554

*** Research partially supported by the National Science Foundation under Grant PHY-85-13554

1. Introduction

We wish to establish the existence of a mass gap for the abelian Higgs model on the subspace of gauge invariant observables. Earlier work on this problem has led to a method to establish these results and to a partial solution [1, 2]. Here we continue this study with the development of a multiscale expansion suitable for the problem. The basic formulation of the model is given in [2]. We consider an action function S^ε which is defined for a gauge theory on a lattice with spacing ε . We use the Wilson form of lattice action, which is gauge invariant. Thus it is important to consider gauge invariant observables such as loop variables

$$u(\gamma) = \exp \left[\sum_{b \in \gamma} ie\varepsilon A(b) \right], \quad (1.1)$$

where γ is a closed curve on the lattice, or string variables

$$s(x, y, \Gamma) = \overline{\phi(x)} \exp \left(ie\varepsilon \sum_{b \in \Gamma} A(b) \right) \phi(y), \quad (1.2)$$

where Γ is a lattice curve from x to y . These variables must be renormalized appropriately, by multiplying or subtracting ε -dependent terms. For gauge invariant operators (but not in general) we expect exponential clustering in the equilibrium state defined by S^ε . This state is given by the limit of normalized finite volume expectations

$$\langle B \rangle = \frac{1}{Z} \int e^{-S^\varepsilon} B(u, \phi) \mathcal{D}u \mathcal{D}\phi. \quad (1.3)$$

(We assume periodic boundary conditions, but this is not crucial since as a corollary we establish the existence of the infinite volume limit.) Thus for gauge invariant functions B, C we expect

$$|\langle BC \rangle - \langle B \rangle \langle C \rangle| \leq O(1) \exp[-m \text{dist}(B, C)], \quad (1.4)$$

where $0 < m$ and $\text{dist}(B, C)$ denotes the distance between the supports of B and C . For unit lattice models, (1.4) was established in [3] and here we investigate the corresponding estimates uniformly in the lattice spacing ε .

The exponential decay or mass gap is intimately connected with the Higgs mechanism. We see the Higgs mechanism at work through the evolution of the effective action as we proceed lower in momentum. The action on the ε -lattice appears almost massless, but as we approach the unit lattice, the Higgs potential exhibits a pronounced ring of minima at $|\phi| = \varrho_0$, which leads to a mass term for the gauge field. The apparently massless rotational degrees of freedom of ϕ can be gauged away.

To obtain decay, we need a convergent expansion with a small parameter. Thus, we restrict the coupling constants (e, λ) to be sufficiently small in order to use cluster expansions. Such methods yield a nonperturbative analysis of the vacuum state, by explicitly displaying the exponential decay (1.4). Classically, the gauge field mass is of order $e/\lambda^{1/2}$, so we choose this ratio to be a fixed number of order unity.

The general ideas of these methods were described in our earlier papers [1, 2, 4]. Gauge invariance enters in a crucial way, both in the Higgs mechanism

described above and in the control of ultraviolet divergences. By separating high and low momentum parts of the interaction in a gauge invariant way, we can choose convenient gauges to discuss renormalization of the high momentum part and to discuss the spectrum of the Hamiltonian in the low momentum region.

In the present paper, we consider clustering properties of the high frequency modes of the model. Our goal is an exact expression for the effective action on the unit lattice. The expression is complicated by the need to treat large field or large action regions differently from the perturbative, small field regions. The effective unit lattice theory is, however, similar in spirit to the one considered in [3]. A final cluster expansion will be performed on this theory in another paper, and the proof of clustering and of the existence of the mass gap will then be complete.

The dual requirements of clustering and of the renormalization group force us to develop a cluster expansion for each frequency mode separately. We formulate an inductive form of the model after k renormalization group transformations, and then the bulk of this paper is devoted to clustering of the $(k + 1)$ -st mode in the next renormalization transformation.

The heart of our method is the way we accomplish changes of gauge without spoiling the exponential decoupling properties of the functional integral. We integrate out each frequency mode using a simple “tree” gauge on blocks (called axial gauge). After a number of modes have been integrated out, such gauges are not sufficiently regular to allow control of all the terms in the expansion. This is expected even in perturbation theory, where only gauges such as the Landau gauge are well behaved in the ultraviolet. Thus we must change the gauge in which those modes are expressed in the effective action. In keeping with the locality requirements of the cluster expansion, the change of gauge must be performed in patches, with slightly different changes on the overlaps. It turns out that the effects of the lack of alignment are small, and this way we avoid building up effects over long distances – something that tends to happen when changing gauges globally.

A similar problem occurs in our treatment of the effective unit lattice model. We have to change from the Landau gauge to the unitary gauge that is best suited for exhibiting the Higgs mechanism. Again, this must be accomplished without spoiling decoupling. Thus the method for changing gauges is the crucial aspect of our analysis, both for high momenta and for low momenta.

This paper is organized as follows. Having discussed Green’s functions in [2] as global operators, we start by introducing localized forms of these operators which are better suited to the cluster expansion. We replace kernels $G(x, y)$ with kernels $G_{\text{loc}}(x, y) = G(x, y)\zeta(x, y)$, where ζ is smooth and supported in some neighborhood of $x = y$. This section also serves to review the roles of the various operators. We then briefly describe the cluster expansion in the first renormalization step. This leads to the formulation of the inductive hypothesis for the form of the model after k renormalization steps. Finally, we describe the expansion in the general renormalization step. Usually we are able to prove the necessary convergence estimates as we describe each part of the expansion. This has the advantage of allowing us to consider each part in isolation, without worrying about the overall structure. Unfortunately this philosophy cannot be applied to the large field estimates. For these we find it necessary to postpone the estimates of convergence until integration over the final set of fields on the unit lattice. However, using the

expected small factors arising from terms in the action with large fields, we show how convergence will eventually be obtained. We also assume estimates similar to those proven in [5] on the perturbation theory for this model.

2. Localized Kernels

In the previous paper [2], a number of operators arose from the application of renormalization transformations to the Gaussian approximation to the Higgs model. (Equation numbers from that paper will be prefixed here by I .) Exponential tails in the kernels of these approximately local operators are unavoidable. However, they are inconvenient for our analysis here, since they interfere with the decoupling of distant regions of space-time. Therefore, we introduce localized versions in which the tail has been cut off at a sufficiently large distance. In addition, for operators depending on an external gauge field, the dependence will be reduced to a bounded region. The use of the localized operators instead of the exact ones will introduce small error terms that are easily controlled.

Let us consider gauge field operators first. The minimizer H_k maps unit lattice bond fields to η -lattice bond fields, $\eta = L^{-k}$. The configuration $A = H_k B$ minimizes the Landau gauge η -lattice gauge field action under the constraint $Q_k A = B$. We define H_k starting with a large but fixed torus $T_{0,\varepsilon}$ of size $O(\varepsilon^{-1})$, say, in each lattice direction. [Recall our convention that subscripts on tori T or T_0 indicate the lattice spacing (in this case ε); superscripts (k) indicate the number of times the initial lattice has been decimated.] This avoids spurious dependence on the lattice T_ε on which we put our model, and so simplifies the infinite volume limit. (Alternatively we could take the limit $T_{0,\varepsilon} \nearrow \mathbb{Z}^d$.) Construct a translation invariant localization function ζ_k such that

$$\zeta_k(b, b') = \begin{cases} 0, & \text{if } \text{dist}(b, b') \geq \frac{1}{8}r(e_k), \\ 1, & \text{if } \text{dist}(b, b') \leq \frac{1}{16}r(e_k), \end{cases} \tag{2.1}$$

and such that ζ_k is a smooth function of b . Here $b \in T_\eta$, $b' \in T_1^{(k)}$, and

$$e_k = (L^k \varepsilon)^{(4-d)/2} e, \quad \lambda_k = (L^k \varepsilon)^{4-d} \lambda, \tag{2.2}$$

$$r(e_k) = |\log e_k^{-1}|^r, \quad r > 1. \tag{2.3}$$

Then the localized version of H_k has a kernel

$$H_{k,\text{loc}}(b, b') = \zeta_k(b, b') H_k(b, b'). \tag{2.4}$$

There is no ambiguity because ζ_k permits a sampling only of b near b' , relative to the size of $T_{0,\eta}$; H_k is also translation invariant. Since ζ_k is smooth, $H_{k,\text{loc}}$ inherits the regularity and decay properties of H_k , see (I.7.2.2). So we have

$$|H_{k,\text{loc}}(b, b')| \leq c e^{-c \text{dist}(b, b')}, \tag{2.5}$$

$$H_{k,\text{loc}}(b, b') = 0 \quad \text{for } \text{dist}(b, b') \geq \frac{1}{8}r(e_k), \tag{2.6}$$

$$|H_{k,\text{loc}}(b, b') - H_k(b, b')| \leq e^{-cr(e_k)} e^{-c \text{dist}(b, b')}, \tag{2.7}$$

and similarly for $\partial H_{k,\text{loc}}$, $\partial^* H_{k,\text{loc}}$, and for Hölder derivatives of $H_{k,\text{loc}}$ of order less than two.

Next we consider $C^{(k)}$, the covariance of the k -th step gauge field. This is defined on the unit lattice $T_{0,1}^{(k)}$. First define

$$\tilde{C}^{(k)}(b_1, b_2) = \begin{cases} C^{(k)}(b_1, b_2), & \text{if } \text{dist}(b_1, b_2) \leq \frac{1}{4}r(e_k), \\ 0, & \text{otherwise,} \end{cases} \quad (2.8)$$

and extend by translation invariance to $T_1^{(k)}$. Then put

$$C_{\text{loc}}^{(k)} = (I - Q^{s*}Q)\tilde{C}^{(k)}(I - Q^*Q^s); \quad (2.9)$$

this insures that $C_{\text{loc}}^{(k)}$, like $C^{(k)}$, satisfies the constraints from the renormalization transformation and from the axial gauge conditions:

$$QC_{\text{loc}}^{(k)} = C_{\text{loc}}^{(k)}Q^* = 0, \quad (2.10)$$

$$\sum_{b \in T_{y,x}} C_{\text{loc}}^{(k)}(b, b_0) = \sum_{b \in T_{y,x}} C_{\text{loc}}^{(k)}(b_0, b) = 0. \quad (2.11)$$

(See [2, Chap. 2] for definitions of the block averaging operators Q , Q^s , and Q^e .) We have estimates analogous to (2.5)–(2.7) for $C_{\text{loc}}^{(k)}$.

From $C_{\text{loc}}^{(k)}$ and $H_{k,\text{loc}}$ we construct a localized η -lattice gauge field propagator analogous to \mathcal{D}_k :

$$\mathcal{D}_{k,\text{loc}} = \sum_{j=0}^{k-1} H_{j,\text{loc}}^{L^j\eta} C_{\text{loc}}^{(j),L^j\eta} H_{i,\text{loc}}^{*L^j} \equiv \sum_{j=0}^{k-1} G_{\text{loc}}^{(j),\eta}. \quad (2.12)$$

Superscripts $L^j\eta$, η , etc. indicate the lattice spacing for operators rescaled to nonstandard lattices. This propagator derives its regularity and decay from that of $C_{\text{loc}}^{(k)}$ and $H_{k,\text{loc}}$. Thus

$$|(\mathcal{D}_{k,\text{loc}}f)(b)| \leq ce^{-c \text{dist}(\text{suppt } f, b)} \|f\|_{\infty}, \quad (2.13)$$

and similarly for derivatives of $\mathcal{D}_{k,\text{loc}}$ and Hölder derivatives of order less than 2. Furthermore,

$$\mathcal{D}_{k,\text{loc}}(b_1, b_2) = 0 \quad \text{for } \text{dist}(b_1, b_2) \geq \frac{1}{2}r(e_k),$$

and $\mathcal{D}_{k,\text{loc}}$ is close to \mathcal{D}_k , see (5.4.3) below.

The operator σ_k gives the quadratic form for the k -th-step field strengths $f^{(k)}(p) = (ie_k)^{-1} \log u(p)$. As before we construct a localized operator on $T_1^{(k)}$ from σ_k on $T_{0,1}^{(k)}$:

$$\sigma_{k,\text{loc}}(p_1, p_2) = \begin{cases} \sigma_k(p_1, p_2), & \text{if } \text{dist}(p_1, p_2) \leq \frac{1}{2L}r(e_{k-1}), \\ 0, & \text{otherwise,} \end{cases} \quad (2.14)$$

where p_1, p_2 are plaquettes of $T_1^{(k)}$. Recall from [2] that

$$\sigma_k = Q_k^e(I - \partial G_{k,Ax} \partial^*) Q_k^{e*} = Q_k^e(I - \partial \mathcal{D}_k \partial^*) Q_k^{e*}, \quad (2.15)$$

the second equality following from the change of gauge, (I.5.2.6). Writing \mathcal{D}_k in hierarchical form as in (2.12) and using the regularity of H_j , $0 \leq j < k$, we see that

$$|\sigma_k(p_1, p_2)| \leq ce^{-c \text{dist}(p_1, p_2)} \quad \text{for } \text{dist}(p_1, p_2) \geq c. \quad (2.16)$$

[The rapid decay of the terms with small j compensates for the scaling factors $(L^j\eta)^{-1}$.] For close p_1, p_2 the kernel of σ_k can be large, of the order of η^{-2} . However,

we shall only encounter situations where $f^{(k)}(p_2) = (\partial A)(p_2)$ for p_2 near p_1 . Then we prove that

$$(\sigma_k f^{(k)})(p_1) \leq \|f^{(k)}\|_\infty \quad (2.17)$$

as follows. Write $f^{(k)} = \partial \square A + f'$, where \square is the characteristic function of a neighborhood of p_2 . The distant part $(\sigma_k f')(p_1)$ is easily estimated by $\|f^{(k)}\|_\infty$ by (2.16). The near part is similarly bounded since σ_k is a bounded operator on curls [2]. It was also shown in [2] that σ_k is bounded from below. In view of (2.16) we have that

$$|\sigma_{k,\text{loc}}(p_1, p_2) - \sigma_k(p_1, p_2)| \leq ce^{-cr(e_k)} e^{-c \text{dist}(p_1, p_2)}, \quad (2.18)$$

so that (2.16), (2.17) hold for $\sigma_{k,\text{loc}}$, and

$$\sigma_{k,\text{loc}} \geq c > 0 \quad (2.19)$$

as well.

Another important kernel is the one generating the gauge transformation:

$$(Q_k^{*} - \mathcal{D}_k \partial^* Q_k^{e*} \partial) A = H_k A + \partial C_k A. \quad (2.20)$$

The kernel C_k is constructed from the basic gauge transformation D_k which changes the minimizer from axial to Landau gauge (I.5.1.1):

$$H_{k, \text{Ax}} B = H_k B + \partial D_k B. \quad (2.21)$$

By changing gauge in each term in the hierarchical sum defining \mathcal{D}_k and applying (I.5.3.1), we obtain

$$C_k = D_k + \sum_{j=0}^{k-1} D_j^{L_j \eta} C^{(j), L_j \eta} H_j^{* L_j \eta} \partial^* Q_k^{e*} \partial. \quad (2.22)$$

The kernels of all these operators have an exponential decay on their respective length scales; for D_k the required estimate is (I.7.2.4). The sum over j is not well controlled for close points; this will not be important for us. For more distant points, however, the rapid decay of terms with small j controls the scalings and the sum over j to yield a uniform bound

$$|C_k(x, b')| \leq ce^{-c \text{dist}(x, b')}, \quad \text{dist}(x, b') > c. \quad (2.23)$$

Here $x \in T_{0, \eta}$, $b' \in T_{0, 1}^{(k)*}$. Of course there is no uniform bound on ∂C_k . The localized version of C_k is defined using another smooth cutoff:

$$\zeta'_k(x, b') = \begin{cases} 1, & \text{for } \text{dist}(x, b') \leq \frac{1}{4} r(e_k) \\ 0, & \text{for } \text{dist}(x, b') \geq \frac{1}{2} r(e_k) \end{cases}. \quad (2.24)$$

We then construct $C_{k,\text{loc}}$ on $T_1^{(k)}$ from C_k on $T_{0, 1}^{(k)}$:

$$C_{k,\text{loc}}(x, b') = \zeta'_k(x, b') C_k(x, b'). \quad (2.25)$$

Then $C_{k,\text{loc}}$ also satisfies (2.23) and

$$|C_{k,\text{loc}}(x, b') - C_k(x, b')| \leq e^{-cr(e_k)} e^{-c \text{dist}(x, b')}. \quad (2.26)$$

In the scalar field sector, we have the η -lattice propagators $G_k(\Omega, u)$ defined on subsets $\Omega \subset T_\eta$ with Neumann boundary conditions. To localize the dependence on

u , we interpolate in a smooth fashion between operators with Neumann boundary conditions on small cubes. Let $\{\square_x\}$ be the collection of $\frac{1}{2L}r(e_{k-1})$ -cubes that can be built from cubes of size $M=O(1)$ as in [6]. Define

$$\tilde{G}_k(u; x_1, x_2) = \sum_x \lambda_x G_k(\square_x, u; x_1, x_2) \tag{2.27}$$

as a convex combination of Neumann propagators. The convex combination varies smoothly with $(x_1 + x_2)/2$; it involves at most 2^d terms and is concentrated on \square_x when $(x_1 + x_2)/2$ is near the center of \square_x . We then put

$$G_{k,loc}(u; x_1, x_2) = \zeta_k''(x_1, x_2) \tilde{G}_k(u; x_1, x_2), \tag{2.28}$$

where $\zeta_k''(x_1, x_2)$ is a smooth function of $x_1 - x_2$,

$$\zeta_k''(x_1, x_2) = \begin{cases} 0, & \text{if } |x_1 - x_2| \geq \frac{1}{4L} r(e_{k-1}) \\ 1, & \text{if } |x_1 - x_2| \leq \frac{1}{8L} r(e_{k-1}) \end{cases}. \tag{2.29}$$

The boundary conditions are always at a distance $O(r(e_k))$ from x_1, x_2 , so a straightforward application of the random walk expansion of [6] shows that

$$\|(G_{k,loc}(u)f)(x)\| \leq ce^{-c \text{dist}(\text{supp}f, x)} \|f\|_\infty, \tag{2.30}$$

$$\|(G_{k,loc}(u)f - G_k(\Omega, u)f)(x)\| \leq e^{-cr(e_k)} e^{-c \text{dist}(\text{supp}f, x)} \|f\|_\infty, \tag{2.31}$$

for $\text{dist}(x, \Omega^c) \geq O(r(e_k))$. [Each $G_k(\square_x, u)$ is close to $G_k(\Omega, u)$ for the relevant x_1, x_2 , therefore the convex combination and $G_{k,loc}$ are close also.] We assume that u is smooth in the \square_x 's entering the sum in (2.27); for (2.31) we assume smoothness throughout the subset $\Omega \subset T_\eta$. This means that in a neighborhood of each \square_x there exists an A, λ such that

$$u = \exp[ie_k \eta(A + \partial\lambda)] \quad \text{with} \quad |\partial A|, |\partial^* A| \leq O(p(e_k)). \tag{2.32}$$

Here

$$p(e_k) = |\log e_k^{-1}|^p, \quad p = O(1) \tag{2.33}$$

is our logarithmic scale for small fields. Bounds analogous to (2.30), (2.31) hold for covariant derivatives and Hölder derivatives of $G_{k,loc}(u)$ of order less than two.

We use $G_{k,loc}$ to define a localized quadratic form for scalar fields,

$$\Delta_{k,loc}(u) = a_k I - a_k^2 Q_k(u) G_{k,loc}(u) Q_k^*(u). \tag{2.34}$$

Here we have simply replaced $G_k(\Omega, u)$ with $G_{k,loc}$ in the definition of $\Delta_k(\Omega, u)$; see (I.4.6.4). Hence

$$|\Delta_{k,loc}(u; x_1, x_2) - \Delta_k(\Omega, u; x_1, x_2)| \leq e^{-cr(e_k)} e^{-c|x_1 - x_2|} \quad \text{for} \quad \text{dist}(\{x_1, x_2\}, \Omega^c) > O(r(e_k)), \tag{2.35}$$

$$|\Delta_{k,loc}(u; x_1, x_2)| \leq ce^{-c|x_1 - x_2|}, \tag{2.36}$$

$$\Delta_{k,loc}(u; x_1, x_2) = 0 \quad \text{for} \quad |x_1 - x_2| \geq \frac{1}{2L} r(e_{k-1}). \tag{2.37}$$

Again we assume u is smooth in the relevant regions; $\Delta_{k,\text{loc}}(u; x_1, x_2)$ depends on u only in an $O(r(e_k))$ -neighborhood of x_1, x_2 . Finally, in view of (2.35), the lower bound (I.7.3.2) applies to $\Delta_{k,\text{loc}}(u)$ as well. Let ϕ be supported in a region having an $r(e_k)$ neighborhood where u is smooth. Then

$$\langle \phi, \Delta_{k,\text{loc}}(u)\phi \rangle \geq c \sum_{b \in T_1^{(k)*}} |u(\langle b_-, b_+ \rangle)\phi(b_+) - \phi(b_-)|^2 - ce_k^2 p(e_k)^2 \sum_{x \in T_1^{(k)}} |\phi(x)|^2. \tag{2.38}$$

Finally, we need to construct a localized version of

$$C_A^{(k)}(\Omega, u) = [(\Delta_k(\Omega, u) + aL^{-2}Q(u)*Q(u)|_A)^{-1}], \tag{2.39}$$

the single-scale propagator for the scalar field in the k -th step. We have already replaced $\Delta_k(\Omega, u)$ with $\Delta_{k,\text{loc}}(u)$. Let us assume u is smooth in a neighborhood of A , the region for the Dirichlet boundary conditions in (2.39). We define

$$C_A^{(k)}(u) = [(\Delta_{k,\text{loc}}(u) + aL^{-2}Q(u)*Q(u)|_A)^{-1}]. \tag{2.40}$$

This is of course a nonlocal operator, but by (2.38), $C_A^{(k)}(u)^{-1}$ is bounded below and a random walk expansion as in [6] can be used to prove that

$$|C_A^{(k)}(u; x_1, x_2)| \leq ce^{-c|x_1 - x_2|}. \tag{2.41}$$

We shall actually use a convenient resummation of this expansion. The basic expansion has the form

$$C_A^{(k)}(u) = \sum_{\omega} C_{\omega}, \tag{2.42}$$

where ω is a walk on a lattice of spacing $M = O(1)$. We define the localized form of $C_A^{(k)}(u)$ to be

$$C_{A,\text{loc}}^{(k)}(u; x_1, x_2) = \sum'_{\omega} C_{\omega}(x_1, x_2), \tag{2.43}$$

where the prime indicates that only ω remaining within $\frac{1}{4}r(e_k)$ of x_1, x_2 are included. Let X be a connected union of $r(e_k)$ -cubes, and let X^0 be the cubes of X not at the boundary of X . We define

$$C_{A,X}^{(k)}(u; x_1, x_2) = \sum'_{\omega} C_{\omega}(x_1, x_2), \tag{2.44}$$

where the sums runs over walks not included in \sum' , which remain within X^0 and which intersect each cube of X^0 . Then we define

$$C_A^{(k)}(u) = C_{A,\text{loc}}^{(k)}(u) + \sum_X C_{A,X}^{(k)}(u), \tag{2.45}$$

and the convergence and locality properties of the random walk expansion imply the following facts about these operators. The local part $C_{A,\text{loc}}^{(k)}(u; x_1, x_2)$ depends only on u in an $O(r(e_k))$ neighborhood of x_1, x_2 ; it vanishes for $|x_1 - x_2| > \frac{1}{2}r(e_k)$ and is bounded as in (2.41). The operator $C_{A,X}^{(k)}(u)$ depends only on u in X . It vanishes unless both arguments are in X , and is estimated as follows:

$$|C_{A,X}^{(k)}(u; x_1, x_2)| \leq e^{-cr(e_k)|X|}. \tag{2.46}$$

Here and elsewhere, $|X|$ refers to the number of $r(e_k)$ -cubes in X , not the volume of X . This estimate can be summed over all connected sets X to show that

$$|C_{A,\text{loc}}^{(k)}(u; x_1, x_2) - C_A^{(k)}(u; x_1, x_2)| \leq e^{-cr(e_k)} e^{-c|x_1 - x_2|}. \quad (2.47)$$

If X does not intersect A^c , then $C_{A,X}^{(k)}(u)$ does not depend on A ; neither does $C_{A,\text{loc}}^{(k)}(u; x_1, x_2)$ depend on A if $\text{dist}(\{x_1, x_2\}, A^c) > \frac{1}{2}r(e_k)$. In this case we write it as

$$C_{\text{loc}}^{(k)}(u; x_1, x_2) = C_{A,\text{loc}}^{(k)}(u; x_1, x_2), \quad A \text{ large enough.} \quad (2.48)$$

Note that all operators introduced through random walk expansions of $C_A^{(k)}(u)$ or $G_k(\Omega, u)$ transform properly under gauge transformations, that is, by the difference of the gauge transformation between the points of evaluation of the kernel.

Lastly we note that the single-step covariance for the gauge field can be given a random walk expansion analogous to (2.45), with similar estimates:

$$C_A^{(k)} = C_{A,\text{loc}}^{(k)} + \sum_X C_{A,X}^{(k)}. \quad (2.49)$$

3. The First Renormalization Step

In this section we briefly and informally describe the sequence of operators performed in the first renormalization step. This will serve to orient the reader in the more detailed descriptions for the general step, and it will motivate the inductive hypothesis for the general step. Most estimates will not be discussed here, since they are special cases of those proven for the general step. We avoid formulae in favor of verbal descriptions, except for the first few operations, which are special to the first step.

We wish to give an expansion for the partition function, or for an unnormalized expectation of an observable F . Thus we consider

$$[F] = \int \mathcal{D}u \mathcal{D}\phi e^{-S^\varepsilon(u, \phi)} F, \quad (3.1)$$

where F is a gauge-invariant function, a product of terms like $|\phi(x)|^2$, $\bar{\phi}(b_-)u(b)\phi(b_+)$, $\text{Re}(i\varepsilon^2)^{-1}(u(p)-1)$. Each such term may need to have an appropriate constant subtracted in order to obtain ε -independent bounds on the full expectation

$$\langle F \rangle = [F]/[1]. \quad (3.2)$$

These ‘‘Wick ordering’’ constants are given by perturbation theory to a low order, and will be discussed carefully in a subsequent paper on the perturbation expansions.

The action on T_ε , the ε -lattice, is

$$S^\varepsilon(u, \phi) = \sum_{p \in T_\varepsilon^{**}} \varepsilon^d \frac{1}{e^2 \varepsilon^4} [1 - \text{Re} u(p)] + \frac{1}{2} \langle \phi, -\Delta_u^\varepsilon \phi \rangle + \sum_{x \in T_\varepsilon} \varepsilon^d P(\phi(x)) + E_0 + E_1. \quad (3.3)$$

Here $-\Delta_u^\varepsilon = D_u^{\varepsilon*} D_u^\varepsilon$, and

$$P(\phi) = \lambda |\phi|^4 - \frac{1}{4} |\phi|^2 + \frac{1}{64\lambda} - \frac{1}{2} \delta m^2 |\phi|^2. \quad (3.4)$$

We have taken the bare scalar field mass [coming from the radial curvature of $P(\phi)$] equal to 1 ; other values can be achieved by scalings. We have included a mass renormalization $\delta m^2 = \delta m^2(e, \lambda, \varepsilon)$ and a vacuum energy renormalization $E_1 = E_1(e, \lambda, \varepsilon, |T_\varepsilon|)$. The constant $E_0 = E_0(\varepsilon, T_\varepsilon)$ normalizes the integral (3.1) so that

$$\lim_{\lambda, \varepsilon \rightarrow 0} [1] = 1. \tag{3.5}$$

We will be considering subsets X of the lattice $T_a^{(k)}$ obtained by decimating T_ε k -times and scaling the resulting lattice spacing to a . We denote by X^* the set of bonds with both endpoints in X ; then X^{**} denotes the set of plaquettes with all four corners in X . A superscript c denotes complement, so that $X^c = T_a^{(k)} \setminus X$, $X^{*c} = T_a^{(k)*} \setminus X^*$, etc. Thus X^{c*c} includes bonds with one or both endpoints in X .

We rescale our expressions from T_ε to the unit lattice T_1 . The scalar field is multiplied by $\varepsilon^{-(d-2)/2}$, and we have

$$[F] = \int \mathcal{D}u \mathcal{D}\phi \varrho_0(u, \phi), \tag{3.6}$$

$$\begin{aligned} \varrho_0(u, \phi) = F \exp \left[- \sum_{p \in T_1^{**}} e_0^{-2} (1 - \text{Re} u(p)) - \frac{1}{2} \langle \phi, -\Delta_u \phi \rangle \right. \\ \left. - \sum_{x \in T_1} P_0(\phi(x)) - \sum_{x \in T_1} \frac{1}{2} \delta m^2 \varepsilon^2 |\phi(x)|^2 - \mathcal{E}_0 - E_1 \right]. \end{aligned} \tag{3.7}$$

Here P_0 is the first in a sequence of scalar potentials forming the dominant term after k steps:

$$P_k(\phi) = \lambda_k |\phi|^4 - \frac{1}{4} (L^k \varepsilon)^2 |\phi|^2 + \frac{1}{64 \lambda} (L^k \varepsilon)^d. \tag{3.8}$$

We use a rescaled coupling constant

$$\lambda_k = (L^k \varepsilon)^{4-d} \lambda, \tag{3.9}$$

and since $e^2/\lambda = O(1)$ we have also $e_k^2/\lambda_k = O(1)$, by (2.2). The constant \mathcal{E}_0 includes the scaling factors,

$$\mathcal{E}_0 = E_0 - (d-2) |T_1| \log \varepsilon^{-1}. \tag{3.10}$$

Each factor ϕ in F acquires a factor $\varepsilon^{-(d-2)/2}$, but we use the same notation. Ultimately these scaling factors will be cancelled by successive rescalings back to the original scale.

We begin to compute $[F]$ by integrating over u, ϕ under constraints given by the block fields v, ψ on the L -lattice. This is the renormalization transformation, described in the previous paper. With the gauge fix $\delta_{Ax}(u)$, it takes the density $\varrho_0(u, \phi)$ to

$$\begin{aligned} \varrho_1^L(v, \psi) = \int \mathcal{D}u \mathcal{D}\phi \delta(v/Qu) \delta_{Ax}(u) F \exp \left[- \sum_p e_0^{-2} (1 - \text{Re} u(p)) \right. \\ \left. - \frac{1}{2} a L^{-2} \langle \psi - Q(u)\phi, \psi - Q(u)\phi \rangle - \frac{1}{2} \langle \phi, -\Delta_u \phi \rangle \right. \\ \left. - \sum_x P_0(\phi(x)) - \sum_x \frac{1}{2} \delta m^2 \varepsilon^2 |\phi(x)|^2 - \mathcal{E}_0 - E^{(0)} - E_1 \right]. \end{aligned} \tag{3.11}$$

Here we define

$$E^{(0)} = -|T_1^{(1)}| \log(aL^{d-2}/2\pi), \tag{3.12}$$

which normalizes the transformation so that

$$[F] = \int dv d\psi \varrho_1^L(v, \psi). \tag{3.13}$$

The first operation is a decomposition of the lattice into large and small field regions. This is accomplished by means of a partition of unity,

$$1 = \sum_{A_0^{(0)}} \zeta_{A_0^{(0)c}} \chi_{A_0^{(0)}}. \tag{3.14}$$

Here $A_0^{(0)} \subset T_1$ is the small field region. It is composed of $r(e_0)$ -cubes, in each of which the factor $\chi_{A_0^{(0)}}$ enforces the following conditions:

$$\begin{aligned} |D_u \phi| \leq p(e_0), \quad |\psi - Q(u)\phi| \leq p(e_0), \quad |\phi| \leq \lambda_0^{-1/4} p(e_0) \\ |f^{(0)}(p)| \leq p(e_0), \quad \text{where } f^{(0)}(p) = (ie_0)^{-1} \log u(p). \end{aligned} \tag{3.15}$$

The factor $\zeta_{A_0^{(0)c}}$ forces at least one of these conditions to be violated somewhere in each $r(e_0)$ -cube of $A_0^{(0)c}$.

Later in this step we will introduce sets $A_1^{(0)}, A_2^{(0)}$, etc., which are obtained from $A_0^{(0)}$ either by deleting $r(e_0)$ -cubes at the boundary of $A_0^{(0)}$, or by deleting $r(e_0)$ -cubes covering regions with “irrelevant” terms from the expansions. (These are terms bounded by a high power of rescaled coupling constants.) In the k -th step we will introduce analogous small field sets $A_0^{(k)}, A_1^{(k)}$, etc.

In the previous paper, we worked with the basic quadratic form $\langle \partial A, \partial A \rangle$. This is obtained now by expanding the Wilson action in powers of e_0 . In $A_0^{(0)**}$ we have small $f^{(0)}$, so we write

$$\begin{aligned} e_0^{-2} [1 - \text{Re} u(p)] &= \frac{1}{2} f^{(0)}(p)^2 + \sum_{n=2}^{\bar{n}/2} (-1)^n \frac{e_0^{2n-2} (f^{(0)}(p))^{2n}}{(2n)!} + W_0(p) \\ &= \frac{1}{2} f^{(0)}(p)^2 + V_0(p) + W_0(p). \end{aligned} \tag{3.16}$$

We consider the expansion up to order \bar{n} in e_0 explicitly, the remainder is called “irrelevant” because it is bounded by $ce_0^{\bar{n}} p(e_0)^{\bar{n}+2} \leq ee^{d+1}$ for \bar{n} large enough. The first term, summed over $A_0^{(0)**}$, gives rise to the quadratic form $\frac{1}{2} \langle A_0^{(0)**} f^{(0)}, A_0^{(0)**} f^{(0)} \rangle$. (We use A^{**} to denote the set of plaquettes with all four corners in A ; A^* denotes the bonds with both endpoints in A . The same symbols are used for the corresponding characteristic functions.) The low order terms in e_0 are new interaction vertices.

For factors $(ie\varepsilon^2)^{-1} (u(p) - 1) = (ie_0)^{-1} \varepsilon^{-d/2} (u(p) - 1)$ in the observable $F, p \in A_0^{(0)**}$, we expand:

$$\begin{aligned} (ie\varepsilon)^{-1} \varepsilon^{-d/2} (u(p) - 1) &= \sum_{n=1}^{\infty} \varepsilon^{-d/2} \frac{(ie_0)^{n-1}}{n!} (f_0(p))^n \\ &= F_{\text{rel}}(p) + F_{\text{irr}}(p). \end{aligned} \tag{3.17}$$

The first three terms are relevant (for observables this means they do not go to zero with ε .) The others are included in $F_{\text{irr}}(p)$. (Our use of the terms “relevant” and

“irrelevant” is different from standard renormalization group language.) We sum over these two terms for $p \in \pi(F) \cap A_0^{(0)**}$, where $\pi(F)$ is the set of plaquettes having factors $\text{Re}(ie\varepsilon^2)^{-1}(u(p)-1)$ in F (with multiplicity). Denote by $\beta(F)$, $\xi(F)$ the bond, sites having factors $:\bar{\phi}(b_-)u(b)\phi(b_+)$; or $:\phi(x)^2$: in F . The result is the following expansion for F :

$$F = \sum_{S_\pi \subset \pi(F) \cap A_0^{(0)**}} \prod_{p \in S_\pi} F_{\text{irr}}(p) \prod_{p \in S_\pi^c} F_{\text{rel}}(p) \prod_{p \in \pi(F) \setminus A_0^{(0)**}} \\ \times \text{Re}(ie\varepsilon^2)^{-1}(u(p)-1) \prod_{b \in \beta(F)} :\bar{\phi}(b_-)u(b)\phi(b_+): \prod_{x \in \xi(F)} :|\phi(x)|^2:. \quad (3.18)$$

The irrelevant part of the gauge field action is Mayer-expanded:

$$\exp \left[- \sum_{p \in A_0^{(0)**}} W_0(p) \right] = \sum_{S_p \subset A_0^{(0)**}} \prod_{p \in S_p} (e^{-W_0(p)} - 1). \quad (3.19)$$

We group together large-field regions and regions with irrelevant terms. Anticipating the structure of the induction, we define $A_3^{(-1)c}$ as the union of $r(e_0)$ -cubes covering $A_0^{(0)c}$ and all plaquettes in S_p or S_π . We divide $A_3^{(-1)c}$ into connected components $\{X_\omega\}$, and define

$$g_0(X_\omega) = \sum_{S_\pi \subset \pi(F) \cap A_0^{(0)**} \cap X_\omega^{**}} \sum_{S_p \subset A_0^{(0)**} \cap X_\omega^{**}} \prod_{p \in S_\pi} \\ \times F_{\text{irr}}(p) \prod_{p \in S_\pi^c \cap X_\omega^{**}} F_{\text{rel}}(p) \prod_{p \in (\pi(F) \cap X_\omega^{**}) \setminus A_0^{(0)**}} \\ \times \text{Re}(ie\varepsilon^2)^{-1}(u(p)-1) \prod_{b \in \beta(F) \cap X_\omega^*} :\bar{\phi}(b_-)u(b)\phi(b_+): \\ \times \prod_{X \in \xi(F) \cap X_\omega} :|\phi(x)|^2: \prod_{p \in S_p} (e^{-W_0(p)} - 1) \\ \times \exp \left[- \sum_{p \in X_\omega^{**} \setminus A_0^{(0)**}} \frac{1}{e_0^2} (1 - \text{Re}u(p)) \right. \\ \left. - \sum_{x \in X_\omega \setminus A_0^{(0)}} (P_0(\phi(x)) + \frac{1}{2} \delta m^2 \varepsilon^2 |\phi(x)|^2 + E_1(x)) \right]. \quad (3.20)$$

We have written $E_1 = \sum_{x \in T_1} E_1(x)$, $E_1(x)$ defined by fixing one vertex at x for each diagram defining E_1 .

The remaining $r(e_0)$ -cubes covering the support of F are divided into connected components $\{X_\sigma\}$, and we put

$$F_{0,\text{loc}}(X_\sigma) = \prod_{p \in X_\sigma^* \cap \pi(F)} F_{\text{rel}}(p) \prod_{b \in X_\sigma^* \cap \beta(F)} :\phi(b_-)u(b)\phi(b_+): \\ \times \prod_{x \in X_\sigma \cap \xi(F)} :|\phi(x)|^2:. \quad (3.21)$$

Our density now takes the following form:

$$\varrho_1^L(v, \psi) = \sum_{A_0^{(0)}} \sum_{\{X_\omega\}} \int \mathcal{D}u \mathcal{D}\phi \delta(v/Qu) \delta_{\text{Ax}}(u) \zeta_{A_0^{(0)}} \chi_{A_0^{(0)}} \\ \times \prod_{\omega} g_0(X_\omega) \prod_{\sigma} F_{0,\text{loc}}(X_\sigma) \exp \left[-\frac{1}{2} \langle A_0^{(0)**} f^{(0)}, A_0^{(0)**} f^{(0)} \rangle \right. \\ \left. -\frac{1}{2} aL^{-2} \langle \psi - Q(u)\phi, \psi - Q(u)\phi \rangle -\frac{1}{2} \langle \phi, (-\Delta_u)\phi \rangle \right. \\ \left. - \mathcal{P}_{0,\text{loc}}(A_0^{(0)}) - \mathcal{E}_0 - E^{(0)} \right], \quad (3.22)$$

where the basic interaction terms have been included in

$$\mathcal{P}_{0,\text{loc}}(A_0^{(0)}) = \sum_{x \in A_0^{(0)}} (P_0(\phi(x)) + \frac{1}{2} \delta m^2 \varepsilon^2 |\phi(x)|^2 + E_1(x)) + \sum_{p \in A_0^{(0)**}} V_0(p). \quad (3.23)$$

These expressions have a form similar to that of the inductive hypothesis for the general renormalization step, introduced in Sect. 4.1. Nevertheless, we continue in an informal fashion with the first step, in order to outline the conceptual ideas whose details are treated in the general case of the next chapter.

We begin with a translation of the gauge field which takes the block field v out of the δ -functions of the renormalization transformation. Thus we put

$$u_b = \begin{cases} u'_b u_{b'}, & \text{if } b \in B^s(b') \cap A_1^{(0)*}, \\ u'_b, & \text{otherwise.} \end{cases} \\ \equiv u'_b (A_1^{(0)*} Q^{s*} v)_b, \quad (3.24)$$

where the prefactor $A_1^{(0)*}$ indicates that what follows is present only for $b \in A_1^{(0)*}$. From the restrictions on the fields, we have that $u'_b = e^{ie_0 A_b}$, with $|A'_b| \leq cp(e_0)$ in $A_1^{(0)*}$. The axial gauge δ -functions are invariant under this translation. In $A_1^{(0)}$, $\delta(v/Q u)$ becomes proportional to $\delta(QA')$.

In the general step, a gauge transformation is needed at this point. However, it is unnecessary here.

The quadratic form for the gauge field in $A_1^{(0)**}$ is

$$\langle A_1^{(0)**} f^{(0)}, A_1^{(0)**} f^{(0)} \rangle \\ = \langle A_1^{(0)**} (\partial A' + L^{-2} Q^{e*} f), A_1^{(0)**} (\partial A' + L^{-2} Q^{e*} f) \rangle, \quad (3.25)$$

where

$$f(p) = (ie_0)^{-1} \log v(p). \quad (3.26)$$

A second translation is needed to remove the term linear in A' . We put

$$A' = A^{(0)} - A_4^{(0)*} L^{-2} C_{\text{loc}}^{(0)} \partial^* Q^{e*} f, \quad (3.27)$$

which does not precisely eliminate the linear term. However, it is local, and away from $\partial A_4^{(0)}$ the linear term is extremely small. If we neglect terms at $\partial A_4^{(0)}$ and localized terms of the order of $e^{-cr(e_0)}$, we obtain the main quadratic forms for block and fluctuation fields:

$$\langle A_1^{(0)**} \partial A^{(0)}, A_1^{(0)**} \partial A^{(0)} \rangle + \langle A_5^{(0)'**} f, \sigma_{1,\text{loc}}^L A_5^{(0)'**} f \rangle.$$

The prime denotes decimation (taking the corners of blocks only); the superscript L indicates the block lattice spacing.

Let us write the background gauge field in $A_3^{(0)*}$ in terms of $A^{(0)}$. It is

$$u = (Q^{s*} v) \exp[ie_0 (A^{(0)} - A_4^{(0)*} L^{-2} C_{\text{loc}}^{(0)} \partial^* Q^{e*} f)]. \quad (3.28)$$

We wish to expand in $A^{(0)}$ in the scalar field quadratic forms, and in the observables $F_{0,\text{loc}}$, where this gauge field appears. This will give us scalar field forms that depend only on the block gauge field. Let θ_0 be the characteristic function of $A_6^{(0)*}$. We expand in $\theta_0 A^{(0)}$. For terms of zeroth order in $\theta_0 A^{(0)}$ we have

a background gauge field

$$u_1 = (A_1^{(0)*} Q^{s*v}) (A_6^{(0)*c} u^{(0)}) \exp(i e_0 A_4^{(0)*} L^{-2} C_{\text{loc}}^{(0)} \partial^* Q^{e*f}).$$

Here $u^{(0)} = \exp(i e_0 A^{(0)})$. Terms of first or higher order in $\theta_0 A^{(0)}$ will be treated as interactions.

The expansion yields for the scalar field forms

$$\begin{aligned} \frac{1}{2} a L^{-2} \langle \psi - Q(u)\phi, \psi - Q(u)\phi \rangle + \frac{1}{2} \langle \phi, -\Delta_u \phi \rangle &= \frac{1}{2} a L^{-2} \langle \psi - Q(u_1)\phi, \psi - Q(u_1)\phi \rangle \\ + \frac{1}{2} \langle \phi, -\Delta_{u_1} \phi \rangle + R^{(0)}(u_1, \theta_0 A^{(0)}) + \sum_{\square} W_1^{(0)}(\square), \end{aligned} \quad (3.29)$$

where $R^{(0)}$ contains the first \bar{n} orders in $A^{(0)}$ (or in e_0) and the higher order, irrelevant, local terms are incorporated in $W_1^{(0)}(\square)$. (These terms can always be localized to some $r(e_0)$ -cube \square .) Similarly each factor $F_{0,\text{loc}}(X_\sigma)$ is written as a sum of two terms: the first \bar{m} orders in e_0 , and the remainder which is bounded uniformly in ε for an appropriate choice of \bar{m} .

The next step is a scalar field translation to remove the term linear in ϕ in (3.29). Again we make a local translation,

$$\phi = \phi^{(0)} + a L^{-2} A_7^{(0)} C_{\text{loc}}^{(0)}(u_1) Q^*(u_1) \psi. \quad (3.30)$$

Neglecting terms at $\partial A_7^{(0)}$ and local terms [range $O(r(e_0))$] of the order of $e^{-cr(e_0)}$, we obtain the basic quadratic forms in $\phi^{(0)}$ and ψ :

$$\frac{1}{2} \langle \phi^{(0)}, (-\Delta_{u_1} + a L^{-2} Q(u_1)^* Q(u_1)) \phi^{(0)} \rangle + \frac{1}{2} \langle A_8^{(0)'} \psi, A_{1,\text{loc}}^L(u_1) A_8^{(0)'} \psi \rangle. \quad (3.31)$$

In the small field region $A_0^{(0)}$ we have small block fields:

$$\begin{aligned} |v(p) - 1| \leq c e_0 p(e_0), \quad |\psi(y)| \leq c p(e_0) \lambda_0^{-1/4}, \\ |(D_{\bar{u}_1} \psi)(b')| \leq c p(e_0), \quad b' \in A_0^{(0)*}, \end{aligned} \quad (3.32)$$

where $\bar{u}_1(b') = \bar{u}_1(\langle b'_-, b'_+ \rangle)$. We change nothing, then, by inserting a factor $\chi_{1,A_0^{(0)}}(v, \psi)$ which enforces these conditions by means of approximate characteristic functions. Similarly, it can be shown that

$$|A^{(0)}| \leq c p(e_0) \quad \text{in } A_1^{(0)*}, \quad |\phi^{(0)}| \leq c p(e_0) \quad \text{in } A_1^{(0)*}, \quad (3.33)$$

and we inset a factor $\chi'_{A_7^{(0)}}$ enforcing these bounds in $A_7^{(0)}$.

We now consider now the interaction terms $\mathcal{P}_{0,\text{loc}}(A_0^{(0)})$ and $R^{(0)}(u_1, \theta_0 A^{(0)})$, and reorganize them as follows. Vertices are restricted to $A_8^{(0)}$, and terms whose combined order in e and $\lambda^{1/2}$ is greater than \bar{n} are removed. The result is a standard set of terms which will appear at each iteration step. Here they are grouped into an interaction $V^{(0)}(A_8^{(0)}, u_1, A^{(0)}, \phi^{(0)})$, a polynomial in $A^{(0)}$ and $\phi^{(0)}$. All other terms are either localized near $A_8^{(0)c}$ or else are of high order in couplings. The other terms are written as $\sum_{\square} W_3^{(0)}(\square)$, each term localized at an $r(e_0)$ -cube, and we have a bound

$$|W_3^{(0)}(\square)| \leq e^{\bar{n}\beta} (\varepsilon/\varepsilon_0)^\kappa. \quad (3.34)$$

Here $\beta > 0$ is a fixed small power, $\kappa > d$ is a fixed large power, and ε_0 is the lattice spacing to terminate the induction.

$$\varepsilon_0 = \varepsilon^\beta \min\{1, (8\lambda/e^2)^{1/2}\}. \quad (3.35)$$

The bound (3.34) is sufficient for a rough treatment of such terms, since $(L^k \varepsilon / \varepsilon_0)^k$ is summable on k even with an entropy factor $(L^k \varepsilon / \varepsilon_0)^{-d}$. Terms satisfying such bounds are called irrelevant.

We avoid any further consideration of the irrelevant terms by Mayer-expanding them as in (3.19). This includes terms $W_1^{(0)}(\square)$, $W_3^{(0)}(\square)$, as well as the small terms neglected in obtaining the quadratic forms for $A^{(0)}$, f , $\phi^{(0)}$, and ψ . Grouping all irrelevant terms localized in \square into $W_4^{(0)}(\square)$, we write

$$\exp\left(-\sum_{\square} W_4^{(0)}(\square)\right) = \sum_{S_4} \prod_{\square \in S_4} (\exp(-W_4^{(0)}(\square)) - 1). \tag{3.36}$$

In a similar fashion we break off the low order terms in the observables, and we obtain a sum of terms, depending on whether the relevant or irrelevant parts of $F_{k, \text{loc}}(X_\sigma)$ are chosen. (For observable terms, irrelevant means bounded independently of ε .)

We then avoid regions with irrelevant terms $\exp(-W_4^{(0)}(\square)) - 1$ or from $F_{k, \text{loc}}(X_\sigma)$. Subtract from $A_8^{(0)}$ all such regions; call the result $\tilde{A}_8^{(0)}$ and define $A_9^{(0)}$ by deleting an $r(e_0)$ -collar from it.

The original characteristic functions $\chi_{A_8^{(0)}}$ are inconvenient for our subsequent analysis because they couple block and fluctuation fields. We remove them, relying only on $\chi_{A_1, A_9^{(0)}}$ and $\chi'_{A_9^{(0)}}$ for restrictions. This means we expand each characteristic function as $\chi = 1 - \chi^c$. We obtain a sum of regions $\tilde{A}_9^{(0)}$ which contains only 1-terms:

$$\chi_{A_9^{(0)}} = \sum_{\tilde{A}_9^{(0)}} \zeta'_{\tilde{A}_9^{(0)c}}. \tag{3.37}$$

Here the function $\zeta'_{\tilde{A}_9^{(0)c}}$ forces some field to be large (χ^c) in each $r(e_0)$ -cube of $A_9^{(0)} \setminus \tilde{A}_9^{(0)}$. Then $A_{10}^{(0)}$ is defined by deleting a collar from $\tilde{A}_9^{(0)}$.

We now are prepared to calculate the integral over $\phi^{(0)}$, $A^{(0)}$ in $A_{10}^{(0)}$. We write the integrals there as normalized Gaussian integrals with conditioning at the boundary of $A_{10}^{(0)}$. This conditioning is given by $\phi^{(0)}$, $A^{(0)}$ in $A_{10}^{(0)c}$, and is a source of some nonlocal effects which must be dealt with. First of all the normalization factors for the Gaussian integral depend on the fields in $A_{10}^{(0)c}$. These can be written as the normalization factors without conditioning, $\mathcal{Z}_{A_{10}^{(0)c} * e}$, $\mathcal{Z}_{A_{10}^{(0)}(u_1)}$ times quadratic forms in $\phi^{(0)}|_{A_{10}^{(0)c}}$, $A^{(0)}|_{A_{10}^{(0)c}}$. These forms are nonlocal and they must be given random walk expansions. Secondly, there are cross terms between the fields in $A_{10}^{(0)}$ and in $A_{10}^{(0)c}$ in the exponent in the normalized Gaussian integral. We take care of most of these with a translation localized near the boundary of $A_{10}^{(0)}$. The residual linear terms, of the order of $e^{-cr(e_0)}$, are left (resulting in an uncentered Gaussian) and produce small effects in the cluster expansion.

The result is a small-field integral of the following form:

$$\int d\mu_{A_{10}^{(0)}}(A^{(0)'}, \phi^{(0)'}) \chi'_{A_9^{(0)}} \prod_{\sigma_1} F_{0, \text{loc}}^{\ddot{m}}(X_{\sigma_1}) \exp \left[-V^{(0)}(A_8^{(0)}, u_1, A^{(0)}, \phi^{(0)}) - \sum_X W_5^{(0)}(X) \right]. \tag{3.38}$$

Here $A^{(0)'}$, $\phi^{(0)'}$ are the translated fields, and the terms $W_5^{(0)}(X)$ come from the random walk expansion mentioned above. The complete expression for our density is of course much more complicated; we focus on this because it is the only remaining nonlocal effect. We give a cluster expansion for it now.

Without going into details, it is worth remarking that if we pull out the terms in $V^{(0)}$ which are independent of $\phi^{(0)}$, $A^{(0)}$ (call these $V_{\text{const}}^{(0)}$), all other terms are

uniformly small (bounded by a power of e_0) because of the restrictions on the fields. The cluster expansion puts the integral (3.38) in polymer form,

$$e^{-V_{\text{const}}^{(0)}(A_8^{(0)})} \sum_{\{X_\alpha\}} \prod_\alpha g_2(X_\alpha). \tag{3.39}$$

The polymer functions g_2 depend only on fields in X_α , and exhibit exponential decay in $|X_\alpha|$.

The clusters X_α intersection $A_{11}^{(0)c}$ have some dependence on $\phi^{(0)}|_{A_{10}^{(0)c}}$, $A^{(0)}|_{A_{10}^{(0)c}}$. The remaining clusters have completely decoupled from the large field region. We denote the region they cover by $A_{12}^{(0)}$. In this region we resum the cluster expansion and use perturbative expansions to calculate the effective action for v, ψ .

The resummed integral in $A_{12}^{(0)}$ is written as

$$z_F(A_{12}^{(0)}) = \frac{z_F(A_{12}^{(0)})}{z(A_{12}^{(0)})} \exp(\log z(A_{12}^{(0)})). \tag{3.40}$$

The first factor is the expectation of the portion of the observable in $A_{12}^{(0)}$ in the interacting fluctuation measure. The exponent is the effective action, which is calculated as follows. We interpolate the interaction $V^{(0)} - V_{\text{const}}^{(0)}$ with a prefactor t . At the same time we interpolate away the characteristic functions χ' in $A_{12}^{(0)}$. The perturbative part of the effective action is

$$\tilde{\mathcal{P}}_1(A_{12}^{(0)}) = \sum_{\alpha=1}^{\bar{n}} -\frac{1}{\alpha!} \frac{d^\alpha}{dt^\alpha} \log z_t(A_{12}^{(0)})|_{t=0}, \tag{3.41}$$

and the remainder involves truncated expectation values in the interacting fluctuation measure with parameter $t \in [0, 1]$. These truncated expectation values can be given a cluster expansion exhibiting their locality properties. Since they involve at least $\bar{n}+1$ interactions, the estimate on the resulting clusters is improved; there is a high power of couplings or a large field effect from a derivative of χ' . Thus the remainder is expressed as $\sum_X W_6^{(0)'}(X)$, a sum of localized, irrelevant terms.

The perturbative terms involve a set of diagrams, the propagators of which are fluctuation covariances $C_{A_{12}^{(0)}}^{(0)}$, $C_{A_{12}^{(0)}(u_1)}^{(0)}$ with Dirichlet boundary conditions on $A_{12}^{(0)}$. These nonlocal covariances are replaced with our standard localized ones, $C_{\text{loc}}^{(0)}$ and $C_{\text{loc}}^{(0)}(u_1)$, with the difference given a random walk expansion. Any term involving a covariance other than $C_{\text{loc}}^{(0)}$ or $C_{\text{loc}}^{(0)}(u_1)$ is extremely small, $O(e^{-cr(e_0)})$, and localized with an exponential decay. For simplicity we extend the range of integration of vertices to all of $A_8^{(0)}$; the difference involves only small, local terms in $A_8^{(0)} \cap A_{12}^{(0)c}$. As a result of these changes we have

$$V_{\text{const}}^{(0)}(A_8^{(0)}) + \tilde{\mathcal{P}}_1(A_{12}^{(0)}) = \mathcal{P}_{1,\text{loc}}^L(A_8^{(0)}) + \sum_X W_6^{(0)''}(X). \tag{3.41}$$

A perturbative contribution to $z_F(A_{12}^{(0)})/z(A_{12}^{(0)})$ is also extracted through integration by parts. When the order in couplings is high enough, the expectation is calculated with the cluster expansion. Nonlocal covariances in the perturbative part are replaced with local ones as above.

In a final operation, we Mayer-expand the irrelevant terms $W_6^{(0)'}$ and $W_6^{(0)''}$. The region $A_{13}^{(0)}$ is defined as the part of $A_{12}^{(0)}$ free of irrelevant terms, either from the

exponent or from the observable. All the terms associated with the connected components $\{X_\omega\}$ of the large-field region $A_{13}^{(0)c}$ are grouped into large field functions $g_1(X_\omega)$. We rescale the L -lattice of blocks to unit lattice spacing. From the block field ψ we get a contribution to the normalization energy:

$$E^{(0)'} = (d-2)(\log L) |T_1^{(1)}|, \quad (3.42)$$

and we put

$$\mathcal{E}_1 = \mathcal{E}_0 + E^{(0)} + E^{(0)'}. \quad (3.43)$$

The result is the following expression for our density:

$$\begin{aligned} \varrho_1(v, \psi) = & \sum_{\{X_\omega\}} \int \mathcal{D}u_{A_{10}^{(0)c}*} \chi_{1, A_0^{(k)'}} \prod_\omega g_1(X_\omega) \prod_\sigma F_{1, \text{loc}}(X_\sigma) Z_{A_{10}^{(0)c}*}^{(0)} Z_{A_{10}^{(0)}}^{(0)}(u_1) \\ & \times \exp\left[-\frac{1}{2} \langle A_5^{(0)'**} f^{(1)}, \sigma_{1, \text{loc}} A_5^{(0)'**} f^{(1)} \rangle \right. \\ & \left. -\frac{1}{2} \langle A_8^{(0)'} \psi, A_{1, \text{loc}}(u_1) A_8^{(0)'} \psi \rangle - \mathcal{P}_{1, \text{loc}}(A_8^{(0)}) - \mathcal{E}_1 \right]. \end{aligned} \quad (3.44)$$

We write explicitly the integral over $u^{(0)}$ in $A_{10}^{(0)c}*$ because in general, normalization factors $Z_{A_{10}^{(0)}}^{(j)}(u_k)$ will depend on $u^{(0)}$ through the background field u_k . We have also introduced the rescaled field strength $f^{(1)}(p) = (e_1)^{-1} \log v(p)$. The expression (3.44) will serve as a model for our starting point for the general step.

4. The Inductive Hypothesis

Our starting point is an expression like (3.44) which depends on fields, u, ϕ on the unit lattice. These were the fields, v, ψ on the L -lattice in the previous step, but we have rescaled and renamed them. We assume that we have already performed k renormalization transformations and expansions of the type we are about to describe. Thus the unit lattice here corresponds to the $L^k \varepsilon$ lattice if we had done no rescalings. The original lattice T_ε is now T_η , $\eta = L^{-k}$, we assume that $L^k \varepsilon < \varepsilon_0 = \min\{1, (8\lambda/e^2)^{1/2}\} e^\beta$, with $\beta > 0$ small and $e \ll 1$. Thus we are stopping the inductive expansion somewhat before either of the two lengths in the problem are reached. The length 1 comes from the curvature of the scalar potential, the length $(8\lambda/e^2)^{1/2}$ comes from the curvature of the vector field potential when ϕ is replaced by a value minimizing its potential. When $L^k \varepsilon \geq \varepsilon_0$, we apply a final cluster expansion designed to exhibit the Higgs mechanisms. This will be the subject of the next paper in the series. The expected correlation length is of order $l = \max\{1, (8\lambda/e^2)^{1/2}\}$.

Our k -step density has the form

$$\begin{aligned} \varrho_k(u, \phi) = & \sum_{\{X_\omega\}} \int \prod_{j=0}^{k-1} [\mathcal{D}u^{(j)}]_{A_{10}^{(j)c}*} \varrho'_k(u, \phi, \{X_\omega\}, \{u^{(j)}\}), \\ \varrho'_k(u, \phi, \{X_\omega\}, \{u^{(j)}\}) = & \chi_{k, A_0^{(k-1)'}} \prod_\omega g_k(X_\omega) \prod_\sigma F_{k, \text{loc}}(X_\sigma) \prod_{j=0}^{k-1} [Z_{A_{10}^{(j)c}*}^{(j)} Z_{A_{10}^{(j)}}^{(j)}(u_k)] \\ & \times \exp\left[-\frac{1}{2} \langle A_5^{(k-1)'**} f^{(k)}, \sigma_{k, \text{loc}} A_5^{(k-1)'**} f^{(k)} \rangle \right. \\ & \left. -\frac{1}{2} \langle A_8^{(k-1)'} \phi, A_{k, \text{loc}}(u_k) A_8^{(k-1)'} \phi \rangle - \mathcal{P}_{k, \text{loc}}(A_8^{(k-1)}) - \mathcal{E}_k \right]. \end{aligned} \quad (4.1)$$

If we integrate this density over the u, ϕ variables, we obtain our original unnormalized expectation $[F]$. The measure $du^{(j)}$ is the normalized measure on $U(1)$, $\int du^{(j)} = 1$.

We now explain the various elements of this formula. Each X_ω is a union of $r(e_{k-1})$ -cubes of the L^{-1} -lattice, and the X_ω 's do not overlap. Each X_ω also specifies subsets $A_x^{(j)} \cap X_\omega$ for $0 \leq j \leq k-1$, $0 \leq \alpha \leq 13$. These are unions of $r(e_j)$ -cubes of the $L^j \eta$ -lattice. These sets satisfy compatibility conditions arising from our constructions. In particular, with $A_x^{(j)c} = \bigcup_\omega (A_x^{(j)c} \cap X_\omega)$, we have $A_0^{(j)} \subset A_{13}^{(j-1)}$ for $j \geq 1$. We have covered already the case $j=0$, which is slightly different. For $j \geq 1$, $\alpha = 1, \dots, 8$, 11 the sets $A_x^{(j)}$ are determined by $A_{x-1}^{(j)}$ by subtracting collar neighborhoods of width $r(e_j)$ in the $L^j \eta$ -lattice. We have $A_x^{(j)} \subset A_{x-1}^{(j)}$. The sets $A_{12}^{(j)}, A_{13}^{(j)}$ need not lose anything from $A_{11}^{(j)}, A_{12}^{(j)}$, though they may be smaller. The sets $A_9^{(j)}, A_{10}^{(j)}$ lose a collar from $\bar{A}_9^{(j)}, \bar{A}_{10}^{(j)}$, which may be smaller than $A_9^{(j)}, A_{10}^{(j)}$. These sets will be defined below in a manner analogous to that in the first step. We define $\bar{A}_x^{(j)}$ as the set in $T_{L^{-j}}$ obtained as the union of L^j -blocks at the points of $A_x^{(j)}$. The factors $g_k(X_\omega)$ represent the effect of large fields or irrelevant interactions from all previous steps. The factors $g_k(X_\omega)$ depend on $u^{(j)}$, $0 \leq j \leq k-1$ and on u, ϕ .

The external gauge field appearing throughout the initial density is u_k . It depends on all the $u^{(j)}$ [or equivalently, the $A^{(j)} = (ie_j)^{-1} \log u^{(j)}$]; but in $\bar{A}_6^{(k-1)*}$ it simplifies to

$$u_k = (Q_k^* u) \exp(-ie_k \eta \mathcal{D}_{k, \text{loc}} \delta^* Q_k^{e*} f^{(k)}), \quad (4.2)$$

where $f^{(k)}(p) = (ie_k)^{-1} \log u(p)$. This is just a localized version of (I.4.5.4).

The form of u_k in $\bar{A}_6^{(k-1)*c}$ is quite complicated; we will see it as we construct u_{k+1} in the induction step. It is important now only to know that $u_{k,b}$ depends only on the fields $u^{(j)}$, u in a neighborhood of b of size $r(e_{k-1})/2L$ on $T_1^{(k)}$. Furthermore, the configuration is smooth in the sense that for each $j < k$ (and lattice spacing $\zeta = L^{-j}$), and for each $r(e_j)$ -cube \square in $\bar{A}_1^{(j)}$, there exists a gauge transformation $u_k \rightarrow u_k^{\zeta}$ such that

$$u_{k,b}^{\zeta} = \exp(ie_j L^{-j} A_b^{\zeta}) \quad \text{with} \quad |A_b^{\zeta}|, |(\partial^{\zeta} A^{\zeta})(p)|, |(\partial^{\zeta*} A^{\zeta})(x)| \leq cp(e_j)r(e_j) \quad (4.3)$$

in \square . In the k -th step the behavior of u_k in $\bar{A}_6^{(k-1)*c}$ matters only in operations involving the Gaussian normalization factors.

The configuration u_k on T_η^* gives a configuration \bar{u}_k on $T_1^{(k)*}$ by taking a product along the bond in $T_1^{(k)*}$, i.e.,

$$\bar{u}_{k,b} = u_k(\langle b_-, b_+ \rangle). \quad (4.4)$$

The factor $\chi_{k, A_0^{(k-1)}}$ gives restrictions on u, ϕ in $A_0^{(k-1)^\nu}$. The following are implied by the smoothed characteristic functions in $\chi_{k, A_0^{(k-1)}}$:

$$\begin{aligned} |f^{(k)}(p)| &\leq cp(e_k), & p \in A_0^{(k-1)\nu**}, \\ |(D_{\bar{u}_k} \phi)(b)| &\leq cp(e_k), & b \in A_0^{(k-1)\nu*}, \\ |\phi(x)| &\leq c\lambda_k^{-1/4} p(e_k), & \text{if } (L^{k-1}\varepsilon)^d < \lambda, \quad x \in A_0^{(k-1)^\nu}, \\ \|(\phi(x)) - (8\lambda)^{-1/2} (L^k \varepsilon)^{d-2/2}\| &\leq c(L^k \varepsilon)^{-1} p(e_k), & \text{if } (L^{k-1}\varepsilon)^d \geq \lambda, \quad x \in A_0^{(k-1)^\nu}. \end{aligned} \quad (4.5)$$

We have incorporated some rescaling factors (powers of L) and the difference between $p(e_k)$ and $p(e_{k-1})$ into the constant c .

The Gaussian normalization factors are given now, in a rescaled form.

$$Z_{A_{10}^{(j)c*c}}^{(j)} = \int \mathcal{D}A_{A_{10}^{(j)c*c}} \delta_{A_x, A_{10}^{(j)}}(A) \delta_{A_{10}^{(j)c*c}}(QA_{10}^{(j)c*c}A) \times \exp\left(-\frac{1}{2} \langle A_{10}^{(j)c*c}A, \partial^* \sigma_{j,\text{loc}}^{L^j \eta} \partial A_{10}^{(j)c*c}A \rangle - E_{k,v}^{(j)} \|A_{10}^{(j)c*c}\| \right), \quad (4.6)$$

where A lies on the $L^j\eta$ -lattice. The subscript to $\mathcal{D}A$ indicates where an A -field is integrated; the subscripts to δ_{A_x} and $\delta(QA)$ indicate which blocks have axial gauge conditions and which block bonds have conditions on QA . We have Dirichlet boundary conditions in $A_{10}^{(j)c*c}$. We have included a constant factor to take care of the scalings and make this independent of k . It is defined using

$$E_{k,v}^{(j)} = -\log \left[\frac{e_j}{2\pi} (L^j \eta)^{(d-2)/2} \right], \quad (4.7)$$

$$\|A_{10}^{(j)c*c}\| = |A_{10}^{(j)c*c}| - |A_{10}^{(j)'c*c}| (L^d - 1) - |A_{10}^{(j)c*c}|. \quad (4.8)$$

Here $\|A_{10}^{(j)c*c}\|$ is the number of free integrations in $A_{10}^{(j)c*c}$ after enforcing the δ -functions.

Similarly for the scalar field we have

$$Z_{A_{10}^{(j)}}^{(j)}(u_k) = \int \mathcal{D}\phi_{A_{10}^{(j)}} \exp\left(-\frac{1}{2} \langle A_{10}^{(j)}\phi, (A_{j,\text{loc}}^{L^j \eta}(u_k) + aL^{-2}P(u_k))A_{10}^{(j)}\phi \rangle - E_{k,s}^{(j)} |A_{10}^{(j)}|\right), \quad (4.9)$$

with

$$P(u_k) = Q(u_k) * Q(u_k), \quad (4.10)$$

$$E_{k,s}^{(j)} = -(d-2) \log L^j \eta. \quad (4.11)$$

The interactions of u, ϕ are in $\mathcal{P}_{k,\text{loc}}(A_8^{(k-1)})$. The subscript loc indicates that the terms therein couple fields no farther than $O(r(e_{k-1}))$ apart. $\mathcal{P}_{k,\text{loc}}$ is given by a perturbation expansion up to some fixed order \bar{n} , which we describe in detail in a later paper. For the present analysis, it is sufficient to describe a few basic features of $\mathcal{P}_{k,\text{loc}}$.

The gauge field propagator in $\mathcal{P}_{k,\text{loc}}$ is $\mathcal{D}_{k,\text{loc}}$ [except for some renormalization transformation vertices, where it is $\sum_{j=1}^{k-1} G_{\text{loc}}^{(j),\eta}$, see (2.12)] and the scalar field propagator is $G_{k,\text{loc}}(u_k)$. The fields u, ϕ appear in the diagrams through the η -lattice minimizers u_k and

$$\phi_k = a_k G_{k,\text{loc}}(u_k) Q_k^*(u_k) \phi, \quad (4.12)$$

$$f_k(p) = (ie_k \eta^2)^{-1} \log u_k(p). \quad (4.13)$$

Propagators and external fields are connected together at vertices which arise from an expansion of the η -lattice action. Vertices are restricted to $\bar{A}_8^{(k-1)}$; for vertices involving the gauge field the restriction is accomplished by means of a function h_k multiplying each vector field leg at the vertex. The function h_k changes smoothly from 0 to 1 in a neighborhood of $\bar{A}_8^{(k-1)*c}$.

The dominant term for the scalar field is $P_k(\phi_k)$, where

$$P_k(\phi) = \lambda_k |\phi|^4 - \frac{1}{4} (L^k \varepsilon)^2 |\phi|^2 + \frac{1}{64\lambda} (L^k \varepsilon)^d. \tag{4.14}$$

Under the restrictions in χ_k , $|P_k(\phi_k)| \leq c p(e_k)^4$. At P_k -vertices with l external legs, we have its l -th derivative

$$\begin{aligned} |P_k^{(l)}(\phi_k)| &\leq c \lambda_k^{l/4} p(e_k)^{4-l}, \quad \text{for } (L^k \varepsilon)^d \leq \lambda, \\ |P_k^{(l)}(\phi_k)| &\leq c (L^k \varepsilon)^l p(e_k)^{4-l}, \quad \text{for } (L^k \varepsilon)^d > \lambda. \end{aligned} \tag{4.15}$$

In fact all terms except $P_k(\phi_k)$ in $\mathcal{P}_{k,\text{loc}}$ obey bounds $O(e^{\beta-\alpha} (L^k \varepsilon / \varepsilon_0)^{1/4-\alpha})$, with $\alpha > 0$, small, $\alpha < \beta$. If ϕ or u obey better bounds, then there is a corresponding improvement in bounds on terms in $\mathcal{P}_{k,\text{loc}}$. We will prove a general theorem on estimates on perturbation expansions in a later paper.

Also in $\mathcal{P}_{k,\text{loc}}$ are vacuum energy and mass renormalization counterterms, properly localized. In \mathcal{E}_k we keep track of normalization energies occurring over the whole lattice. This includes the basic normalization counterterm E_0 , and factors from scaling and from normalization of renormalization transformations.

The observable is treated in a manner analogous to $\mathcal{P}_{k,\text{loc}}$. Each factor $F_{k,\text{loc}}(X_\sigma)$ is a perturbative expansion to order \bar{m} of some of the factors in F [those located in X_σ , a connected union of $r(e_{k-1})$ -cubes] with the same propagators, vertices, and external fields as before. The only difference is that the connected diagrams have at least one vertex from the observable $F(X_\sigma)$. Also the expansion is taken to a lower order in coupling constants for most F 's. The order depends on how singular F is. The sets X_σ are the connected components of the smallest union of $r(e_{k-1})$ -cubes covering all vertices of all diagrams in the expansion for $F(A_{13}^{(k-1)})$.

As in the case of the effective action, the remainders from the perturbation expansion for $F(X_\sigma)$ were included in the hole functional $g_k(X_\omega)$. In the case of the effective action remainder terms, this was possible because of a sufficiently high power of $e^{\beta-\alpha} (L^k \varepsilon / \varepsilon_0)^{1/4-\alpha}$; in the case of the observable it is possible when terms obey bounds uniform in k and ε . The bounds may depend on the numbers of fields of various kinds in F , and how close they approach one another. The perturbative terms in $F(X_\sigma)$ are considered more carefully to show that they obey bounds independent of ε . Cancellations with ‘‘Wick ordering’’ subtractions must be performed to obtain bounds which depend only on $L^k \varepsilon$. For example, as long as $(L^k \varepsilon)^d < \lambda$, we expect for the expansion arising from $|\phi(x)|^2$: a bound of the order of $(L^{k-1} \varepsilon)^2)^{2-d} + \lambda_k^{-1/2} p(e_{k-1})^2$ (with the first factor replaced by $\log L^{k-1} \varepsilon$ if $d = 2$). According to our convention, the perturbative terms in $F(X_\sigma)$ and $\mathcal{P}_{k,\text{loc}}$ are called ‘‘relevant’’ because in each case they contain insufficiently many powers of coupling constants for brute force estimation.

Finally, we assume that every factor or term in our starting expression is gauge invariant in the following senses. Gauge transformations

$$u_{k,b} \rightarrow u_{k,b} e^{-ie_k \eta(\hat{e}^{\eta\lambda})^{(b)}}, \quad \phi(x) \rightarrow \phi(x) e^{ie_k \eta \lambda(x)} \tag{4.16}$$

leave each expression invariant. We will need to use only gauge transformations supported in $\bar{A}_{13}^{(k-1)}$, so the terms in question are scalar field forms, interaction terms in $\mathcal{P}_{k,\text{loc}}$ renormalized observables $F_{k,\text{loc}}$, characteristic functions $\chi_{k,A_6^{k-1}}$,

and the normalization factors $Z_{A_{10}^{(j)}}^{(j)}(u_k)$. However, all expressions possess this invariance, even those that are buried in the inductive definition of g_k . Note that λ above is any real function on T_n , although only its values on $T_1^{(k)}$ are relevant for ϕ . We call these transformations background gauge transformations, because the integration variables $u^{(j)}$, u are not involved. In fact, transforming these fields would affect the axial gauge conditions and the gauge field renormalization transformations. These are invariant under only a very restricted class of transformations, which we describe now.

The second kind of gauge invariance is called block field gauge invariance, and is invariance under

$$\begin{aligned} u_b &\rightarrow u_b e^{-ie_k(\partial\lambda)(b)}, & \phi(x) &\rightarrow e^{ie_k\lambda(x)}, \\ u_b^{(j)} &\rightarrow u_b^{(j)} \exp[-ie_k L^j \eta(\partial^{L^j} Q_{k-j}^{\prime*} \lambda)(b)], & \text{if } b &\in A_1^{(j)*c}, \\ &u_b^{(j)} \rightarrow u_b^{(j)}, & \text{otherwise,} \end{aligned} \quad (4.17)$$

for λ a function on $T_1^{(k)}$. Here Q'_k denotes the averaging operator for real-valued functions on sites. The dependence of u_k and u and the $u^{(j)}$ is such that the above transformations induce the gauge transformation $u_{k,b} \rightarrow u_{k,b} \exp[-ie_k \eta(\partial^n Q_k^{\prime*} \lambda)]$, and thus we have invariance in the previous sense. Here, however, the variables u and $u^{(j)}$ are also transformed, but in a way that does not affect the δ -functions giving the axial gauge conditions and gauge field renormalization transformations. We remark that the first translation of $u^{(j)}$ in $A_1^{(j)*}$ accounts for the lack of a transformation there in (4.17).

In both types of gauge transformations we would have rotations of the earlier fields $\phi^{(j)}|_{A_{10}^{(j)c}}$ which are integrated over in g_k . But since the measure $d\phi^{(j)}$ is rotationally invariant, no account need be made of these rotations.

5. Renormalization and Decoupling in the General Step

5.1. Renormalization Transformation

A density of $\tilde{Q}_{k+1}^L(v, \psi)$ is obtained by applying the renormalization transformations of [1] to Q'_k as follows:

$$\begin{aligned} \tilde{Q}_{k+1}^L(v, \psi) &= \sum_{\{X_\omega\}} T_L \left[\int \prod_{j=0}^{k-1} du_{A_{10}^{(j)c}*} T_{a,L,u_k} Q'_k(u, \phi, \{X_\omega\}, \{u^{(j)}\}) \right] \\ &\equiv \sum_{\{X_\omega\}} \int \mathcal{D}u \delta(v/Qu) \int \prod_{j=0}^{k-1} \mathcal{D}u_{A_{10}^{(j)c}*} \int \mathcal{D}\phi \\ &\quad \times \exp \left[-\frac{1}{2} aL^{-2} \langle \psi - Q(u_k)\phi, \psi - Q(u_k)\phi \rangle - E^{(k)} \right] Q'_k(u, \phi, \{X_\omega\}, \{u^{(j)}\}). \end{aligned} \quad (5.1.1)$$

Here $a \approx 1$ is fixed throughout, and the normalization is

$$E^{(k)} = -\log(aL^{d-2}/2\pi). \quad (5.1.2)$$

We normalize the δ -function on $U(1)$ so that

$$\int du \delta(u) f(u) = f(1), \quad \int du = 1. \quad (5.1.3)$$

Under gauge transformations λ of $u, \phi, u^{(j)}$ that vanish on points of $T_L^{(k+1)}$, we see that the δ -functions and ϱ'_k are invariant. Since u_k also transforms by λ , we have $Q(u_k)\phi$ invariant as well. Thus no change is made if we insert the axial gauge conditions

$$\delta_{Ax}(u) = \prod_{y \in T_1^{(k)'}} \prod_{x \in B(y), x \neq y} \delta(u(\Gamma_{y,x})) \tag{5.1.4}$$

into the u -integral above.

5.2. Restrictions on the Fields

We insert a partition of unity under the integrals:

$$\begin{aligned} 1 &= \sum_{P_x \subset A_{13}^{(k-1)'}} \sum_{P_y \subset A_{13}^{(k-1)''}} \sum_{P_b \subset A_{13}^{(k-1)'''}} \sum_{P_p \subset A_{13}^{(k-1)''''}} \\ &\times \prod_{x \in P_x} \chi_x^c \prod_{x \in A_{13}^{(k-1)'}\setminus P_x} \chi_x \prod_{y \in P_y} \chi_y^c \prod_{y \in A_{13}^{(k-1)''}\setminus P_y} \chi_y \\ &\times \prod_{b \in P_b} \chi_b^c \prod_{b \in A_{13}^{(k-1)'''}\setminus P_b} \chi_b \prod_{p \in P_p} \chi_p^c \prod_{p \in A_{13}^{(k-1)''''}\setminus P_p} \chi_p, \end{aligned} \tag{5.2.1}$$

where we denote

$$\begin{aligned} \chi_x &= \begin{cases} \chi(\lambda_k p(e_k), |\phi(x)|), & \text{if } (L^k \varepsilon)^d < \lambda \\ \chi((L^k \varepsilon)^{-1} p(e_k), ||\phi| - (8\lambda)^{-1/2} (L^k \varepsilon)^{(d-2)/2}|), & \text{if } (L^k \varepsilon)^d \geq \lambda \end{cases} \\ &= 1 - \chi_x^c, \\ \chi_y &= \chi(p(e_k), |(y - Q(u_k)\phi)(y)|) = 1 - \chi_y^c, \\ \chi_b &= \chi(p(e_k), |(D_{\bar{u}_k}\phi)(b)|) = 1 - \chi_b^c, \\ \chi_p &= \chi(e_k p(e_k), |u(p) - 1|) = 1 - \chi_p^c. \end{aligned} \tag{5.2.2}$$

At each $x, y, b,$ or p where a χ^c factor is present, we expect to obtain small factors $\exp(-cp(e_k)^2) \leq e_k^\kappa$, for any κ , using the positivity of terms in the action.

The function $\chi(p, x)$ is defined as follows: We let $\chi(1, x)$ be an even, C^∞ function, equal to zero for $|x| \geq 1$, and equal to one for $|x| \leq 9/10$, and with

$$\left| \frac{d^n}{dx^n} \chi(1, x) \right| \leq c^n n^{cn} \quad \text{for all } n, x. \tag{5.2.3}$$

Then we put

$$\chi(p, x) = \chi(1, x/p). \tag{5.2.4}$$

The restrictions on $|\phi|$ are best understood by looking at the leading term in $\mathcal{P}_{k, \text{loc}}, P_k(\phi_k) \cong P_k(\phi)$, where

$$\begin{aligned} P_k(\phi) &= \lambda_k |\phi|^4 - \frac{1}{4} (L^k \varepsilon)^2 |\phi|^2 + \frac{1}{64\lambda} (L^k \varepsilon)^d \\ &= \lambda_k (|\phi| - \varrho_0)^4 + (2\lambda_k)^{1/2} (|\phi| - \varrho_0)^3 + \frac{1}{2} (L^k \varepsilon)^2 (|\phi| - \varrho_0)^2, \\ \varrho_0 &= (8\lambda)^{-1/2} (L^k \varepsilon)^{\frac{d-2}{2}}. \end{aligned} \tag{5.2.5}$$

For $(L^k \varepsilon)^d < \lambda$, the quartic term gets larger before the quadratic term, whereas for $(L^k \varepsilon)^d > \lambda$, the quadratic term gets large first. It is easy to see that for $|\phi(x)|$ in the support of χ_x^c , $P_k(\phi(x)) \geq O(p(e_k)^2)$.

We define the small field region $A_{\bullet}^{(k)}$ as the union of $r(e_k)$ -blocks, none of whose points are in $A_{13}^{(k-1) \prime c}$, P_x , or in bonds, plaquettes, or blocks in P_b , P_p , P_y . The regions $A_x^{(k)}$, $0 \leq \alpha \leq 8$ are thus determined. We resum the partition of unity to obtain

$$1 = \sum_{A_0^{(k)}} \zeta_{A_0^{(k)c}} \chi_{A_0^{(k)}}, \tag{5.2.6}$$

where

$$\begin{aligned} \zeta_{A_0^{(k)c}} &= \sum_{P_x, P_y, P_b, P_p} \prod_{x \in P_x} \chi_x^c \prod_{x \in A_{13}^{(k-1) \prime c} \setminus P_x \setminus A_0^{(k)}} \chi_x \\ &\cdots \prod_{p \in P_p} \chi_p^c \prod_{p \in A_{13}^{(k-1) \prime **} \setminus P_p \setminus A_0^{(k) **}} \chi_p, \\ \chi_{A_0^{(k)}} &= \prod_{x \in A_0^{(k)}} \chi_x \cdots \prod_{p \in A_0^{(k) **}} \chi_p. \end{aligned} \tag{5.2.7}$$

Here the sum is over subsets compatible with $A_{\bullet}^{(k)c}$, $A_{13}^{(k-1)}$.

Our density now has the form

$$\begin{aligned} \tilde{Q}_{k+1}^L(v, \psi) &= \sum_{\{X_\omega\}} \sum_{A_0^{(k)}} \int \mathcal{D}u \mathcal{D}\phi \delta_{A_x}(u) \delta(v/Qu) \int \prod_{j=0}^{k-1} \mathcal{D}u_{10}^{(j)c**} \zeta_{A_0^{(k)c}} \chi_{A_0^{(k)}} \\ &\times \chi_{k, A_0^{(k-1) \prime}} \prod_{\omega} g_k(X_\omega) \prod_{\sigma} F_{k, \text{loc}}(X_\sigma) \prod_{j=0}^{k-1} [Z_{A_{10}^{(j)c**}}^{(j)} Z_{A_{10}^{(j)}}^{(j)}(u_k)] \\ &\times \exp[-\frac{1}{2} \langle A_5^{(k-1) \prime **} f^{(k)}, \sigma_{k, \text{loc}} A_5^{(k-1) \prime **} f^{(k)} \rangle \\ &- \frac{1}{2} aL^{-2} \langle \psi - Q(u_k)\phi, \psi - Q(u_k)\phi \rangle \\ &- \frac{1}{2} \langle A_8^{(k-1) \prime} \phi, \Delta_{k, \text{loc}}(u_k) A_8^{(k-1) \prime} \phi \rangle - \mathcal{P}_{k, \text{loc}}(A_8^{(k-1)}) - \mathcal{E}_k - E^{(k)}]. \end{aligned} \tag{5.2.8}$$

Let us remark that having imposed the axial gauge conditions, we resign from all but the following restricted block field gauge invariance:

$$\begin{aligned} \psi_y &\rightarrow \psi_y e^{ie_k \lambda(y)}, & \phi_x &\rightarrow \phi_x e^{ie_k(Q^* \lambda)(x)}, \\ v_{b'} &\rightarrow v_{b'} e^{-ie_k L(\partial^T \lambda)(b')}, & u_b &\rightarrow u_b e^{-ie_k(\partial Q^* \lambda)(b)}, \\ u_b^{(j)} &\rightarrow u_b^{(j)} \exp(-ie_k Lj \eta(\partial^{Lj} Q_{k-j+1}^* \lambda)(b)), & & b \in A_1^{(j)*c} \text{ only.} \end{aligned} \tag{5.2.9}$$

These transformations represent exactly the gauge invariance that was not broken by the axial gauge conditions but was broken by the renormalization transformation. By compensating with transformations of the block fields v, ψ , we again have an invariance. This restricted gauge invariance we intend to preserve in all subsequent operations. For example, it is easily seen that the characteristic functions we have inserted are invariant. After integrating over ϕ , the ϕ -rotation becomes irrelevant and we will obtain the block field invariance at the next scale, as described in the induction hypothesis.

In an analogous fashion, ψ must be rotated when performing a general background gauge transformation. After integrating over ϕ we will obtain the invariance (4.17) at the next scale.

5.3. First Gauge Field Translation

The first translation is done in $A_1^{(k)*}$, and it removes the v -field from the δ -functions there. As in (3.24) we put

$$u = u'(A_1^{(k)*} Q^{s*} v), \quad (5.3.1)$$

cf. also (I.6.2). (The reader may wish to refer to chapter 6 of [2], where the effects of the translations are followed without the complications of the large field regions.) Using the restrictions $|u(p) - 1| \leq e_k p(e_k)$ in $A_0^{(k)**}$, and the axial gauge conditions, we obtain that

$$u'_b = e^{ie_k A'_b} \quad \text{with} \quad |A'_b| \leq cp(e_k), \quad \text{for} \quad b \in A_1^{(k)*}.$$

Let us define $f(p) = (ie_k)^{-1} \log v(p)$. The restrictions on $u(p)$ and the fact that $v = Qu$ imply that $|f(p)| \leq cp(e_k)$ for $p \in A_0^{(k)**}$. Under the translation we have

$$f^{(k)}(p) = A_1^{(k)**c}(ie_k)^{-1} \log u'(p)v(p'_0) + A_1^{(k)**}(\partial A' + L^{-2} Q^{e*} f)(p), \quad (5.3.2)$$

where p_0 is the portion of p intersecting some $B^s(b')$, $b' \in A_1^{(k)'\ast}$, and p'_0 is formed by replacing each bond in p_0 with the block bond b' in $A_1^{(k)'\ast}$, whose $B^s(b')$ contains it.

After this translation the background gauge field is

$$u_k = (\bar{A}_1^{(k)*c} Q_{k+1}^{s*} v) \exp ie_k \eta [Q_k^{s*} A_1^{(k)*} A' - \mathcal{D}_{k,\text{loc}} \partial^* Q_k^{e*} (A_1^{(k)**} (ie_k)^{-1} \log u'(p)v(p'_0) + A_1^{(k)**}(\partial A' + L^{-2} Q^{e*} f))], \quad (5.3.3)$$

for $b \in \bar{A}_6^{(k-1)*}$. The background field f_k appearing at some vertices in $\mathcal{D}_{k,\text{loc}}$ and in $F_{k,\text{loc}}$ is transformed accordingly. In $\bar{A}_2^{(k)*}$ this simplifies to

$$u_k = (Q_{k+1}^{s*}) \exp ie_k \eta [Q_k^{s*} A' - \mathcal{D}_{k,\text{loc}} \partial^* Q_k^{e*} (\partial A' + L^{-2} Q^{e*} f)], \quad (5.3.4)$$

cf. (I.6.2.3). The quadratic form $f^{(k)}$ transforms into

$$\begin{aligned} & \frac{1}{2} \langle A_5^{(k-1)'\ast} f^{(k)}, \sigma_{k,\text{loc}} A_5^{(k-1)'\ast} f^{(k)} \rangle \\ &= \frac{1}{2} \langle A_5^{(k-1)'\ast} A_1^{(k)**c}(ie_k)^{-1} \log u(p)v(p'_0) \\ & \quad + 2A_1^{(k)**} A_2^{(k)c**}(\partial A' + L^{-2} Q^{e*} f), \sigma_{k,\text{loc}} A_5^{(k-1)'\ast} A_1^{(k)**c}(ie_k)^{-1} \log u(p)v(p'_0) \rangle \\ & \quad + \frac{1}{2} \langle A_1^{(k)**}(\partial A' + L^{-2} Q^{e*} f), \sigma_{k,\text{loc}} A_1^{(k)**}(\partial A' + L^{-2} Q^{e*} f) \rangle \\ &= \mathcal{Q}_1 + \mathcal{Q}'_1. \end{aligned} \quad (5.3.5)$$

The translation affects the δ -functions as follows.

$$\delta_{\text{Ax}}(u) = \delta_{\text{Ax}}(u'), \quad \delta(v/Qu) = \delta_{A_1^{(k)'\ast c}(v/Qu)} \delta_{A_1^{(k)'\ast} \left(\frac{e_k}{2\pi} Q A' \right)}, \quad (5.3.6)$$

where

$$\delta_{A_1^{(k)'\ast} \left(\frac{e_k}{2\pi} Q A' \right)} = \prod_{b' \in A_1^{(k)'\ast}} \delta \left(\frac{e_k}{2\pi} (Q A')(b') \right). \quad (5.3.7)$$

The factor $e_k/2\pi$ arises because $du_b = (e_k/2\pi) dA_b$.

5.4. Gauge Transformation

We need to make an A' -dependent gauge transformation to put u_k into a proper form. The purpose of this operation is to keep the operators H_k appearing in u_k and in $G_{k,\text{loc}}$ in a good gauge, i.e., Landau gauge and not axial gauge. The axial gauge operator $H_{k,\text{Ax}}$ would arise more naturally in our procedure, but it does not have the necessary regularity properties. This operation is not performed in the first step, since $H_0 = H_{0,\text{Ax}} = I$.

The unlocalized form of the gauge transformation is based on the identity (2.20),

$$Q_k^{s*} A' - \mathcal{D}_k \partial^* Q_k^{e*} \partial A' = H_k A' + \partial C_k A'. \quad (5.4.1)$$

The operator C_k has an exponential decay, but C_k and ∂C_k can have local singularities which is why the term ∂C_k must be removed.

To do this in a way that does not introduce nonlocal dependence on A' , and in a way that does not change u_k in $\bar{A}_3^{(k)c*}$, we make background gauge transformations on individual terms that depend on u_k in $\bar{A}_3^{(k)c*c}$. Up to some small errors, the scalar field rotations can be removed using the rotational invariance of $\mathcal{D}\phi$, $\mathcal{D}\psi$.

The expressions $\chi_{A_0^{(k)}}$, $\chi_{k, A_0^{(k-1)}}$, $F_{k,\text{loc}}(X_\sigma)$, $Z_{A_{10}^{(k)}}(u_k)$, $\langle \psi - Q(u_k)\phi, \psi - Q(u_k)\phi \rangle$, $\langle A_8^{(k-1)'}\phi, \Delta_{k,\text{loc}}(u_k)A_8^{(k-1)'}\phi \rangle$, and $\mathcal{P}_{k,\text{loc}}(A_8^{(k-1)'})$ are the ones depending on u_k in $\bar{A}_3^{(k)c*c}$. The dependence is through some simple, localized expressions like

$$\begin{aligned} & (\bar{\psi}Q(u_k)\phi)(y), \quad G_{k,\text{loc}}(u_k; b_-, b_+)u_k(b), \\ & (\bar{\phi}Q_j(u_k)G_{j,\text{loc}}(u_k)Q_j^*(u_k)\phi)(x_1, x_2), \quad f_k(p)^4, \end{aligned}$$

or in similar expressions for the diagrams in $\mathcal{P}_{k,\text{loc}}$ or $F_{k,\text{loc}}$. The Gaussian normalization factors are written as in (4.9), and the dependence on u_k is in the operators

$$A_{j,\text{loc}}(u_k) = a_j I - a_j^2 Q_j(u_k) G_{j,\text{loc}}(u_k) Q_j^*(u_k) \quad \text{and} \quad P(u_k),$$

and we have terms of the above type. However, a slightly different procedure is applied to normalization factors; we describe it later. Let us fix a set of sites where fields ψ or ϕ sit; then the dependence on $u_{k,b}$ is only for b in some cube \square_0 enclosing all points closer than $\frac{1}{2L}r(e_{k-1})$ to the fixed sites. $\left[\text{There are at most some fixed number of propagators } G_{j,\text{loc}}(u_k) \text{ or } \mathcal{D}_{k,\text{loc}}, \text{ and each has a range less than } \frac{1}{4L}r(e_{k-1}). \right]$ Thus we can choose L such that \square_0 is a cube of size $\frac{1}{4}r(e_k)$. For the diagrams without external ϕ, ψ fields, we have to localize one vertex in a unit cube and consider the localized diagram as a separate term. We define an appropriate \square_0 containing all relevant bonds for the propagators in the localized diagram.

The gauge field appearing in any one term can be written as

$$u_k = (Q_{k+1}^{s*} v) \exp i e_k \eta [(Q_k^{s*} - \mathcal{D}_{k,\text{loc}} \partial^* Q_k^{e*} \partial) \square A' - L^{-2} \mathcal{D}_{k,\text{loc}} \partial^* Q_{k+1}^{e*} f]_b, \quad (5.4.2)$$

where \square is a $\frac{1}{2}r(e_k)$ -cube in $T_1^{(k)*}$ containing a collar neighborhood around \square_0 . We extend \square to a component of a $\frac{1}{2}r(e_k)$ -neighborhood of $A_3^{(k)*c} \setminus A_3^{(k)c*}$ for all terms such that \square_0 intersects $A_3^{(k)*c} \setminus A_3^{(k)c*}$. The values of A' outside \square do not matter

because $\mathcal{D}_{k,\text{loc}}$ has a range $\frac{1}{2L}r(e_{k-1})$, and so $u_{k,b}$ doesn't depend on them for $b \in \square_0$. We now write

$$(Q_k^{s*} - \mathcal{D}_{k,\text{loc}} \partial^* Q_k^{e*} \partial) \square A' = (Q_k^{s*} - \mathcal{D}_k Q_k^{e*} \partial) \square A' + w'_1 A'. \tag{5.4.3}$$

The kernel $w'_1 = (\mathcal{D}_k - \mathcal{D}_{k,\text{loc}}) \partial^* Q_k^{e*} \partial \square$ involves only the tails not included in the expansion (2.12). Using the regularity and exponential decay of $H_j, H_{j,\text{loc}}$, along with (2.7) and scaling properties of these kernels, we find that

$$\begin{aligned} |(\partial w'_1)(p, b')| &\leq \sum_{j=1}^{k-1} (L^j \eta)^{-1 - (d-2) - 1 + (d-2)} e^{-cr(e_j)} e^{-c \text{dist}(p, b')} \\ &\leq e^{-cr(e_k)} e^{-c \text{dist}(p, b')}, \end{aligned}$$

and similarly for $w'_1, \partial^* w'_1$. Also, w'_1 is finite ranged in the sense that $w'_1 = w'_1 \square$; we use $w'_1(b, b')$ only for b in $\square_0 \subset \square$.

The nonlocal gauge transformation (5.4.1) is now applied and we have

$$\begin{aligned} (Q_k^{s*} - \mathcal{D}_{k,\text{loc}} \partial^* Q_k^{e*} \partial) \square A' &= (H_k + \partial C_k) \square A' + w'_1 A' \\ &= H_{k,\text{loc}} A' + \partial C_k \square A' + w_1 A'. \end{aligned} \tag{5.4.4}$$

We have put $w_1 = w'_1 + H_k \square - H_{k,\text{loc}}$ and it satisfies the same bounds as w'_1 .

The background gauge transformation

$$\begin{aligned} u_{k,b} &\rightarrow u'_{k,b} = u_{k,b} \exp(-ie_k \eta (\partial \bar{A}_3^{(k)} C_k \square A')(b)), \\ \phi(x) &\rightarrow \phi(x) \exp(ie_k (\bar{A}_3^{(k)} C_k \square A')(x)), \\ \psi(y) &\rightarrow \psi(y) \exp(ie_k (\bar{A}_3^{(k)} C_k \square A')(y)), \end{aligned} \tag{5.4.5}$$

is now performed on the term localized in \square_0 . The background field becomes

$$\begin{aligned} u'_k &= (Q_{k+1}^{s*} v) \exp ie_k \eta [H_{k,\text{loc}} A' + \partial \bar{A}_3^{(k)c} C_k \square A' \\ &\quad + w_1 A' - L^{-2} \mathcal{D}_{k,\text{loc}} \partial^* Q_{k+1}^{e*} f], \end{aligned} \tag{5.4.6}$$

for $b \in \bar{A}_3^{(k)c*c}$; in $\bar{A}_3^{(k)c*}$ it is unchanged from the expression (5.3.3), obtained after the first translation. This field depends on the term considered, but we shall remove the term $w_1 A'$ from this expression later (only in $\bar{A}_3^{(k)c*c}$). Without $w_1 A'$ the field u'_k is independent of the term.

There are still the phase factors at ϕ and ψ . We define

$$\phi'(x) = \phi(x) e^{ie_k (\bar{A}_3^{(k)} C_{k,\text{loc}} A')(x)}, \quad \psi'(y) = \psi(y) e^{ie_k (\bar{A}_3^{(k)} C_{k,\text{loc}} A')(y)}.$$

By (2.26), $C_{k,\text{loc}}(x, b')$ approximates the phase factors in (5.4.5), while being independent of \square_0 . The measure $d\phi$ is rotationally invariant, so we can replace $d\phi$ with $d\phi'$ and drop the prime. We have not yet integrated over ψ , so a different density is obtained by replacing ψ' with ψ . However, the new density $\mathcal{Q}_{k+1}^L(v, \psi)$ still has the property that $\int dv d\psi \mathcal{Q}_{k+1}^L(v, \psi) = [F]$.

After these rotations, the scalar field still have small, term-dependent phase factors. The scalar fields appear as $\phi(x) \exp[ie_k (w_2 A')(x)], \psi(y) \exp[ie_k (w_2 A')(y)]$, where $w_2 = \bar{A}_3^{(k)} C_{k,\text{loc}} - \bar{A}_3^{(k)} C_k \square$ satisfies a bound

$$|w_2(x, b)| \leq \exp(-cr(e_k)) \exp(-c \text{dist}(x, b)), \tag{5.4.7}$$

[this follows from (2.26)]. Like the $w_1 A'$ terms, the $w_2 A'$ terms will be expanded out of all expressions.

The same constructions apply to $f_k(p) = (ie_k \eta^2)^{-1} \log u_k(p)$, and we end up with $f'_k(p) = (ie_k \eta^2)^{-1} \log u'_k(p)$. Of course, the term $\partial \bar{A}_3^{(k)c} C_k \square A'$ disappears. The terms involving $w_1 A'$ will be separated out later.

The constructions described above were motivated by a desire to preserve locality, and to avoid effects of the gauge transformation from reaching the hole functionals $g_k(X_\omega)$ or the large-field regions. The Gaussian normalization factors $Z_{A_{10}^{(j)}}(u_k)$ are intrinsically nonlocal objects; all regions are essentially tied together. Only after some expansions can some small terms be localized. Thus at this point we must resign from a local form of the gauge transformation. Recall from (5.3.4) that u_k has been written as

$$(\bar{A}_2^{(k)*c} u_k) (\bar{A}_2^{(k)*} Q_{k+1}^{s*} v) \exp i e_k \eta \bar{A}_2^{(k)*} (Q_k^{s*} A' - \mathcal{D}_{k, \text{loc}} \partial^* Q_k^{e*} (\partial A' + L^{-2} Q^{e*} f)). \quad (5.4.8)$$

We put

$$\begin{aligned} \bar{A}_2^{(k)*} (Q_k^{s*} - \mathcal{D}_{k, \text{loc}} \partial^* Q_k^{e*} \partial) A' &= \bar{A}_2^{(k)*} (Q_k^{s*} - \mathcal{D}_{k, \text{loc}} \partial^* Q_k^{e*} \partial) A_3^{(k)*c} A' \\ &+ (Q_k^{s*} - \mathcal{D}_{k, \text{loc}} \partial^* Q_k^{e*} \partial) A_3^{(k)*} A' = \bar{A}_2^{(k)*} (Q_k^{s*} - \mathcal{D}_{k, \text{loc}} \partial^* Q_k^{e*} \partial) A_3^{(k)*c} A' \\ &+ H_{k, \text{loc}} A_3^{(k)*} A' + \partial C_k A_3^{(k)*} A' + w_5 A'. \end{aligned} \quad (5.4.9)$$

Here

$$w_5 = [(\mathcal{D}_{k, \text{loc}} - \mathcal{D}_k) \partial^* Q_k^{e*} \partial + H_k - H_{k, \text{loc}}] A_3^{(k)*}$$

is another small, exponentially decaying kernel. We can gauge away the term $\partial C_k A_3^{(k)*} A'$, leaving us with the following background gauge field for the normalization factors:

$$\begin{aligned} (\bar{A}_2^{(k)*c} u_k) (\bar{A}_2^{(k)*} Q_{k+1}^{s*} v) \exp i e_k \eta [\bar{A}_2^{(k)*} (Q_k^{s*} - \mathcal{D}_{k, \text{loc}} \partial^* Q_k^{e*} \partial) A_3^{(k)*c} A' \\ - L^{-2} \bar{A}_2^{(k)*} \mathcal{D}_{k, \text{loc}} \partial^* Q_{k+1}^{e*} f + H_{k, \text{loc}} A_3^{(k)*} A' + w_5 A']. \end{aligned} \quad (5.4.10)$$

The term $w_5 A'$ will be removed later on; it couples A' to bonds everywhere in T_η .

5.5. Second Gauge Field Translation

We translate a second time to eliminate most of the term in \mathcal{L}_1 linear in A' . This is analogous to what is done in Sect. I.6.1. The linear term is almost equal to $\langle A_1^{(k)*} L^{-2} Q^{e*} f, Q_k^e \partial H_k A' \rangle$, since by (I.6.1.5), (2.15) we have

$$\sigma_k \partial A' = Q_k^e \partial H_k A'. \quad (5.5.1)$$

So we eliminate most of the linear term by a translation approximately equal to $A_4^{(k)*} L^{-2} C^{(k)} H_k^* \partial^* Q_{k+1}^{e*} f$.

The translation we actually use is localized, and is given by

$$A' = A^{(k)} - A_4^{(k)*} L^{-2} C_{\text{loc}}^{(k)} H_{k, \text{loc}}^* \partial^* Q_{k+1}^{e*} f. \quad (5.5.2)$$

Our construction of a $C_{\text{loc}}^{(k)}$ satisfying (2.10), (2.11) ensures that $\delta_{\text{Ax}}(A') = \delta_{\text{Ax}}(A^{(k)})$, $\delta(QA') = \delta(QA^{(k)})$. The quadratic form \mathcal{Q}'_1 becomes

$$\begin{aligned} \mathcal{Q}'_1 &= \frac{1}{2} \langle A_1^{(k)**} \partial A^{(k)}, \sigma_{k,\text{loc}} A_1^{(k)**} \partial A^{(k)} \rangle \\ &\quad + \langle A_1^{(k)**} L^{-2} Q^{e*} f - \partial A_4^{(k)*} L^{-2} C_{\text{loc}}^{(k)} H_{k,\text{loc}}^* \partial^* Q_{k+1}^{e*} f, \sigma_{k,\text{loc}} A_1^{(k)**} \partial A^{(k)} \rangle \\ &\quad + \frac{1}{2} L^{-4} \langle A_1^{(k)**} Q^{e*} f - \partial A_4^{(k)*} C_{\text{loc}}^{(k)} H_{k,\text{loc}}^* \partial^* Q_{k+1}^{e*} f, \sigma_{k,\text{loc}} (A_1^{(k)**} Q^{e*} f \\ &\quad - \partial A_4^{(k)*} C_{\text{loc}}^{(k)} H_{k,\text{loc}}^* \partial^* Q_{k+1}^{e*} f) \rangle. \end{aligned} \quad (5.5.3)$$

In the second term we isolate a term localized near $A_5^{(k)c}$ and a small term. We write

$$\mathcal{Q}'_2 = \langle A_1^{(k)**} L^{-2} Q^{e*} f, \sigma_{k,\text{loc}} A_1^{(k)**} \partial A_2^{(k)*c} A^{(k)} \rangle,$$

and in the term with $A_2^{(k)*}$ instead of $A_2^{(k)*c}$ we put

$$\begin{aligned} \sigma_{k,\text{loc}} \partial A_2^{(k)*} A^{(k)} &= \sigma_{k,\text{loc}} \partial A_2^{(k)*} \square A^{(k)} = \sigma_k \partial A_2^{(k)*} \square A^{(k)} + w'_3 A^{(k)} \\ &= Q_k^e \partial^\eta H_k A_2^{(k)*} \square A^{(k)} + w'_3 A^{(k)} \\ &= Q_k^e \partial^\eta H_{k,\text{loc}} A_2^{(k)*} A^{(k)} + w''_3 A^{(k)}. \end{aligned} \quad (5.5.4)$$

The $\frac{1}{2}r(e_k)$ -cube \square is centered near the plaquette that we are evaluating $\sigma_{k,\text{loc}} \partial A_2^{(k)*} A^{(k)}$ at. The kernels w'_3 , w''_3 have range less than $\frac{1}{2}r(e_k)$, and we have

$$|w'_3(p, b)|, \quad |w''_3(p, b)| \leq e^{-cr(e_k)}. \quad (5.5.5)$$

In (5.5.4) we have applied our usual method for obtaining formulas for localized kernels analogous to those valid for unlocalized ones [in this case, (5.5.1)]. The precise form of the error terms will be unimportant; only bounds like (5.5.5) will matter. The $f \cdot A^{(k)}$ cross-term is now

$$\begin{aligned} \mathcal{Q}'_2 &+ \langle A_1^{(k)**} L^{-2} Q_1^{e*} f, w''_3 A^{(k)} \rangle - \langle f, Q_{k+1}^e \partial^\eta H_{k,\text{loc}} C_{\text{loc}}^{(k)} L^{-2} A_4^{(k)*} \partial^* \sigma_{k,\text{loc}} \partial A^{(k)} \rangle \\ &+ \langle f, L^{-2} Q_{k+1}^e \partial^\eta H_{k,\text{loc}} A_2^{(k)*} A^{(k)} \rangle. \end{aligned} \quad (5.5.6)$$

We insert the decomposition $f = A_5^{(k)'} ** f + A_5^{(k)'} ** c f$ in the last two terms. The terms with $A_5^{(k)'} ** c f$ will be denoted by \mathcal{Q}'_2 . The first $A_5^{(k)'} **$ term involves

$$C_{\text{loc}}^{(k)} \partial^* \sigma_{k,\text{loc}} \partial = I + w''_3, \quad (5.5.7)$$

with another small, short-ranged kernel w''_3 . The term with the identity operator cancels the second $A_5^{(k)'} **$ term. Thus if we define

$$\begin{aligned} \langle f, w_3 A^{(k)} \rangle &= \langle A_1^{(k)**} L^{-2} Q_1^{e*} f, w''_3 A^{(k)} \rangle \\ &\quad - \langle A_5^{(k)'} ** L^{-2} f, Q_{k+1}^e \partial^\eta H_{k,\text{loc}} w''_3 A^{(k)} \rangle, \end{aligned} \quad (5.5.8)$$

$$\mathcal{Q}_2 = \mathcal{Q}'_2 + \mathcal{Q}_2', \quad (5.5.9)$$

then we have written the cross-term as $\mathcal{Q}_2 + \langle f, w_3 A^{(k)} \rangle$, with \mathcal{Q}_2 large but localized in a $\frac{1}{2}r(e_k)$ -neighborhood of $A_5^{(k)c}$, and with w_3 very small and having a range $\frac{1}{2}r(e_k)$.

Next we do a similar analysis on the third term in \mathcal{Q}'_1 , the term quadratic in f . The important contribution is when f is localized in $A_5^{(k)'} **$, in which case we obtain the quadratic form $\sigma_{k+1,\text{loc}}^L$ for f , plus small errors. The analysis here

parallels that of [2], Sect. 6.1, with adjustments for localized kernels. Using (5.5.4), (5.5.7), and

$$\sigma_{k,\text{loc}} \approx Q_k^e (I - \partial \mathcal{D}_{k,\text{loc}} \partial^*) Q_k^{e*}, \quad (5.5.10)$$

$$\mathcal{D}_{k,\text{loc}} + H_{k,\text{loc}} C_{\text{loc}}^{(k)} H_{k,\text{loc}}^* = \mathcal{D}_{k+1,\text{loc}}^n, \quad (5.5.11)$$

we may write the $A_5^{(k)'\ast\ast f}$ terms as

$$\frac{1}{2} \langle A_5^{(k)'\ast\ast f}, \sigma_{k+1,\text{loc}}^L A_5^{(k)'\ast\ast f} \rangle + \frac{1}{2} \langle f, w_4 f \rangle,$$

with w_4 a small, short-ranged kernel. All terms involving at least one $A_5^{(k)'\ast\ast f}$ are assembled into a quadratic form \mathcal{Q}_3 .

To summarize the effect of the two translations, we have

$$\begin{aligned} \frac{1}{2} \langle A_5^{(k-1)'\ast\ast f^{(k)}}, \sigma_{k,\text{loc}} A_5^{(k-1)'\ast\ast f^{(k)}} \rangle &= \mathcal{Q}_1 + \mathcal{Q}'_1 \\ &= \mathcal{Q}_1 + \frac{1}{2} \langle A_1^{(k)'\ast\ast} \partial A^{(k)}, \sigma_{k,\text{loc}} A_1^{(k)'\ast\ast} \partial A^{(k)} \rangle + \frac{1}{2} \langle A_5^{(k)'\ast\ast f}, \sigma_{k+1,\text{loc}}^L A_5^{(k)'\ast\ast f} \rangle \\ &\quad + \mathcal{Q}_2 + \mathcal{Q}_3 + \langle f, w_3 A^{(k)} \rangle + \frac{1}{2} \langle f, w_4 f \rangle. \end{aligned} \quad (5.5.12)$$

Here \mathcal{Q}_i , $i=1, 2, 3$, are large linear or quadratic forms, localized near $A_5^{(k)c}$. They can be written as sums over the components X_μ of $A_6^{(k)c}$, i.e., $\mathcal{Q}_i = \sum_\mu \mathcal{Q}_i(X_\mu)$. The kernels w_3, w_4 are not localized near $A_5^{(k)c}$, but are small, have a range approximately $r(e_k)$, and become independent of the $A_x^{(k)}$ in $A_6^{(k)}$, say.

After the translation, the background gauge field looks as follows. For $b \in \bar{A}_3^{(k)c\ast c}$, (5.4.6) becomes

$$\begin{aligned} u'_k &= (Q_{k+1}^{s*} v) \exp i e_k \eta [(w_1 + H_{k,\text{loc}})(A^{(k)} - A_4^{(k)*} L^{-2} C_{\text{loc}}^{(k)} H_{k,\text{loc}}^* \partial^* Q_{k+1}^{e*} f) \\ &\quad + \partial \bar{A}_3^{(k)c} C_k \square A^{(k)} - L^{-2} \mathcal{D}_{k,\text{loc}} \partial^* Q_{k+1}^{e*} f]. \end{aligned} \quad (5.5.13)$$

In $A_5^{(k)*}$ we apply (5.5.11) to obtain

$$u'_k = (Q_{k+1}^{s*} v) \exp i e_k \eta [H_{k,\text{loc}} A^{(k)} - L^{-2} \mathcal{D}_{k+1,\text{loc}}^\eta \partial^* Q_{k+1}^{e*} f + w_1 A']. \quad (5.5.14)$$

The same formula holds in $\bar{A}_5^{(k)*}$ for the gauge field in the normalization factors, except that we have w_5 instead of w_1 .

5.6. Expansion with Respect to the Fluctuation Field

Let θ_k be a function on T_η^* that equals 1 in $\bar{A}_6^{(k)*}$, 0 in $\bar{A}_5^{(k)*c}$, and changes smoothly from 0 to 1 in a neighborhood of $\bar{A}_6^{(k)*}$ of thickness $M = O(1)$. We expand most terms in our density with respect to $\theta_k H_{k,\text{loc}} A^{(k)} = \theta_k A_k$, and with respect to $w_1 A'$. This produces a number of important vertices for $A^{(k)}$, as well as irrelevant terms. We also expand in the small kernel w_2 appearing in phase factors before scalar fields. This produces only irrelevant terms. After these expansions, the background field will have the form required for the next step in $\bar{A}_6^{(k)*}$, with dependence on v only. In the next section we consider the expansion of the normalization factors.

The new background field for the action and observables is denoted \tilde{u}_{k+1} , and for $b \in \bar{A}_5^{(k)*}$ it is given by

$$\tilde{u}_{k+1} = (Q_{k+1}^{s*} v) \exp i e_k \eta [(1 - \theta_k) H_{k,\text{loc}} A^{(k)} - L^{-2} \mathcal{D}_{k+1,\text{loc}}^\eta \partial^* Q_{k+1}^{e*} f]. \quad (5.6.1)$$

In $\bar{A}_6^{(k)*}$ the $A^{(k)}$ term is absent, and except for scaling, this reduces to the form in the induction hypothesis, (4.2).

To summarize all the changes we have made on the background field, the final form of \tilde{u}_{k+1} is given by (5.3.3) in $\bar{A}_6^{(k-1)*} \cap \bar{A}_3^{(k)c}$, by (5.5.13) in $\bar{A}_3^{(k)c*c} \cap \bar{A}_5^{(k)*c}$ (but without the w_1 terms), and by (5.6.1) in $\bar{A}_5^{(k)*}$. We will not need to define \tilde{u}^{k+1} anywhere else. The corresponding background field strength is given by

$$\tilde{J}_{k+1}^\eta(p) = (ie_k \eta^2)^{-1} \log \tilde{u}_{k+1}(p). \tag{5.6.2}$$

Later we will define u_{k+1} which will be slightly changed from \tilde{u}_{k+1} in $\bar{A}_2^{(k)*} \cap \bar{A}_4^{(k)*c}$, but which will be defined everywhere because it appears in the normalization factors.

We need to check the regularity condition on \tilde{u}_{k+1} . It states that there exists a gauge transformation $\tilde{u}_{k+1} \rightarrow \tilde{u}_{k+1}^\lambda$ in each $r(e_k)$ -cube $\square \subset \bar{A}_1^{(k)}$ such that $\tilde{u}_{k+1}^\lambda = \exp(ie_k \eta A^\lambda)$ with $|A^\lambda|, |\partial A^\lambda|, |\partial^* A^\lambda| \leq cp(e_k)r(e_k)$. In $\bar{A}_1^{(k)*c}$ we have $\tilde{u}_{k+1} = u_k$, so the condition follows from the induction hypothesis (4.3).

We verify the bound by first checking it for u_k , then noticing that all the operations changing u_k into \tilde{u}_{k+1} did not destroy the bound. We use the new bounds on $u(p)$ in $\bar{A}_0^{(k)**}$ to estimate

$$u_k = (Q_k^{*}u) \exp[-ie_k \eta \mathcal{D}_{k, \text{loc}} \partial^* Q_k^{*} f^{(k)}] \tag{5.6.3}$$

with constants uniform in k . [There are bounds from the $(k-1)$ -st step, but these would not yield uniform constants.] Thus we can assume that $|f^{(k)}(p)| \leq cp(e_k)$ for $p \in A_0^{(k)**}$.

Fix $\square \subset \bar{A}_1^{(k)}$ for estimating \tilde{u}_{k+1} . In \square' [a neighborhood of \square of width $\frac{1}{2}r(e_k)$] we can write $u = \exp[ie_k(\partial\lambda + B)]$ with $|B(b)| \leq cp(e_k)r(e_k)$. We have $f^{(k)} = \partial B$ in the cube, and so

$$u_k = (Q_k^{*} e^{ie_k \partial\lambda}) \exp ie_k \eta [(Q_k^{*} - \mathcal{D}_{k, \text{loc}} \partial^* Q_k^{*} \partial) B]. \tag{5.6.4}$$

We have used the fact that $\mathcal{D}_{k, \text{loc}}$ has range $\frac{1}{2L} r(e_{k-1})$. Note that $Q_k^{*} e^{ie_k \partial\lambda}$ is a gauge transformation (generated by $Q_k^{*} \lambda$), so we can delete it from u_k .

Our desired bound now follows because by (5.4.4),

$$(Q_k^{*} - \mathcal{D}_{k, \text{loc}} \partial^* Q_k^{*} \partial) \square' B = (H_{k, \text{loc}} + \partial C_k + w_1) \square' B. \tag{5.6.5}$$

The kernels $H_{k, \text{loc}}$, w_1 and their derivatives are bounded, so $A^\lambda, \partial A^\lambda, \partial^* A^\lambda$ are finally all bounded by $cp(e_k)r(e_k)$.

The first operation we performed was a translation, which of course does not spoil the regularity of u_k . We then made a gauge transformation and removed the small kernel w_1 . The gauge transformation does not change the regularity, and $\partial w_1, \partial^* w_1$ are small, so the bounds remain valid. After another translation we removed the field $\theta_k H_{k, \text{loc}} A^{(k)}$. This field satisfies $\partial(\theta_k H_{k, \text{loc}} A^{(k)}) \leq cp(e_k)$ because $A^{(k)} \leq cp(e_k)$ and because $\partial H_{k, \text{loc}}, H_{k, \text{loc}}$, and derivatives of θ_k are bounded. Similarly $\partial^*(\theta_k H_{k, \text{loc}} A^{(k)})$, $\theta_k H_{k, \text{loc}} A^{(k)}$ are bounded by $cp(e_k)$. Thus removing $\theta_k H_{k, \text{loc}} A^{(k)}$ does not spoil the regularity, and \tilde{u}_{k+1} satisfies the regularity condition. In an analogous fashion we can check that the j -th regularity condition for $r(e_k)$ -cubes remains valid.

We now proceed with the expansions. We do not expand in the characteristic functions $\chi_{A_8^{(k)}}$, $\chi_{k, A_8^{(k+1)}}$. We take terms which came from the original expressions $F_{k, \text{loc}}$,

$$\langle \psi - Q(u_k)\phi, \psi - Q(u_k)\phi \rangle, \quad \langle A_8^{(k-1)'} \phi, \Delta_{k, \text{loc}}(u_k) A_8^{(k-1)'} \phi \rangle,$$

and $\mathcal{P}_{k, \text{loc}}(A_8^{(k-1)'})$, and write the background field as

$$\tilde{u}_{k+1} \tilde{u} = \tilde{u}_{k+1} \exp i e_k \eta [\theta_k H_{k, \text{loc}} A^{(k)} + w_1 A'] = \tilde{u}_{k+1} e^{i e_k \eta \tilde{A}}. \quad (5.6.6)$$

The scalar fields appear with factors $e^{i e_k w_2 A'}$ before then. All expressions depend on $A^{(k)}$, f only locally, or at most within a component of $A_4^{(k)c}$.

The first expansion we give is for $\tilde{u}_{k+1} \tilde{u}$ itself. (We derive some expansions for the j -th step objects for use in the next section. The expansions are modeled after ones in [7], so we will be brief.)

We have with $\zeta = L^{-j}$, \tilde{A} scaled to the ζ -lattice,

$$\tilde{u}_{k+1} \tilde{u} = \tilde{u}_{k+1} \left(1 + \sum_{n=1}^{\infty} (i e_j \zeta \tilde{A})^n / n! \right) = \tilde{u}_{k+1} (1 + F_{1, j}(\tilde{A})). \quad (5.6.7)$$

Next we expand $Q_l(\tilde{u}_{k+1} \tilde{u})$, $l=1$ or $j, j < k$,

$$\begin{aligned} (Q_l(\tilde{u}_{k+1} \tilde{u})\phi)(y) &= \sum_{x \in B_l(y)} L^{-ld} \tilde{u}_{k+1}(\Gamma_{y, x}^{(l)}) \phi(x) \left(1 + \sum_{n=1}^{\infty} (i e_j \zeta A(\Gamma_{y, x}^{(l)}))^n / n! \right) \\ &= (Q_l(\tilde{u}_{k+1})\phi)(y) + (F_{2, l}(\tilde{A}, \tilde{u}_{k+1})\phi)(y). \end{aligned} \quad (5.6.8)$$

Inserting this formula into $|\psi - Q(\tilde{u}_{k+1} \tilde{u})\phi|^2$, we obtain the vertices new to this step. For the covariant derivative on the ζ -lattice, we have

$$(\mathcal{D}_{\tilde{u}_{k+1} \tilde{u}} \phi)(b) = (D_{\tilde{u}_{k+1}} \phi)(b) + \tilde{u}_{k+1, b} F_{1, j}(\tilde{A}) \phi(b). \quad (5.6.9)$$

For the basic quadratic form with Neumann boundary conditions on Ω giving rise to $G_j(\Omega)$, we have

$$- \Delta_{\tilde{u}_{k+1}, \tilde{u}, \Omega}^N + a_j P_j(\tilde{u}_{k+1} \tilde{u}) = - \Delta_{\tilde{u}_{k+1}, \Omega}^N + a_j P_j(\tilde{u}_{k+1}) - V_j(\Omega), \quad (5.6.10)$$

where $V_j(\Omega)$ is obtained by inserting (5.6.8), (5.6.9) into the left-hand side. This leads to an expansion of the scalar field propagator in a fixed region Ω :

$$G_j(\Omega, \tilde{u}_{k+1} \tilde{u}) = G_j(\Omega, \tilde{u}_{k+1}) + G_j(\Omega, \tilde{u}_{k+1}) V_j G_j(\Omega, \tilde{u}_{k+1} \tilde{u}). \quad (5.6.11)$$

The terms in V_j are small ($O(e_j^{1-\alpha})$), bounded kernels, either alone or applied to $D_{\tilde{u}_{k+1}}$ or $D_{\tilde{u}_{k+1}}^*$. Thus the regularity properties of $G_j(\tilde{u}_{k+1})$, $D_{\tilde{u}_{k+1}} G_j(\tilde{u}_{k+1})$ imply that we can develop this expansion to any order.

We insert this into $G_{j, \text{loc}}(\tilde{u}_{k+1} \tilde{u})$ to obtain

$$\begin{aligned} G_{j, \text{loc}}(\tilde{u}_{k+1} \tilde{u})(x_1, x_2) &= G_{j, \text{loc}}(\tilde{u}_{k+1})(x_1, x_2) \\ &\quad + \zeta_j''(x_1, x_2) \sum_x \lambda_x (G_j(\square_x, \tilde{u}_{k+1}) V_j G_j(\square_x, \tilde{u}_{k+1} \tilde{u}))(x_1, x_2). \end{aligned}$$

The second term can be changed slightly by changing the set \square_x in G_j and changing the tails of the operators. The difference is w'_6 , a small ($O(e^{-cr(e_j)})$), local kernel with small covariant derivatives, and depending only locally on \tilde{u}_{k+1} , \tilde{u} . We obtain

$$G_{j, \text{loc}}(\tilde{u}_{k+1} \tilde{u}) = G_{j, \text{loc}}(\tilde{u}_{k+1}) + G_{j, \text{loc}}(\tilde{u}_{k+1}) V_j G_{j, \text{loc}}(\tilde{u}_{k+1} \tilde{u}) + w'_6.$$

This is now iterated to yield

$$G_{j,\text{loc}}(\tilde{u}_{k+1}\tilde{u}) = \sum_{n=0}^{\bar{n}} G_{j,\text{loc}}(\tilde{u}_{k+1}) [V_j G_{j,\text{loc}}(\tilde{u}_{k+1})]^n + G_{j,\text{loc}}(\tilde{u}_{k+1}) [V_j G_{j,\text{loc}}(\tilde{u}_{k+1})]^{\bar{n}} V_j G_{j,\text{loc}}(\tilde{u}_{k+1}\tilde{u}) + w_6, \quad (5.6.12)$$

with another small kernel w_6 .

These expansions are inserted in $F_{k,\text{loc}}, \langle \psi - Q(u_k)\phi, \psi - Q(u_k)\phi \rangle, \mathcal{P}_{k,\text{loc}}$, and in

$$\Delta_{k,\text{loc}}(\tilde{u}_{k+1}\tilde{u}) = a_k I - a_k^2 Q_k(\tilde{u}_{k+1}\tilde{u}) G_{k,\text{loc}}(\tilde{u}_{k+1}\tilde{u}) Q_k^*(\tilde{u}_{k+1}\tilde{u}).$$

We expand the phase factors as $1 + (e^{ie_k w_2 A'} - 1)$, the second term being extremely small. We also have the field strength expanded as

$$f'_k = \tilde{f}_{k+1}^{\bar{n}} + \partial w_1 A' + \partial(\theta_k H_{k,\text{loc}} A^{(k)}).$$

Any term involving w_1 or w_2 , and terms of higher than \bar{n} -th order in e_k are irrelevant and will be treated separately. The lower order terms are polynomials in $A^{(k)}$. All terms are local.

Let us summarize these expansions as follows. In the action we have written

$$\begin{aligned} & \frac{1}{2} a L^{-2} \langle \tilde{\psi} - Q(u'_k)\tilde{\phi}, \tilde{\psi} - Q(u'_k)\tilde{\phi} \rangle \\ & + \frac{1}{2} \langle A_8^{(k-1)'} \tilde{\phi}, \Delta_{k,\text{loc}}(u'_k) A_8^{(k-1)'} \tilde{\phi} \rangle + \mathcal{P}_{k,\text{loc}}(A_8^{(k-1)'} \tilde{\phi}, u'_k) \\ & = \frac{1}{2} a L^{-2} \langle \psi - Q(\tilde{u}_{k+1})\phi, \psi - Q(\tilde{u}_{k+1})\phi \rangle + \frac{1}{2} \langle A_8^{(k-1)'} \phi, \Delta_{k,\text{loc}}(\tilde{u}_{k+1}) A_8^{(k-1)'} \phi \rangle \\ & + \mathcal{P}_{k,\text{loc}}(A_8^{(k-1)'} \phi, \tilde{u}_{k+1}) + R^{(k)}(u_{k+1}, \theta_k H_{k,\text{loc}} A^{(k)}) + \sum_{\square} W_1^{(k)}(\square). \end{aligned} \quad (5.6.13)$$

The tildes on ϕ and ψ indicate the presence of the phase factors. Here $W_1^{(k)}(\square)$ is localized near \square , an $r(e_k)$ -cube in $A_2^{(k)}$, and $|W_1^{(k)}(\square)| \leq e_k^{\bar{n}-1-\alpha}$. (Two powers of e_k may be needed to beat the bounds on ϕ .) If we define

$$\begin{aligned} & \tilde{R}^{(k)}(\tilde{u}_{k+1}, \theta_k H_{k,\text{loc}} A^{(k)}) \\ & = \sum_{n=1}^{\bar{n}} \left[\frac{d^n}{de^n} \left(\frac{1}{2} a L^{-2} \|\psi - Q(\tilde{u}_{k+1}) e^{ie' e_k \eta \theta_k H_{k,\text{loc}} A^{(k)}} \phi\|^2 \right. \right. \\ & \quad \left. \left. + \frac{1}{2} \langle A_8^{(k-1)'} \phi, \Delta_{k,\text{loc}}(\dots) A_8^{(k-1)'} \phi \rangle + \mathcal{P}_{k,\text{loc}}(A_8^{(k-1)'} \phi, \dots) \right) \right]_{e'=0}, \end{aligned} \quad (5.6.14)$$

then $R^{(k)}$ can be obtained by replacing propagators $G_k(\square, \tilde{u}_{k+1})$ with $G_{k,\text{loc}}(\tilde{u}_{k+1})$ and eliminating extra kernels ζ_k'' explicitly (not in $G_{k,\text{loc}}(\tilde{u}_{k+1})$).

In a similar fashion we put

$$F_{k,\text{loc}}(X_\sigma) = F_{k,\text{loc}}^{(\bar{m})}(X_\sigma) + \tilde{F}_{k,\text{loc}}(X_\sigma), \quad (5.6.14)$$

where $F_{k,\text{loc}}^{(\bar{m})}(X_\sigma)$ is defined by replacing $G_k(\square, \tilde{u}_{k+1})$, ζ_k'' in

$$\sum_{m=0}^{\bar{m}} \left[\frac{d^m}{de^m} F_{k,\text{loc}}(X_\sigma, \tilde{u}_{k+1}) e^{ie' e_k \eta \theta_k H_{k,\text{loc}} A^{(k)}} \right]_{e'=0}.$$

All remainder terms are in $\tilde{F}_{k,\text{loc}}(X_\sigma)$, and we have $|\tilde{F}_{k,\text{loc}}(X_\sigma)| \leq c(F)$.

5.7. The Gaussian Normalization Factors

The expansion with respect to the fluctuation field $A^{(k)}$ must be performed with special care in the Gaussian normalization factors. Nonlocal terms naturally arise, which must then be organized properly and treated with random walk expansions. At this point, the background field in the normalization factors is given by (5.4.10), which simplifies to (5.5.14) in $\bar{A}_5^{(k)*}$. We express it as $u_{k+1}\tilde{u}$, with

$$\tilde{u} = \exp i e_k \eta [\theta_k H_{k, \text{loc}} A^{(k)} + w_5 A'] \equiv \exp i e_k \eta \tilde{A}. \quad (5.7.1)$$

In this way the field u_{k+1} is defined, and after \tilde{A} is expanded away, it remains in the normalization factors for the next renormalization transformation. One could apply (5.4.10) inductively to obtain a complete formula for u_{k+1} on the whole lattice. We will only need to use the fact that it agrees with \tilde{u}_{k+1} in $\bar{A}_5^{(k)*}$, cf. (5.6.1). The regularity conditions can be checked for u_{k+1} in the same manner as for \tilde{u}_{k+1} .

In the integral representation for the normalization factor $Z_{A_{10}^{(j)}}^{(j)}(u_k)$, rescaled to the unit lattice, we have the quadratic form (with Dirichlet boundary conditions)

$$\begin{aligned} C_{A_{10}^{(j)}}^{(j)}(u_{k+1}\tilde{u})^{-1} &= \Delta_{j, \text{loc}}(u_{k+1}\tilde{u}) + aL^{-2}P(u_{k+1}\tilde{u}) \\ &= \Delta_{j, \text{loc}}(u_{k+1}) + aL^{-2}P(u_{k+1}) - W^{(j)} \\ &= C_{A_{10}^{(j)}}^{(j)}(u_{k+1})^{-1} - W^{(j)}. \end{aligned} \quad (5.7.2)$$

All the terms from our expansions of the last section for $\Delta_{j, \text{loc}}P$, with \tilde{u}_{k+1} replaced with u_{k+1} , \tilde{u} replaced with \tilde{u} , are included in $W^{(j)}$. Thus we have

$$\begin{aligned} C_{A_{10}^{(j)}}^{(j)}(u_{k+1}\tilde{u})^{-1} &= C_{A_{10}^{(j)}}^{(j)}(u_{k+1})^{-1/2} (I - C_{A_{10}^{(j)}}^{(j)}(u_{k+1})^{1/2} W^{(j)} \\ &\quad \times C_{A_{10}^{(j)}}^{(j)}(u_{k+1})^{1/2}) C_{A_{10}^{(j)}}^{(j)}(u_{k+1})^{-1/2}, \end{aligned}$$

and so

$$Z_{A_{10}^{(j)}}^{(j)}(u_k) = Z_{A_{10}^{(j)}}^{(j)}(u_{k+1}) [\det(I - C_{A_{10}^{(j)}}^{(j)}(u_{k+1})^{1/2} W^{(j)} C_{A_{10}^{(j)}}^{(j)}(u_{k+1})^{1/2})^{-1/2}]. \quad (5.7.3)$$

Each term in $W^{(j)}$ has at least one factor e_j , and all fields are logarithmically bounded. Thus the operator after the identity is bounded by a very small number. Thus the determinant can be expanded as

$$\exp \left[\sum_{l=1}^{\infty} \frac{1}{2l} \text{tr}(C_{A_{10}^{(j)}}^{(j)}(u_{k+1}) W^{(j)})^l \right]. \quad (5.7.4)$$

The operator $C_{A_{10}^{(j)}}^{(j)}(u_{k+1})$ is our first encounter with nonlocal effects. To treat it we apply the generalized random walk expansion (2.45), modified slightly to use cubes of size $L^{k-j}r(e_k)$ in T_{L-j} . We need a similar expansion for $W^{(j)}$ into terms defined in regions X with appropriate decay estimates.

For example, we have in $F_{1, (\tilde{A}_b)}$ a series involving powers of $(w_5 A')_b$, with a nonlocal kernel w_5 . We put these powers in the form of a sum on X of quantities defined in X only. To each $b \in T_\eta$ and each collection of bonds $b_1, \dots, b_m \in T^{(k)}$ we associate in some arbitrary manner a set X (a connected union of $r(e_k)$ -cubes containing them). Then we put

$$(w_5 A')_b^m = \sum_X w_{b, m}(X), \quad (5.7.5)$$

$$w_{b, m}(X) = \sum_{(b_1 \dots b_m) \text{ compatible with } b, X} \prod_{l=1}^m (w_5(b, b_l) A'(b_l)),$$

where compatible means that b_1, \dots, b_m, b were associated to X as above. We have $|A'(b_i)| \leq cp(e_k)$, $|w_s(b, b_i)| \leq e^{-cr(e_k)} e^{-c \text{dist}(b, b_i)}$ and so

$$|w_{b,m}(X)| \leq e^{-cr(e_k)|X|}. \tag{5.7.6}$$

We may use (5.7.5) to analyze the interaction terms generated in the expansion with respect to $\tilde{A} = \theta_k H_{k,\text{loc}} A^{(k)} + w_s A'$. Treating for the moment only the high order terms, we obtain an expansion for $F_{1,j}$ in (5.6.7):

$$F_{1,j}(\tilde{A}_b^\zeta) = \sum_{n=1}^{\bar{n}} \zeta^{-1} (ie_j \zeta \tilde{A}_b^\zeta)^n / n! + \sum_X F_{1,j,b}(X),$$

with $|F_{1,j,b}(X)| \leq e_j^{\bar{n}+1-\alpha} e^{-cr(e_k)|X|^-}$. (Here $|X|^-$ is defined as $\max\{0, |X| - 1\}$, and \tilde{A} has been rescaled to the $\zeta = L^{-j}$ lattice.) In a similar fashion we can write

$$\begin{aligned} (F_{2,j}(\tilde{A}^\zeta, u_{k+1})\phi)(y) &= \sum_{x \in B_j(y)} \zeta^d u_{k+1}(\Gamma_{y,x}^{(j)}) \phi(x) \\ &\times \sum_{n=1}^{\bar{n}} (ie_j \zeta \tilde{A}(\Gamma_{y,x}^{(j)}))^n / n! + \sum_X (F_{2,j}(X)\phi)(y), \end{aligned}$$

with

$$|F_{2,j}(X; y, x)| \leq \zeta^d e_j^{\bar{n}+1-\alpha} e^{-cr(e_k)|X|^-}.$$

These expansions can be inserted into V_j , yielding

$$V_j = V_j^{(\bar{n})} + \sum_X V_j(X), \tag{5.7.7}$$

the first term containing the expansions to order \bar{n} in e_j , the second containing the remaining terms. Both terms involve small, bounded kernels (of order $e_j^{1-\alpha}$ for $V^{(\bar{n})}$, of order $e_j^{\bar{n}+1-\alpha} e^{-cr(e_k)|X|^-}$ for $V(X)$), alone or applied to $D_{u_{k+1}}$ or $D_{u_{k+1}}^*$.

Next we examine the propagators, and expand in V_j to all orders:

$$G_j(\Omega, u_{k+1} \tilde{u}) = G_j(\Omega, u_{k+1}) + \sum_{n=1}^{\bullet} G_j(\Omega, u_{k+1}) [V_j G_j(\Omega, u_{k+1})]^n.$$

Thus we have

$$\begin{aligned} G_{j,\text{loc}}(u_{k+1} \tilde{u})(x_1, x_2) &= G_{j,\text{loc}}(u_{k+1})(x_1, x_2) + \zeta_j''(x_1, x_2) \sum_{n=1}^{\infty} (G_j(\square(x_1, x_2), u_{k+1}) \\ &\times [V_j G_j(\square(x_1, x_2), u_{k+1})]^n)(x_1, x_2). \end{aligned}$$

We insert the expansion for V_j into this formula, and insert expansions for $Q_j(u_{k+1} \tilde{u})$, $Q(u_{k+1} \tilde{u})$, $G_{j,\text{loc}}(u_{k+1} \tilde{u})$ into $\Delta_{j,\text{loc}}(u_{k+1} \tilde{u}) + aL^{-2}P(u_{k+1} \tilde{u})$. Terms whose order in e_j (or equivalently in \tilde{A}) is between 1 and \bar{n} are considered as part of $-W^{(j,\bar{n})}$. Terms of higher order, or involving $F_{2,j}(X)$ or $V_j(X)$ are grouped into an expansion $\sum_X -W^{(j)}(X)$, with $|W^{(j)}(X; x_1, x_2)| \leq e_j^{\bar{n}+1-\alpha} e^{-cr(e_k)|X|^-}$. Thus we have written the interaction term in (5.7.2) as

$$W^{(j)} = W^{(j,\bar{n})} + \sum_X W^{(j)}(X).$$

The lower order terms need to be resummed by gathering terms with different j into a perturbative expression. This is because $e^{-cr(e_k)}$ is not small enough to

compensate for having only a few powers of e_j . We must replace $G_j(\square(x_1, x_2), u_{k+1})$ with $G_j(\Omega, u_{k+1})$ for some fixed Ω . This is accomplished with a random walk expansion for $G_j(\Omega, u)$. Such an expansion is given in [6]. It takes the form

$$G_j(\Omega, u) = \sum_{\omega} G(\omega),$$

where ω is a walk on the lattice of M -cubes in T_L^{j-1} . Each $G(\omega)$ has regularity as in (2.30), as well as an exponential decay in the length of the walk. By summing over an appropriate subset of walks that remain inside X , a union of $L^{k-j}r(e_k)$ -cubes, we obtain $G_j(\Omega, X, u)$. [An analogous construction for $C_A^{(k)}(u)$ is described in (2.42)–(2.45).] Walks that stay within $\square(x_1, x_2) \cap \Omega$ define $G_{j, \text{loc}}(\Omega, u; x_1, x_2)$, which is then nonzero only if $|x_1 - x_2| \leq O(L^{k-j}r(e_k))$. A convex combination as in (2.27) is used to preserve regularity across boundaries of M -cubes. The result is the expansion

$$G_j(\Omega, u) = G_{j, \text{loc}}(\Omega, u) + \sum_X G_j(\Omega, X, u). \quad (5.7.8)$$

Of course, $G_j(\Omega, X, u; x_1, x_2) = 0$ unless both x_1 and x_2 are in X . All operators obey the usual regularity bounds, provided $\text{dist}(\{x_1, x_2\}, \Omega^c) > c$. The bound on $G_j(\Omega, X, u)$ has in addition a factor $e^{-cr(e_k)L^{k-j}|X|}$. The dependence on u is in X only; for $G_{j, \text{loc}}$ it is only in an $L^{k-j}r(e_k)$ -neighborhood of x_1, x_2 . Also, when x_1 and x_2 are farther than $L^{k-j}r(e_k)$ from Ω^c , $G_{j, \text{loc}}$ is independent of Ω .

We have developed expansions for $C_A^{(j)}(u)$, $W^{(j)}$, and $G_j(\Omega, u)$. We now put them together to analyze the expansion of the normalization factors. In the expansion (5.7.4) we put $W^{(j)} = W^{(j, \bar{n})} + \sum_X W^{(j)}(X)$. In terms with $l \leq \bar{n}$ we separate from

$\frac{1}{2l} \text{tr}(C_{A_{10}^{(j)}}^{(j)}(u_{k+1})W^{(j, \bar{n})})$ the terms of order $\leq \bar{n}$ in e_j . We can write the sum of all these terms as

$$\sum_{n=1}^{\bar{n}} \frac{1}{n!} \left[\frac{d^n}{de^n} \log Z_{A_{10}^{(j)}}^{(j)}(u_{k+1} \exp(i e' e_j \tilde{A}^c)) \right]_{e'=0}.$$

These will be treated carefully by a resummation. In the other terms we insert the expansion for $C_{A_{10}^{(j)}}^{(j)}(u_{k+1})$; they then take the form $\sum_X W^{(j)'}(X)$, with

$$|W^{(j)'}(X)| \leq e_j^{\bar{n}+1-\alpha} e^{-cr(e_k)|X|} (r(e_k)L^{k-j})^d |X| \leq e_j^\kappa e^{-cr(e_k)|X|}. \quad (5.7.9)$$

We can take κ arbitrarily large by increasing \bar{n} . The high power of e_j beats the big factor $(r(e_k)L^{k-j})^d$, the volume of an elementary cube measured on the j -th scale. This is to account for one free summation on $T_1^{(j)}$; all but one such summation is controlled by exponential decay on the j -th scale.

We return to the perturbative terms. Resummation in j will be possible only if \tilde{A} is localized to sets like $\bar{A}_5^{(j)} \cap \bar{A}_6^{(j+1)c}$. Thus we write

$$\tilde{A} = \theta_k(H_{k, \text{loc}} A^{(k)} + w_5 A') + \sum_{j=-1}^{k-1} \theta_j(1 - \theta_{j+1})w_5 A' = \tilde{A}_k + \sum_{j=0}^{k-1} \tilde{A}_j.$$

Here $\theta_{-1} = 1$, and each \tilde{A}_j is a smooth, small field supported in $\bar{A}_5^{(j)} \cap \bar{A}_6^{(j+1)c}$ ($\bar{A}_5^{(k)}$ if $j = k$). The low order terms can be written as

$$\sum_{n=1}^{\bar{n}} \frac{1}{n!} \left[\prod_{\alpha=1}^n \left(\sum_{j_\alpha=0}^k \frac{d}{de'_{j_\alpha}} \right) \log Z_{A_{10}^{(j)}}^{(j)} \left(u_{k+1} \exp \left(ie'_j \zeta \sum_{l=0}^k e'_l \tilde{A}_l^\zeta \right) \right) \right]_{e'_l=0}.$$

Note that $Z_{A_{10}^{(j)}}^{(j)}(\dots)$ depends only on e_l for $l \geq j$. Thus we can write the last expression as

$$\sum_{m=j}^k \sum_{n=1}^{\bar{n}} \frac{1}{n!} \sum_{\{j_\alpha\}_{\alpha=1}^n: \min_\alpha j_\alpha = m} \times \left[\prod_{\alpha=1}^n \frac{d}{de'_{j_\alpha}} \log Z_{A_{10}^{(j)}}^{(j)} \left(u_{k+1} \exp \left(ie'_j \zeta \sum_{l=j}^k e'_l \tilde{A}_l^\zeta \right) \right) \right]_{e'_l=0}. \tag{5.7.10}$$

We write all terms in the form of the expansions derived in this section, except that we write a new expansion analogous to one we gave above for $(w_5 A')_b^m$:

$$\prod_{\alpha \in \underline{\alpha}} (\theta_{j_\alpha} (1 - \theta_{j_\alpha+1}) w_5 A')_b = \sum_X w_{b, \underline{\alpha}}(X).$$

Here $\alpha \subset \{1, \dots, n\}$, and $w_{b, \underline{\alpha}}(X)$ is also bounded as in (5.7.6).

We insert this expansion in

$$\left[\prod_{\alpha \in \underline{\alpha}} \frac{d}{de'_{j_\alpha}} F_{1, j} \left(\sum_{l=j}^k e'_l \tilde{A}_l^\zeta \right) \right]_{e'_l=0}.$$

The result is a localized expansion $\sum_X F_{1, j, b}^\alpha(X)$, with $|F_{1, j, b}^\alpha(X)| \leq e_j^{1-\alpha} e^{-cr(e_k)|X|}$ (unless $j_\alpha = k$ for all $\alpha \in \underline{\alpha}$, in which case $|X|$ is replaced by $|X|^-$). We make the same expansions in $F_{2, j}$. The expansions for $F_{1, j}$, $F_{2, j}$ are inserted in the low order terms in V_j , $G_{j, \text{loc}}(u_{k+1} \tilde{u})$, and $A_{k, \text{loc}}(u_{k+1} \tilde{u}) + aL^{-2}P(u_{k+1} \tilde{u})$. Finally, they are inserted into (5.7.10), using (5.7.4) for $\log Z$.

The term $m = j$ is special; we bound that term directly without resummation. The random walk expansion is inserted for $C_{A_{10}^{(j)}}^{(j)}(u_{k+1})$, and we obtain an expansion $\sum_X W^{(j)''}(X)$, with

$$|W^{(j)''}(X)| \leq e_j^{1-\alpha} e^{-cr(e_k)|X|} |X' \cap A_5^{(j)'} \cap A_6^{(j+1)c}|.$$

Each term contributing to $W^{(j)''}(X)$ must contain at least one kernel w_5 . There is a summation in $A_{10}^{(j)}$, but since at least one field \tilde{A}_j is present, there is an exponential decay on the j -th scale localizing summations near $A_5^{(j)'} \cap A_6^{(j+1)c}$. This gives rise to the volume factor in the above bound. The volume divergence will be beaten by small factors coming from large fields near $A_5^{(j)'} \cap A_6^{(j+1)c}$; we will have available some $e_j^{\kappa|A_5^{(j)'} \cap A_6^{(j+1)c}|/r(e_j)^d}$, and since $\kappa(\log e_j^{-1})r(e_j)^{-d} > \sum_{k=j+1}^{\log_{\mathbb{Z}}(\epsilon_0/\epsilon)} e_j^{1-\alpha}$, this is sufficient.

Next we take an $m > j$ and we try to replace each $C_{A_{10}^{(j)}}^{(j)}(u_{k+1})$ with $C_{B_{m-j}(A_3^{(m)})}^{(j)}(u_{k+1})$. Using the random walk expansions we can write the difference as $\sum_X C^{(j)'}(X)$, with $|C^{(j)'}(X, x_1, x_2)| \leq e^{-c|x_1 - x_2|} e^{-cr(e_k)|X|}$ for x_1, x_2 in $B_{m-j}(A_4^{(m)})$. For $m > j$, all operators $C_{A_{10}^{(j)}}^{(j)}(u_{k+1})$ in our low order expansion satisfy this restriction. Terms with all $C_{B_{m-j}(A_3^{(m)})}^{(j)}$'s will be considered below. In all other terms, we

random-walk expand any $C_{B_{m-j}(\Lambda_3^{(m)})}^{(j)}(u_{k+1})$'s, sum over m , and obtain

$$\sum_X W^{(j)'''}(X), \quad \text{with} \quad |W^{(j)'''}(X)| \leq e^{-cr(e_j)|X|}.$$

We need to replace the propagators $G_{j,\text{loc}}(u_{k+1})$, $G_j(\square, u_{k+1})$ by $G_j(\bar{\Lambda}_2^{(m)}, u_{k+1})$. They appear in the expansion of $\Delta_{j,\text{loc}}(u_{k+1})$ and indirectly in $C^{(j)}(u_{k+1})$. We write

$$G_{j,\text{loc}}(u_{k+1}) = G_j(\bar{\Lambda}_2^{(m)}, u_{k+1}) + (G_{j,\text{loc}}(u_{k+1}) - G_{j,\text{loc}}(\bar{\Lambda}_2^{(m)}, u_{k+1})) + \sum_X G_j(\bar{\Lambda}_2^{(m)}, X, u_{k+1}),$$

and similarly for $G_{j,\text{loc}}(\square, u_{k+1})$. We only need to look at this operator in $\bar{\Lambda}_3^{(m)}$. There a random walk expansion on the j -th scale will yield the usual regularity bounds on the second term on the right with an extra factor $e^{-cr(e_j)}$. We insert this expansion into $C^{(j)-1}$ to obtain

$$C_{B_{m-j}(\Lambda_3^{(m)})}^{(j)}(u_{k+1})^{-1} = C_{B_{m-j}(\Lambda_3^{(m)})}^{(j)}(\bar{\Lambda}_2^{(m)}, u_{k+1})^{-1} + \sum_X \Delta_j(X),$$

where

$$\begin{aligned} & C_{B_{m-j}(\Lambda_3^{(m)})}^{(j)}(\bar{\Lambda}_2^{(m)}, u_{k+1})^{-1} \\ &= a_j I - a_j^2 Q_j(u_{k+1}) G_j(\bar{\Lambda}_2^{(m)}, u_{k+1}) Q_j^*(u_{k+1}) + aL^{-2} P(u_{k+1}), \quad (5.7.11) \\ & |\Delta_j(X, x_1, x_2)| \leq e^{-cr(e_j)|X|}, \quad = 0 \quad \text{if} \quad x_1 \quad \text{or} \quad x_2 \notin X. \end{aligned}$$

Thus we have

$$\begin{aligned} C_{B_{m-j}(\Lambda_3^{(m)})}^{(j)}(u_{k+1}) &= C_{B_{m-j}(\Lambda_3^{(m)})}^{(j)}(\bar{\Lambda}_2^{(m)}, u_{k+1}) \\ &\quad - C_{B_{m-j}(\Lambda_3^{(m)})}^{(j)}(\bar{\Lambda}_2^{(m)}, u_{k+1}) \left(\sum_X \Delta_j(X) \right) C_{B_{m-j}(\Lambda_3^{(m)})}^{(j)}(u_{k+1}). \end{aligned}$$

This expansion is inserted at each appearance of $C^{(j)}$ in our low order terms. The same analysis is performed when $G_j(\square, u_{k+1})$ appears instead of $G_{j,\text{loc}}(u_{k+1})$. In the leading terms (terms with no $e^{-cr(e_j)}$ from the random walk expansions) we put $\zeta_j'' = 1 + (\zeta_j'' - 1)$. The leading terms are now

$$\begin{aligned} & \sum_{n=1}^{\infty} \frac{1}{n!} \sum_{\{j_\alpha\}; \min_\alpha j_\alpha = m} \\ & \times \left[\prod_{\alpha=1}^n \frac{d}{de'_{j_\alpha}} \log Z_{B_{m-j}(\Lambda_3^{(m)})}^{(j)} \left(\bar{\Lambda}_2^{(m)}, u_{k+1} \exp \left(ie_j \zeta \sum_{l=j}^k e'_l \tilde{A}_l^{\zeta} \right) \right) \right]_{e'_l=0}, \quad (5.7.12) \end{aligned}$$

where this $Z^{(j)}$ uses the quadratic form in (5.7.11). Remainder terms are again localized – there will be typically some delocalized operators and some localized ones. Thus we random walk expand any $G_j(\bar{\Lambda}_2^{(m)}, u_{k+1})$. Also, we expand any $C_{B_{m-j}(\Lambda_3^{(m)})}^{(j)}(\bar{\Lambda}_2^{(m)}, u_{k+1})$ as

$$\sum_{p=0}^{\infty} C_{B_{m-j}(\Lambda_3^{(m)})}^{(j)}(u_{k+1}) \left[\sum_X \Delta_j(X) C_{B_{m-j}(\Lambda_3^{(m)})}^{(j)}(u_{k+1}) \right]^p,$$

and finally we random-walk expand all $C_{B_{m-j}(\Lambda_3^{(m)})}^{(j)}(u_{k+1})$'s. We gather all terms of this rather complicated expansion of the remainders and sum over m , to yield

$$\sum_X W^{(j)(iv)}(X), \quad \text{with} \quad |W^{(j)(iv)}(X)| \leq e^{-cr(e_j)|X|}.$$

As always, X is a connected union of $L^{k-j}r(e_k)$ -cubes, and $W^{(j)(iv)}(X)$ has dependence only on fields in X , \bar{X} , or $B_{k-j}(X)$.

We make a final change in the leading terms, namely we replace $C_{B_{m-j}(A_3^{(m)})}^{(j)}(\bar{A}_2^{(m)}, u_{k+1})$ with $C^{(j)}(\bar{A}_2^{(m)}, u_{k+1})$, the covariance without Dirichlet boundary conditions. We give a random walk expansion for the difference, $\sum_X C^{(j)''}(X)$. It is actually a double expansion, since each term in the usual random walk expansion still depends on $\bar{A}_2^{(m)}$ through the basic quadratic form [which involves $G_j(\bar{A}_2^{(m)}, u_{k+1})$] and through operators $C_{\square}^{(j)}(\bar{A}_2^{(m)}, u_{k+1})$. However, each of these can be expanded as described earlier, yielding terms $C^{(j)''}(X)$ with proper locality properties, and obeying the following bounds:

$$|C^{(j)''}(X, x_1, x_2)| \leq e^{-cr|x_1-x_2|} e^{-cr(e_j)|X|} e^{-c \text{dist}(\{x_1, x_2\}, A_3^{(m)c})}$$

Leading terms are now given as in (5.7.12) but with no Dirichlet boundary conditions. Finally, remainder terms are expanded out completely. All remainder terms have at least one operator $C^{(j)''}(X)$, which provides exponential localization to $B_{m-j}(A_3^{(m)c})$. The field \bar{A}_m is supported in $\bar{A}_5^{(m)}$, thus all terms have at least a factor $e^{-cr(e_j)}$. Thus we can sum all the remainder terms into

$$\sum_X W^{(j)(v)}(X), \quad \text{with } |W^{(j)(v)}(X)| \leq e^{-cr(e_j)|X|}$$

Now the leading terms can be rescaled to the L^{j-m} -lattice, and we sum over $j < m$. All the changes we have made allow us now to compose the normalization factors as

$$\prod_{j=0}^{m-1} Z^{(j), L^{j-m}}(\bar{A}_2^{(m)}, u) = Z_m(\bar{A}_2^{(m)}, u) \cdot \text{const},$$

where the m -step Gaussian normalization factor Z_m arises as in Eq. (2.40) of [7]. We obtain the perturbative expansion

$$\sum_{n=1}^{\bar{n}} \sum_{\{j_\alpha\}: \min_\alpha j_\alpha = m} \times \left[\prod_{\alpha=1}^n \frac{d}{de'_{j_\alpha}} \log Z_m \left(\bar{A}_2^{(m)}, u_{k+1} \exp \left(ie_m L^{-m} \sum_{l=m}^k e'_l \tilde{A}_l^{L^{-m}} \right) \right) \right]_{e'_l=0}$$

The diagrams in this expansion are covered by our theorems on the perturbation expansion. The point is that various Ward identities and symmetries necessary to obtain good bounds can only be seen in this resummed form of perturbation theory. We give the random walk expansion for the propagator $G_m(\bar{A}_2^{(m)}, u_{k+1})$. There is at least one factor e_m in all terms, and a free summation in $A_5^{(m)} \cap A_6^{(m+1)c} (m < k)$ or $A_5^{(m)} (m = k)$. Thus we can write the perturbation expansion for $m < k$ as $\sum_X W^{(m)(vi)}(X)$, with

$$|W^{(m)(vi)}(X)| \leq e_m^{1-\alpha} e^{-cr(e_k)|X|} |X \cap A_5^{(m)'} \cap A_6^{(m+1)c}|$$

As for the $W^{(j)''}$ terms, the volume factor will be beaten by convergence factors from the large field region $A_5^{(m)'} \cap A_6^{(m+1)c}$.

The term $m=k$ is treated slightly differently. We decompose \tilde{A}_k^η into $\theta_k H_{k,\text{loc}} A^{(k)} + \theta_k w_5 A'$. Terms with one or more $w_5 A'$ field are expanded as in the $m < k$ case. In terms with all $\theta_k H_{k,\text{loc}} A^{(k)}$ fields, we expand the propagators as before, leaving in the main terms the localized propagator $G_{k,\text{loc}}(u_{k+1})$. The result is our standard perturbative expansion in the field $\theta_k H_{k,\text{loc}} A^{(k)}$, which we denote $Q^{(k)}(u_{k+1}, \theta_k H_{k,\text{loc}} A^{(k)})$. The remainder terms become $\sum_X W^{(k)(vi)}(X)$, with $|W^{(k)(vi)}(X)| \leq e^{-cr(e_k)|X|}$.

We can summarize the results of this section as follows:

$$\begin{aligned} & \prod_{j=0}^{k-1} Z_{A_{10}^{(j)}}^{(j)}(u_{k+1}, \tilde{u}) \\ &= \prod_{j=0}^{k-1} Z_{A_{10}^{(j)}}^{(j)}(u_{k+1}) \exp \left[-Q^{(k)}(u_{k+1}, \theta_k H_{k,\text{loc}} A^{(k)} - \sum_X W_2^{(k)}(X)) \right], \end{aligned} \tag{5.7.13}$$

where X is a connected union of $r(e_k)$ -cubes in $T_1^{(k)}$,

$$W_2^{(k)}(X) = \sum_{j=0}^k W^{(j)'}(X) + \dots + W^{(j)(vi)}(X),$$

$$|W_2^{(k)}(X)| \leq e_k^\kappa e^{-cr(e_k)|X|} + \sum_{j < k} e_j^{1-\alpha} e^{-cr(e_k)|X|} |B_{k-j-1}(X) \cap A_5^{(j)'} \cap A_6^{(j+1)c}|, \tag{5.7.14}$$

and

$$\begin{aligned} & Q^{(k)}(u_{k+1}, \theta_k H_{k,\text{loc}} A^{(k)}) \\ &= \sum_{n=1}^{\bar{n}} \frac{d^n}{d e'^n} \log \left[\int d\phi |_{\bar{A}_4^{(k)}} \exp \left(-\frac{1}{2} \langle \phi, G_{k,\text{loc}}(u_{k+1} e^{ie'e_k \eta \theta_k H_{k,\text{loc}} A^{(k)}})^{-1} \phi \rangle \right) \right]_{e'=0}. \end{aligned} \tag{5.7.15}$$

5.8. Scalar Field Translation

The scalar field quadratic forms, after all our manipulations with the gauge field, are as follows:

$$\frac{1}{2} \langle A_8^{(k-1)'} \phi, A_{k,\text{loc}}(\tilde{u}_{k+1}) A_8^{(k-1)'} \phi \rangle + \frac{1}{2} a L^{-2} \langle \psi - Q(\tilde{u}_{k+1}) \phi, \psi - Q(\tilde{u}_{k+1}) \phi \rangle.$$

To eliminate most of the linear term $\langle \psi, Q(\tilde{u}_{k+1}) \phi \rangle$ in the small field region, we make a translation

$$\phi = \phi^{(k)} + a L^{-2} A_7^{(k)} C_{\text{loc}}^{(k)}(u_{k+1}) Q^*(u_{k+1}) \psi. \tag{5.8.1}$$

(Recall that $u_{k+1} = \tilde{u}_{k+1}$ in $\bar{A}_6^{(k)}$.)

The terms quadratic in $\phi^{(k)}$ are then

$$\frac{1}{2} \langle A_8^{(k-1)'} \phi^{(k)}, (A_{k,\text{loc}}(\tilde{u}_{k+1}) + a L^{-2} P(\tilde{u}_{k+1})) A_8^{(k-1)'} \phi^{(k)} \rangle + \mathcal{Q}_4, \tag{5.8.2}$$

where

$$\mathcal{Q}_4 = \frac{1}{2} a L^{-2} \langle \phi^{(k)}, A_8^{(k-1)'} P(\tilde{u}_{k+1}) \phi^{(k)} \rangle.$$

In the cross terms between $\phi^{(k)}$ and ψ , we write $\psi = A_8^{(k)'} \psi + A_8^{(k)c} \psi$. The terms with $A_8^{(k)c} \psi$ define \mathcal{Q}_5 , a form localized near $A_8^{(k)c}$. The other terms can be written as $\langle \phi^{(k)}, w_6 \psi \rangle$, with w_6 a small kernel with range less than $r(e_k)$. This is because (5.8.1) would eliminate entirely the linear term were it not for the localizations.

In the terms quadratic in ψ , we again combine all terms involving $A_8^{(k)c}\psi$ into a form \mathcal{Q}_6 localized near $A_8^{(k)c}$. The remaining terms become

$$\frac{1}{2}aL^{-2}\langle A_8^{(k)'}\psi, (I - aL^{-2}Q(u_{k+1})C_{\text{loc}}^{(k)}(u_{k+1})Q(u_{k+1})^*)A_8^{(k)'}\psi \rangle + \frac{1}{2}\langle \psi, w'_7\psi \rangle,$$

with w'_7 another small, local kernel. We apply the identity [7]

$$\begin{aligned} & aL^{-2}I - a^2L^{-4}Q(u_{k+1})C^{(k)}(u_{k+1})Q(u_{k+1})^* \\ & = a_kL^{-2}I - a_k^2L^{-4}Q_{k+1}(u_{k+1})G_{k+1}^\eta(u_{k+1})Q_{k+1}(u_{k+1})^*, \end{aligned}$$

but in a localized version with $C_{\text{loc}}^{(k)}$ and $G_{k+1, \text{loc}}^\eta$ and with another small kernel w''_7 on the right. This yields the desired form $\Delta_{k+1, \text{loc}}^L(u_{k+1})$, and so we obtain

$$\frac{1}{2}\langle A_8^{(k)'}\psi, \Delta_{k+1, \text{loc}}^L(u_{k+1})A_8^{(k)'}\psi \rangle + \frac{1}{2}\langle \psi, w_7\psi \rangle, \quad \text{with } w_7 = w'_7 + w''_7.$$

To summarize, we have written

$$\begin{aligned} & \frac{1}{2}\langle A_8^{(k-1)'}\phi, \Delta_{k, \text{loc}}(\tilde{u}_{k+1})A_8^{(k-1)'}\phi \rangle + \frac{1}{2}aL^{-2}\langle \psi - Q(\tilde{u}_{k+1})\phi, \psi - Q(\tilde{u}_{k+1})\phi \rangle \\ & = \mathcal{Q}_4 + \mathcal{Q}_5 + \mathcal{Q}_6 + \frac{1}{2}\langle A_8^{(k-1)'}\phi^{(k)}, (\Delta_{k, \text{loc}}(\tilde{u}_{k+1}) + aL^{-2}P(\tilde{u}_{k+1}))A_8^{(k-1)'}\phi^{(k)} \rangle \\ & \quad + \frac{1}{2}\langle A_8^{(k)''}\psi, \Delta_{k+1, \text{loc}}^L(u_{k+1})A_8^{(k)''}\psi \rangle + \langle \phi^{(k)}, w_6\psi \rangle + \frac{1}{2}\langle \psi, w_7\psi \rangle, \end{aligned} \tag{5.8.3}$$

with w_6, w_7 small local kernels, and with $\mathcal{Q}_4, \mathcal{Q}_5, \mathcal{Q}_6$ localized near $A_8^{(k)c}$.

5.9. Bounds on Fluctuation and Block Fields

As we remarked earlier, the restrictions on $u(p)$ and the gauge field renormalization transformations imply that

$$\left| \frac{1}{ie_k} \log v(p) \right| = |f(p)| \leq cp(e_k), \quad p \in A_0^{(k)''**}. \tag{5.9.1}$$

Also, bounds on ϕ and $\psi - Q(u_k)\phi$ imply that for $y \in A_0^{(k)'}$,

$$\begin{aligned} & |\psi(y)| \leq cp(e_k)\lambda_k^{-1/4}, \quad (L^k\varepsilon)^d < \lambda, \\ & \left| |\psi(y)| - (8\lambda)^{-1/2}(L^k\varepsilon)^{(d-2)/2} \right| \leq cp(e_k)(L^k\varepsilon)^{-1}, \quad (L^k\varepsilon)^d \geq \lambda. \end{aligned} \tag{5.9.2}$$

Next, we wish to prove that

$$|u_{k+1}(\langle b_-, b_+ \rangle)\psi(b_+) - \psi(b_-)| \equiv |(D_{\bar{u}_{k+1}}\psi)(b)| \leq cp(e_k), \quad b \in A_0^{(k)''*}. \tag{5.9.3}$$

We prove the bound first for $D_{\bar{u}_k}\psi$ (before the gauge transformation of Sect. 5). Our bounds on $\psi - Q(u_k)\phi$ reduce this to estimating

$$|u_k(\Gamma_{b_-, x})\phi(x) - u_k(\langle b_-, b_+ \rangle)u_k(\Gamma_{b_+, x'})\phi(x')|$$

for any $x \in B(b_-), x' \in B(b_+)$. This is proven with several applications of our bounds on $D_{\bar{u}_k}\phi$. In going from u_k to u_{k+1} we made a gauge transformation and removed some small fields. Also, the gauge transformation was not quite compensated by a rotation of ψ . Thus in going from the old $|D_{\bar{u}_k}\psi|$ to the new $|D_{\bar{u}_{k+1}}\psi|$ we make errors of the order of $ce_k p(e_k)^3\lambda_k^{-1/4}, (L^k\varepsilon)^d < \lambda$ or

$$ce_k p(e_k)^3(\lambda^{-1/2}(L^k\varepsilon)^{(d-2)/2} + (L^k\varepsilon)^{-1}), \quad (L^k\varepsilon)^d \geq \lambda.$$

In both cases this is bounded by $e^\beta(L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha}$ – see the discussion below of the bounds on interaction terms. The desired bound follows.

These bounds allow us to insert the following characteristic functions:

$$\begin{aligned} \chi_{k+1, A_0^{(k)'}} &= \prod_{p \in A_0^{(k)'**}} \chi(cp(e_k)p(e_k), |v(p) - 1|) \\ &\quad \times \prod_{y \in A_0^{(k)'}} \chi(cp(e_k)\lambda_k^{-1/4}, |\psi(y)|) \prod_{b \in A_0^{(k)'*}} \chi(cp(e_k), |(D_{\tilde{u}_{k+1}}\psi)(b)|) \end{aligned}$$

and the integral is unchanged. If $(L^k \varepsilon)^d \geq \lambda$, the bound on ψ is replaced with $\chi(cp(e_k)(L^k \varepsilon)^{-1}, (|\psi(y)| - (8\lambda)^{-1/2}(L^k \varepsilon)^{(d-2)/2}))$.

We remarked earlier that A' is small in $A_1^{(k)*}$. We then defined $A^{(k)} = A' + A_4^{(k)*} L^{-2} C_{\text{loc}}^{(k)} H_{k, \text{loc}}^* \partial^* Q_{k+1}^* f$. Since f is small and $H_{k, \text{loc}}$ is regular, we have that

$$|A_b^{(k)}| \leq cp(e_k), \quad b \in A_1^{(k)*}. \quad (5.9.4)$$

We want a similar bound for $\phi^{(k)}(x)$, $x \in A_7^{(k)}$. Note that $C_{\text{loc}}^{(k)}(u_{k+1})$ is almost equal to $C^{(k)}(u_{k+1})$. Thus we have that in $A_7^{(k)}$, say

$$aL^{-2} C_{\text{loc}}^{(k)}(u_{k+1}) Q(u_{k+1})^* \psi = Q(u_{k+1})^* \psi + O(p(e_k)),$$

(the corresponding statement with $C^{(k)}(u_{k+1})$ was proven in [8, Eq. (2.113)]. Using arguments like the ones we used to bound $D_{\tilde{u}_{k+1}}\psi$, we can replace $Q(u_{k+1})^* \psi$ with ϕ in this bound. This proves that

$$|\phi^{(k)}(x)| \leq cp(e_k), \quad x \in A_7^{(k)*}. \quad (5.9.5)$$

The bounds (5.9.4), (5.9.5) allow us to insert the characteristic functions

$$\chi'_{A_7^{(k)}} = \prod_{b \in A_7^{(k)*}} \chi(cp(e_k), A^{(k)}) \prod_{x \in A_7^{(k)}} \chi(cp(e_k), \phi^{(k)})$$

without changing anything.

We note that the restrictions implied by $\chi_{A_0^{(k)}}$ are stronger than the corresponding restrictions in $\chi_{k, A_0^{(k-1)'}}$ in $A_0^{(k)}$. [When $(L^k \varepsilon)^d \geq \lambda$, we use the inequality $(8\lambda)^{-1/2}(L^k \varepsilon)^{(d-2)/2} + p(e_k)(L^k \varepsilon)^{-1} \leq cp(e_k)\lambda_k^{-1/4}$.] Thus we can replace $\chi_{k, A_0^{(k-1)'}}$ with $\chi_{k, A_0^{(k-1)' \cap A_1^{(k)c}}$ without changing anything.

Let us summarize the operations performed so far by using the concluding formulae in the last several sections to write a complete expression for our density.

$$\begin{aligned} &Q_{k+1}^L(v, \psi) \\ &= \sum_{\{X_\omega\}} \sum_{A_0^{(k)}} \int du^{(k)} d\phi^{(k)} \delta_{A_X}(u^{(k)}) \delta_{A_1^{(k)'**c}}(v/Q u^{(k)}) \delta_{A_1^{(k)'**}} \left(\frac{e_k}{2\pi} Q A^{(k)} \right) \int \prod_{j=0}^{k-1} du^{(j)} |_{A_{10}^{(j)c**}} \\ &\quad \times \zeta_{A_0^{(k)c}} \chi_{A_0^{(k)}} \chi_{k, A_0^{(k-1)' \cap A_1^{(k)c}} \chi_{k+1, A_0^{(k)}} \chi'_{A_7^{(k)}} \prod_{\omega} g_k(X_\omega) \prod_{\sigma} (F_{k, \text{loc}}^{(\bar{m})}(X_\sigma) + \tilde{F}_{\text{loc}}^{(k)}(X_\sigma)) \\ &\quad \times \prod_{j=0}^{k-1} [Z_{A_{10}^{(j)c**}} Z_{A_{10}^{(j)}}(u_{k+1})] \exp \left[-\frac{1}{2} \langle A_1^{(k)**} \partial A^{(k)}, \sigma_{k, \text{loc}} A_1^{(k)**} \partial A^{(k)} \rangle \right. \\ &\quad - \frac{1}{2} \langle A_5^{(k)**} f, \sigma_{k+1, \text{loc}}^L A_5^{(k)'**} f \rangle - \mathcal{Q}_1 - \mathcal{Q}_2 - \mathcal{Q}_3 - \langle f, w_3 A^{(k)} \rangle - \frac{1}{2} \langle f, w_4 f \rangle \\ &\quad - \mathcal{Q}_4 - \mathcal{Q}_5 - \mathcal{Q}_6 - \frac{1}{2} \langle A_8^{(k-1)'} \phi^{(k)}, (A_{k, \text{loc}}(\tilde{u}_{k+1}) + aL^{-2} P(\tilde{u}_{k+1})) A_8^{(k-1)'} \phi^{(k)} \rangle \\ &\quad - \frac{1}{2} \langle A_8^{(k)'} \psi, \Delta_{k+1, \text{loc}}^L(u_{k+1}) A_8^{(k)'} \psi \rangle - \langle \phi^{(k)}, w_6 \psi \rangle \\ &\quad - \frac{1}{2} \langle \psi, w_7 \psi \rangle - \mathcal{E}_k - E^{(k)} - \mathcal{P}_{k, \text{loc}}(A_8^{(k-1)'}, \tilde{u}_{k+1}) - R^{(k)}(u_{k+1}, \theta_k H_{k, \text{loc}} A^{(k)}) \\ &\quad \left. - \sum_{\square} W_1^{(k)}(\square) - Q^{(k)}(u_{k+1}, \theta_k H_{k, \text{loc}} A^{(k)}) - \sum_X W_2^{(k)}(X) \right]. \quad (5.9.6) \end{aligned}$$

5.10. The Interaction for the Fluctuation Fields

Having made the scalar field translation, we regard the terms $\mathcal{P}_{k,\text{loc}}, R^{(k)}, Q^{(k)}, F_{k,\text{loc}}^{(\bar{m})}$ as polynomials in $\phi^{(k)}, A^{(k)}$. We make some small changes and localizations in order to obtain the standard form of the fluctuation field interaction in $A_8^{(k)}$.

We have the external field

$$a_k a L^{-2} G_{k,\text{loc}}(\tilde{u}_{k+1}) Q_k^*(\tilde{u}_{k+1}) A_7^{(k)} C_{\text{loc}}^{(k)}(u_{k+1}) Q^*(u_{k+1}) \psi$$

appearing in the diagrams in $\mathcal{P}_{k,\text{loc}}, R^{(k)}, F_{k,\text{loc}}^{(\bar{m})}$. The first we leave alone, whereas in the second we localize the field to $\bar{A}_8^{(k)}$ and replace it with

$$a_{k+1} L^{-2} G_{k+1,\text{loc}}^{\eta}(u_{k+1}) Q_{k+1}^*(u_{k+1}) \psi + w_8 \psi.$$

The kernel w_8 is local and small with small derivatives and Hölder derivatives. This is accomplished in the usual fashion by replacing $G_{k,\text{loc}}(u_{k+1}), C_{\text{loc}}^{(k)}(u_{k+1})$ with the corresponding operators with Neumann boundary conditions on an $r(e_k)$ -cube \square . The propagator composition formula [7, Eq.(2.41)] is applied, and $G_{k+1}^{\eta}(\square, u_{k+1})$ is localized again.

We localize all vertices to $\bar{A}_8^{(k)}$; vector field legs at a vertex are multiplied by a smooth function $\bar{\theta}_k$ changing from 0 to 1 in a neighborhood of $\bar{A}_8^{(k)c}$. We also remove all diagrams whose combined order in $\lambda^{1/2}$ and e is greater than \bar{n} . We still consider all P_k vertices together; any $P_k^{(l)}$ vertex is considered as one power of λ . Each mass renormalization counterterm is written graphically and powers counted accordingly. The result is the interaction $V^{(k)}(A_8^{(k)}, u_{k+1}, A^{(k)}, \phi^{(k)})$, and

$$\begin{aligned} & \mathcal{P}_{k,\text{loc}}(A_8^{(k-1)}, \tilde{u}_{k+1}) + R^{(k)}(u_{k+1}, \theta_k H_{k,\text{loc}} A^{(k)}) + Q^{(k)}(u_{k+1}, \theta_k H_{k,\text{loc}} A^{(k)}) \\ &= \mathcal{P}_{k,\text{loc}}(A_8^{(k-1)}, A_8^{(k)c}, \tilde{u}_{k+1}) + R^{(k)}(A_8^{(k)c}, u_{k+1}, \theta_k H_{k,\text{loc}} A^{(k)}) \\ &+ Q^{(k)}(A_8^{(k)c}, u_{k+1}, \theta_k H_{k,\text{loc}} A^{(k)}) + V^{(k)}(A_8^{(k)}, u_{k+1}, A^{(k)}, \phi^{(k)}) + \sum_{\square} W_3^{(k)}(\square). \end{aligned}$$

Here in writing $A_8^{(k)c}$ we mean that only the terms without proper localizations are included. The terms $W_3^{(k)}(\square)$ contain terms localized near the $r(e_k)$ -cube \square which involve the small kernel w_8 or have high powers of coupling constants. We have an estimate

$$|W_3^{(k)}(\square)| \leq [e^{\beta(L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha}}]^{\bar{n} + 1} \leq e^{\bar{n} \beta (L^k \varepsilon / \varepsilon_0)^{\kappa}},$$

with $\kappa > d$ as large as desired if $\bar{n} > \bar{n}(\kappa)$. This estimate comes from our analysis of the perturbation expansion and the restrictions on the fields. We find that each vertex results in at least a factor $e^{\beta(L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha}}$.

Estimates on $V^{(k)}, Q^{(k)}, R^{(k)}$ follow from the same analysis. When localized for example to a cube of size $r(e_k)$, all terms [except for $P_k(a_{k+1} L^{-2} G_{k+1,\text{loc}}^{\eta}(u_{k+1}) Q_{k+1}^*(u_{k+1}) \psi)$] are bounded by $e^{\beta(L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha}}$, with α, β small and positive.

In a similar fashion we modify the external scalar fields in $F_{k,\text{loc}}^{(\bar{m})}$ and eliminate diagrams of order higher than \bar{m} . Thus we write

$$F_{k,\text{loc}}^{(\bar{m})}(X_{\sigma}) = F_{k,\text{loc}}^{\bar{m}}(X_{\sigma}) + \tilde{F}_{k,\text{loc}}^{\bar{m}}(X_{\sigma}),$$

with $\tilde{F}_{k,\text{loc}}^{\bar{m}}(X_{\sigma})$ containing the w_8 terms and the higher order terms, and satisfying $\tilde{F}_{k,\text{loc}}^{\bar{m}}(X_{\sigma}) \leq c(F)$. We regard $F_{k,\text{loc}}^{\bar{m}}$ as a polynomial in $A^{(k)}, \phi^{(k)}$.

5.11. Mayer Expansion I

In this section we expand irrelevant terms down from the exponent. This operation is done to simplify the structure of the integral in the region free of irrelevant terms.

Let us combine the irrelevant terms as follows:

$$\begin{aligned} &\langle f, w_3 A^{(k)} \rangle + \frac{1}{2} \langle f, w_4 f \rangle + \langle \phi^{(k)}, w_6 \psi \rangle + \frac{1}{2} \langle \psi, w_7 \psi \rangle + \sum_{\square} W_1^{(k)}(\square) + \sum_X W_2^{(k)}(X) \\ &+ \sum_{\square} W_3^{(k)}(\square) = \sum_X W_4^{(k)}(X), \end{aligned}$$

with $W_4^{(k)}(X)$ containing terms with dependence in X . We combine the estimates on the above terms to obtain

$$\begin{aligned} |W_4^{(k)}(X)| &\leq [e^{\beta(L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha}}]^{\bar{n} + 1} e^{-cr(e_k)|X|} \\ &+ \sum_{j < k} e_j^{1 - \alpha} e^{-cr(e_k)|X|} |B_{k-j-1}(X) \cap A_5^{(j)'} \cap A_6^{(j+1)c}|. \end{aligned}$$

We can only Mayer-expand small terms, therefore we parcel up $W_4^{(k)}(X)$ into manageable chunks. It is a simple matter to decompose $W_4^{(k)}(X)$ as follows

$$W_4^{(k)}(X) = \sum_{j < k} \sum_{x_j \in B_{k-j-1}(X) \cap A_5^{(j)'} \cap A_6^{(j+1)c}} W_{4,j}^{(k)}(x_j, X) + W_{4,k}^{(k)}(x_k, X).$$

Here x_k is some distinguished point in X (for unity of notation) and

$$\begin{aligned} |W_{4,j}^{(k)}(x_j, X)| &\leq e_j^{1 - \alpha} e^{-cr(e_k)|X|}, \\ |W_{4,k}^{(k)}(x_k, X)| &\leq [e^{\beta(L^k \varepsilon / \varepsilon_0)}]^{\bar{n} + 1} e^{-cr(e_k)|X|}. \end{aligned} \tag{5.11.1}$$

The Mayer expansion is the usual identity

$$\exp\left(-\sum_X W_4^{(k)}(X)\right) = \sum_{S_4} \prod_{(j, x_j, X) \in S_4} (e^{-W_{4,j}^{(k)}(x_j, X)} - 1). \tag{5.11.2}$$

Let \tilde{S}_4 be the set of all triplets (j, x_j, X) that arise in the above decomposition of $W_4^{(k)}(X)$, for any X . Then S_4 is summed over subsets of \tilde{S}_4 . Note that $e^{-W_{4,j}^{(k)}(x_j, X)} - 1$ satisfies the same bound as $W_{4,j}^{(k)}(x_j, X)$.

To see what kind of control we have over this expansion, let us do a typical estimate of the type we need:

$$\begin{aligned} &\left| \sum_{S_4 = \{(j_\alpha, x_{j_\alpha}, X_\alpha)\} : \bigcup_\alpha X_\alpha = X} \prod (e^{-W_{4,j}^{(k)}(x_j, \alpha, X_\alpha)} - 1) \right| \\ &\leq \exp\left(\sum_{j < k} e_j^{1 - \alpha} e^{-cr(e_k)|B_{k-j-1}(X) \cap A_5^{(j)'} \cap A_6^{(j+1)c}|} [e^{\beta(L^k \varepsilon / \varepsilon_0)}]^{(\bar{n} + 1)|X|}\right). \end{aligned} \tag{5.11.3}$$

We consider first sums over X_α such that $x_{j_\alpha} = x_j$. A combinatoric factor $c^{|X_\alpha|}$ controls each sum over X_α , and can be absorbed into the factors $e^{-cr(e_k)|X|}$ in our bounds on $e^{-W^{(k)}(x_j, X)} - 1$. If there are n such sets, we use n factors of $e_j^{1 - \alpha} E^{-cr(e_k)}$, $j < k$. The resulting estimate has a factor

$$\sum_{n=0}^{\infty} (e_j^{1 - \alpha} e^{-cr(e_k)})^n \leq \exp(e_j^{1 - \alpha} e^{-cr(e_k)})$$

at each x_j , $j < k$, or $O(1)$, $j = k$. There remains a factor $[e^{\beta(L^k \varepsilon / \varepsilon_0)}]^{(\bar{n} + 1)|X|}$ from a worst-case analysis of the unused small factors, and (5.11.3) follows.

We also expand out the observable:

$$\prod_{\sigma} (F_{k,\text{loc}}^{\tilde{m}}(X_{\sigma}) + \tilde{F}_{k,\text{loc}}(X_{\sigma}) + \tilde{\tilde{F}}_{k,\text{loc}}(X_{\sigma})) = \sum_{\tilde{\sigma}_1} \prod_{\sigma_1 \in \tilde{\sigma}_1} F_{k,\text{loc}}^{\tilde{m}}(X_{\sigma_1}) \prod_{\sigma \notin \tilde{\sigma}_1} F'_{k,\text{loc}}(X_{\sigma}),$$

with $F'_{k,\text{loc}} = \tilde{F}_{k,\text{loc}} + \tilde{\tilde{F}}_{k,\text{loc}}$, and with $\tilde{\sigma}_1$ summed over subsets of the index set for σ on the left-hand side.

We now fix $\tilde{\sigma}_1$, S_4 , and define $\tilde{A}_8^{(k)}$ as follows:

$$\tilde{A}_8^{(k)} = A_8^{(k)} \setminus \bigcup_{\sigma \notin \tilde{\sigma}_1} X_{\sigma} \setminus \bigcup_{(j,x_j,X) \in S_4} X.$$

Then we define $A_9^{(k)}$ by deleting a collar neighborhood of width $r(e_k)$ from $\tilde{A}_8^{(k)}$. In $A_9^{(k)}$ we attempt to remove the characteristic functions $\chi_{A_9^{(k)}}$. Thus we write

$$\chi_{A_9^{(k)}} = \chi_{A_9^{(k)} \cap A_9^{(k)c} \chi_{A_9^{(k)}},$$

and for each type of characteristic function in $\chi_{A_9^{(k)}}$ we expand $\chi = 1 - \chi^c$, as follows:

$$\prod_{x \in A_9^{(k)}} \chi_x = \sum_{S_x \subset A_9^{(k)}} \prod_{x \in S_x} (-\chi_x^c).$$

We have similar sums over $S_y \subset A_9^{(k) \prime}$, $S_b \subset A_9^{(k) *}$, $S_p \subset A_9^{(k) **}$, and we define $\tilde{A}_9^{(k)}$ as the union of all $r(e_k)$ -cubes in $A_9^{(k)}$, none of whose points are in S_x , or in bonds, plaquettes, or blocks in S_b , S_p , S_y . The characteristic function expansion can now be written as

$$\begin{aligned} \chi_{A_9^{(k)}} &= \sum_{\tilde{A}_9^{(k)}} \zeta'_{\tilde{A}_9^{(k)c}}, \\ \zeta'_{\tilde{A}_9^{(k)c}} &= \sum_{\{S_x, S_y, S_b, S_p\} \text{ compatible with } \tilde{A}_9^{(k)c}, A_9^{(k)}} \prod_{x \in S_x} (-\chi_x^c) \\ &\quad \times \prod_{y \in S_y} (-\chi_y^c) \prod_{b \in S_b} (-\chi_b^c) \prod_{p \in S_p} (-\chi_p^c). \end{aligned}$$

Finally we define $A_{10}^{(k)}$ by deleting a collar neighborhood from $\tilde{A}_9^{(k)}$.

These expansions complicate our expression for $\varrho_{k+1}^L(v, \psi)$ in (5.9.6), however the integral in $A_{10}^{(k)}$ is quite simple now. It involves a small, local, polynomial interaction $V^{(k)}$ modifying a Gaussian integral in $\phi^{(k)}$, $A^{(k)}$. The inverse covariance is local and bounded from above and from below. The characteristic functions χ' are simple functions of $\phi^{(k)}$, $A^{(k)}$ keeping them bounded. The observable is a product of polynomial pieces given by low-order perturbation theory. Large field and nonperturbative effects have been separated out.

5.12. Conditional Integration

We exploit the simple structure in $A_{10}^{(k)}$ by doing the integrals there with conditioning on $A_{10}^{(k)c}$, $A_{10}^{(k)c*}$. The formula we use is a generalization of the following identity for scalar fields:

$$\begin{aligned} &\int d\phi|_{A^c} F(\phi|_{A^c}) \int d\phi|_A e^{-\langle A^c \phi, A A \phi \rangle} e^{-1/2 \langle \phi, A A \phi \rangle} G(\phi) \\ &= \int d\phi|_{A^c} F(\phi|_{A^c}) \int d\phi|_A e^{-\langle A^c \phi, A A \phi \rangle} e^{-1/2 \langle \phi, A A \phi \rangle} \\ &\quad \times \frac{\int d\phi|_A G(\phi) e^{-1/2 \langle \phi, A A \phi \rangle} e^{-\langle A^c \phi, A A \phi \rangle}}{\int d\phi|_A e^{-1/2 \langle \phi, A A \phi \rangle} e^{-\langle A^c \phi, A A \phi \rangle}} \\ &= (\int d\phi|_A e^{-1/2 \langle \phi, A A \phi \rangle}) \int d\phi|_{A^c} F(\phi|_{A^c}) e^{1/2 \langle A^c \phi, A A A^{-1} A A^c \phi \rangle} \\ &\quad \times \frac{1}{\mathcal{N}} \int d\phi|_A G(\phi) e^{-1/2 \langle \phi, A A \phi \rangle} e^{-\langle A^c \phi, A A \phi \rangle}. \end{aligned}$$

Here \mathcal{N} is equal to the last integral, without $G(\phi)$. Thus in our expression for $Q_{k+1}^I(v, \psi)$, we have now an “exterior” integral over $u^{(k)}$, $\phi^{(k)}$ in $A_{10}^{(k)c}$, whose Gaussian piece has been replaced by

$$\begin{aligned} & \delta_{\text{Ax}, A_{10}^{(k)c}}(u^{(k)}) \delta_{A_1^{(k)**} \cap A_{10}^{(k)c*}} \left(\frac{e_k}{2\pi} Q A^{(k)} \right) \\ & \times \exp \left[-\frac{1}{2} \langle A_1^{(k)**} \partial A_{10}^{(k)c*} A^{(k)}, \sigma_{k, \text{loc}} A_1^{(k)**} \partial A_{10}^{(k)c} A^{(k)} \rangle \right. \\ & - \frac{1}{2} \langle (A_8^{(k-1)'} \cap A_{10}^{(k)c}) \phi^{(k)}, (\Delta_{k, \text{loc}}(\tilde{u}_{k+1}) + aL^{-2} P(\tilde{u}_{k+1})) (A_8^{(k-1)'} \cap A_{10}^{(k)c}) \phi^{(k)} \rangle \\ & + \frac{1}{2} \langle A_{10}^{(k)c} \phi^{(k)}, \Delta_{k, \text{loc}}(u_{k+1}) C_{A_{10}^{(k)}}^{(k)}(u_{k+1}) A_{k, \text{loc}}(u_{k+1}) A_{10}^{(k)c} \phi^{(k)} \rangle \\ & - \frac{1}{2} \langle A_{10}^{(k)c*} A^{(k)}, L^{-1} Q^* Q^s A_{10}^{(k)c**} \partial^* \sigma_{k, \text{loc}} \partial (L^{-1} A_{10}^{(k)c**} Q^{s*} Q - 2) A_{10}^{(k)c*} A^{(k)} \rangle \\ & + \frac{1}{2} \langle A_{10}^{(k)c*} A^{(k)}, (I - L^{-1} Q^* Q^s A_{10}^{(k)c**}) \partial^* \sigma_{k, \text{loc}} \partial C_{A_{10}^{(k)c**}}^{(k)} \partial^* \sigma_{k, \text{loc}} \partial \\ & \times (I - L^{-1} A_{10}^{(k)c**} Q^{s*} Q) A_{10}^{(k)c*} A^{(k)} \rangle \left. \right] Z_{A_{10}^{(k)c**} c}^{(k)} Z_{A_{10}^{(k)}}^{(k)}(u_{k+1}). \end{aligned} \quad (5.12.1)$$

The “interior” integral is

$$\begin{aligned} & \frac{1}{\mathcal{N}} \int d\phi^{(k)} |_{A_{10}^{(k)}} dA^{(k)} |_{A_{10}^{(k)c**} c} \delta_{\text{Ax}, A_{10}^{(k)}}(A^{(k)}) \delta_{A_{10}^{(k)c**} c} (Q A^{(k)}) \chi'_{A_{10}^{(k)}} \prod_{\sigma_1 \in \hat{\sigma}_1} F_{k, \text{loc}}^{\bar{m}}(X_{\sigma_1}) \\ & \times \exp \left[-\frac{1}{2} \langle A_{10}^{(k)} \phi^{(k)}, (\Delta_{k, \text{loc}}(u_{k+1}) + aL^{-2} P(u_{k+1})) (A_{10}^{(k)} + 2A_{10}^{(k)c}) \phi^{(k)} \rangle \right. \\ & - \frac{1}{2} \langle A_{10}^{(k)c**} A^{(k)}, \partial^* \sigma_{k, \text{loc}} \partial (A_{10}^{(k)c**} + 2A_{10}^{(k)c*}) A^{(k)} \rangle \\ & \left. - V^{(k)}(A_8^{(k)}, u_{k+1}, A^{(k)}, \phi^{(k)}) \right]. \end{aligned} \quad (5.12.2)$$

Here \mathcal{N} is defined by the last integral, but without $\chi'_{A_{10}^{(k)}}$, $F_{k, \text{loc}}^{\bar{m}}$, or $V^{(k)}$.

Let us describe more carefully the calculations leading to (5.12.1). The third form, together with $Z_{A_{10}^{(k)}}^{(k)}(u_{k+1})$, is a calculation of

$$\int d\phi^{(k)} |_{A_{10}^{(k)}} \exp \left[-\frac{1}{2} \langle A_{10}^{(k)} \phi^{(k)}, (\Delta_{k, \text{loc}}(u_{k+1}) + aL^{-2} P(u_{k+1})) (A_{10}^{(k)} + 2A_{10}^{(k)c}) \phi^{(k)} \rangle \right].$$

The 4-th and 5-th forms, with $Z_{A_{10}^{(k)c**} c}^{(k)}$, are a calculation of

$$\begin{aligned} & \int dA^{(k)} |_{A_{10}^{(k)c**} c} \delta_{\text{Ax}, A_{10}^{(k)}}(A^{(k)}) \delta_{A_{10}^{(k)c**} c} (Q A^{(k)}) (e_k/2\pi)^{\|A_{10}^{(k)c**} c\|} \\ & \times \exp \left[-\frac{1}{2} \langle A_{10}^{(k)c**} A^{(k)}, \partial^* \sigma_{k, \text{loc}} \partial (A_{10}^{(k)c**} + 2A_{10}^{(k)c*}) A^{(k)} \rangle \right]. \end{aligned} \quad (5.12.3)$$

The factors $e_k/2\pi$ come from the replacement of $du^{(k)}$ with $dA^{(k)}$ for the free variables; for the constrained variables the replacement is compensated by a removal of the $e_k/2\pi$ factor from the δ -functions, see (4.6)–(4.8).

We calculate (5.12.3) by means of a translation

$$A^{(k)} = A^{(k)'} - A_{10}^{(k)c**} Q^{s*} Q A_{10}^{(k)c*} A^{(k)}, \quad (5.12.4)$$

which removes the dependence on $A_{10}^{(k)c*} A^{(k)}$ in the δ -functions. In fact, $\delta_{\text{Ax}, A_{10}^{(k)}}(A^{(k)}) = \delta_{\text{Ax}, A_{10}^{(k)'}}(A^{(k)'})$, and since $Q Q^{s*} = I$,

$$\delta_{A_{10}^{(k)c**} c} (Q A^{(k)}) = \delta_{A_{10}^{(k)'c**} c} (Q A_{10}^{(k)'c**} A^{(k)'}).$$

The fourth quadratic form above is obtained by collecting the terms in the exponential quadratic in $A_{10}^{(k)c*} A^{(k)}$. There remains a linear form

$$\langle A_{10}^{(k)c**} A^{(k)'}, \partial^* \sigma_{k, \text{loc}} \partial (I - A_{10}^{(k)c**} Q^{s*} Q) A_{10}^{(k)c*} A^{(k)} \rangle, \quad (5.12.5)$$

whose expectation in the Gaussian

$$\exp\left(-\frac{1}{2}\langle A_{10}^{(k)c**c} A^{(k)'}, \partial^* \sigma_{k, \text{loc}} \partial A_{10}^{(k)c**c} A^{(k)'} \rangle\right)$$

gives rise to the fifth form.

We remove the nonlocality in the third and fifth quadratic forms with random walk expansions for $C_{A_{10}^{(k)}}^{(k)}(u_{k+1})$ and for $C_{A_{10}^{(k)c**c}}^{(k)}$ as in (2.45) and (2.48). These obey the usual estimates. We denote the first and second quadratic forms by \mathcal{Q}_7 , and the other three [with $C_{A_{10}^{(k)}}^{(k)}(u_{k+1})$, $C_{A_{10}^{(k)c**c}}^{(k)}$ replaced with $C_{A_{10}^{(k), \text{loc}}}^{(k)}(u_{k+1})$, $C_{A_{10}^{(k)c**c, \text{loc}}}^{(k)}$] by \mathcal{Q}_8 . Altogether the exponential in (5.12.1) has been written as

$$\exp\left[-\mathcal{Q}_7 - \mathcal{Q}_8 - \sum_X W_5^{(k)}(X)\right].$$

Here $W_5^{(k)}(X)$ contains the terms with $C_{A_{10}^{(k)}, X}^{(k)}(u_{k+1})$ or with $C_{A_{10}^{(k)c**c}, X}^{(k)}$, and satisfies $|W_5^{(k)}(X)| \leq e^{-cr(e_k)|X|}$. The quadratic forms in \mathcal{Q}_7 and \mathcal{Q}_8 are localized near $A_{10}^{(k)c}$.

We make the same translation (5.12.4) in both numerator and denominator of the normalized integral in $A_{10}^{(k)}$. Terms quadratic in $A_{10}^{(k)} A^{(k)}$ cancel, but we still have the linear forms (5.12.5) and $\langle A_{10}^{(k)} \phi^{(k)}, A_{k, \text{loc}}(u_{k+1}) A_{10}^{(k)c} \phi^{(k)} \rangle$ as in our last calculation. We remove most of these forms with localized translations

$$\begin{aligned} A^{(k)'} &= A^{(k)''} - C_{A_{10}^{(k)c**c, \text{loc}}}^{(k)} \partial^* \sigma_{k, \text{loc}} \partial (I - L^{-1} A_{10}^{(k)c**c} Q^{s**} Q) A_{10}^{(k)c*} A^{(k)}, \\ \phi^{(k)} &= \phi^{(k)''} - C_{A_{10}^{(k), \text{loc}}}^{(k)}(u_{k+1}) A_{k, \text{loc}}(u_{k+1}) A_{10}^{(k)c} \phi^{(k)}. \end{aligned} \quad (5.12.6)$$

Terms quadratic in $A_{10}^{(k)c*} A^{(k)}$ or $A_{10}^{(k)c} \phi^{(k)}$ cancel as before, leaving the following integral:

$$\int d\mu_{A_{10}^{(k)}}^{(k)}(A^{(k)'}, \phi^{(k)'}) \chi_{A_{10}^{(k)}} \prod_{\sigma_1 \in \bar{\sigma}_1} F_{k, \text{loc}}^m(X_{\sigma_1}) e^{-V^{(k)}(A_8^{(k)}, u_{k+1}, A^{(k)}, \phi^{(k)})}.$$

Here $d\mu_{A_{10}^{(k)}}^{(k)}$ is an uncentered, normalized Gaussian measure,

$$\begin{aligned} & d\mu_{A_{10}^{(k)}}^{(k)}(A^{(k)'}, \phi^{(k)'}) \\ &= \frac{1}{\mathcal{N}} dA^{(k)''} |_{A_{10}^{(k)c**c}} d\phi^{(k)''} |_{A_{10}^{(k)}} \delta_{A_X, A_{10}^{(k)'}}(A^{(k)'}) \delta_{A_{10}^{(k)'c**c}}(Q A_{10}^{(k)c**c} A^{(k)'}) \\ &\quad \times \exp\left[-\frac{1}{2}\langle A_{10}^{(k)c**c} A^{(k)'}, \partial^* \sigma_{k, \text{loc}} \partial A_{10}^{(k)c**c} A^{(k)'} \rangle\right. \\ &\quad - \frac{1}{2}\langle A_{10}^{(k)} \phi^{(k)'}, (A_{k, \text{loc}}(u_{k+1}) + aL^{-2} P(u_{k+1})) A_{10}^{(k)} \phi^{(k)'} \rangle \\ &\quad - \langle A_{10}^{(k)c**c} A^{(k)'}, (I - \partial^* \sigma_{k, \text{loc}} \partial C_{A_{10}^{(k)c**c, \text{loc}}}^{(k)}) \partial^* \sigma_{k, \text{loc}} \partial (I - L^{-1} A_{10}^{(k)c**c} Q^{s**} Q) A_{10}^{(k)c*} A^{(k)'} \rangle \\ &\quad - \langle A_{10}^{(k)} \phi^{(k)'}, (I - (A_{k, \text{loc}}(u_{k+1}) + aL^{-2} P(u_{k+1})) \\ &\quad \times C_{A_{10}^{(k), \text{loc}}}^{(k)}(u_{k+1}) A_{k, \text{loc}}(u_{k+1}) A_{10}^{(k)c} \phi^{(k)'}) \rangle \left. \right]. \end{aligned} \quad (5.12.7)$$

This measure has covariances $C_{A_{10}^{(k)c**c}}^{(k)}$, $C_{A_{10}^{(k)}}^{(k)}(u_{k+1})$, and nonzero means reflecting the terms linear in $A_{10}^{(k)} \phi^{(k)'}$ or $A_{10}^{(k)c**c} A^{(k)'}$.

After the conditioning our density assumes the following form:

$$\begin{aligned}
& Q_{k+1}^L(v, \psi) \\
&= \sum_{\{X_\omega\}} \sum_{A_0^{(k)}} \sum_{S_4} \sum_{\tilde{\sigma}_1} \sum_{A_9^{(k)}} \int \prod_{j=0}^k du^{(j)}|_{A_{10}^{(j)c}} d\phi^{(k)}|_{A_{10}^{(k)c}} \delta_{A_{\lambda, A_{10}^{(k)c}}}(u^{(k)}) \\
&\quad \times \delta_{A_1^{(k)'*c}}(v/Q u^{(k)}) \delta_{A_1^{(k)'*c} \cap A_{10}^{(k)'c*}} \left(\frac{e_k}{2\pi} Q A^{(k)} \right) \zeta_{A_0^{(k)c} \cap A_9^{(k)c}} \zeta_{A_1^{(k)'c} \cap A_0^{(k)'c}} \prod_{\omega} g_k(X_\omega) \\
&\quad \times \prod_{\sigma \notin \tilde{\sigma}_1} F'_{(k), \text{loc}}(X_\sigma) \prod_{(j, x_j, X) \in S_4} (e^{-W_{4,j}^{(k)}(x_j, X)} - 1) \prod_{j=0}^k [Z_{A_{10}^{(j)c}}^{(j)} Z_{A_{10}^{(j)'c}}^{(j)}(u_{k+1})] \\
&\quad \times \exp \left[-\frac{1}{2} \langle A_5^{(k)'**f}, \sigma_{k+1, \text{loc}}^L A_5^{(k)'**f} \rangle \right. \\
&\quad - \frac{1}{2} \langle A_8^{(k)'} \psi, A_{k+1, \text{loc}}^L(u_{k+1}) A_8^{(k)'} \psi \rangle - \sum_{i=1}^8 \mathcal{Q}_i \\
&\quad - \mathcal{E}_k - E^{(k)} - \mathcal{P}_{k, \text{loc}}(A_8^{k+1}, A_8^{(k)c}, \tilde{u}_{k+1}) - R^{(k)}(A_8^{(k)c}, u_{k+1}, \theta_k H_{k, \text{loc}} A^{(k)}) \\
&\quad \left. - Q^{(k)}(A_8^{(k)c}, u_{k+1}, \theta_k H_{k, \text{loc}} A^{(k)}) \right] \int d\mu_{A_{10}^{(k)'}}^{(k)}(A^{(k)'}, \phi^{(k)'}) \zeta_{A_{10}^{(k)'}} \prod_{\sigma_1 \in \tilde{\sigma}_1} F_{k, \text{loc}}^{\tilde{m}}(X_{\sigma_1}) \\
&\quad \times \exp \left[-V^{(k)}(A_8^{(k)}, u_{k+1}, A^{(k)}, \phi^{(k)}) - \sum_X W_5^{(k)}(X) \right]. \tag{5.12.8}
\end{aligned}$$

The next two sections will focus on deriving a cluster expansion for the $d\mu_{A_{10}^{(k)'}}^{(k)}$ integral in (5.12.8)

5.13. Decoupling of the Small Field Region

We give a cluster expansion for the $d\mu_{A_{10}^{(k)'}}^{(k)}$ integral in (5.12.8). The purpose is to remove the dependence of the small field integral on the boundary fields. The cluster expansion has two parts; Mayer expansion of the interaction, and interpolation of the covariances of $d\mu_{A_{10}^{(k)'}}$.

Let us divide $A_9^{(k)}$ into its elementary $r(e_k)$ -cubes $\square^{(\alpha)}$. We assign to $\square^{(\alpha)}$ all bonds $\langle x, x + e_\mu \rangle$ with $x \in \square^{(\alpha)}$. Note that $V^{(k)}(A_8^{(k)}, u_{k+1}, A^{(k)}, \phi^{(k)})$ involves $A^{(k)}|_{A_9^{(k)'}}$, $\phi^{(k)}|_{A_9^{(k)'}}$ only. Thus we localize the fields $\phi^{(k)}$, $A^{(k)}$ in $V^{(k)}$ by writing

$$\phi^{(k)} = \sum_{\alpha} \square^{(\alpha)} \phi^{(k)}, \quad A^{(k)} = \sum_{\alpha} \square^{(\alpha)} A^{(k)}.$$

We associate to any collection of localization cubes a smallest connected union of cubes containing them (call it Y). Summing over all terms in $V^{(k)}$ and over localizations giving rise to Y , we obtain a decomposition.

$$V^{(k)}(A_8^{(k)}, u_{k+1}, A^{(k)}, \phi^{(k)}) = \sum_Y V^{(k)}(Y) + V_{\text{const}}^{(k)}(A_8^{(k)}).$$

The last term includes all terms independent of $A^{(k)}$, $\phi^{(k)}$. We have an estimate $|V^{(k)}(Y)| \leq e^{\beta(L_c/\varepsilon_0)^{1/4-\alpha}}$. Note that Y contains at most a few cubes.

Next we Mayer-expand the interaction

$$\begin{aligned}
& \exp \left[-V^{(k)}(A_8^{(k)}, u_{k+1}, A^{(k)}, \phi^{(k)}) - \sum_X W_5^{(k)}(X) \right] \\
&= \sum_{S_Y} \sum_{S_5} e^{-V_{\text{const}}^{(k)}(A_8^{(k)})} \prod_{Y \in S_Y} (e^{-V^{(k)}(Y)} - 1) \prod_{X \in S_5} (e^{-W_5^{(k)}(X)} - 1).
\end{aligned}$$

Here $S_Y(S_S)$ is the set of all Y 's (X 's) that arise in a term in the Mayer expansion.

We decompose $A_{10}^{(k)}$ into elementary regions $\{\square_i\}_{i \in I}$ which are connected unions of $\square^{(\alpha)}$. Two $\square^{(\alpha)}$ are included into a \square_i if one of the following conditions hold:

- (i) They are both in some Y , $Y \in S_Y$,
- (ii) They are both in some X , $X \in S_S$,
- (iii) They both contain sites or bonds within $r(e_k)$ of some X_{σ_1} , $\sigma_1 \in \tilde{\sigma}_1$.
- (iv) They are both in a connected component of $A_{11}^{(k)c}$.

For the decoupling of the Gaussian measure, we interpolate the covariance with parameters $s_i \in [0, 1]$, $i \in I$, which turn off interactions between \square_i and \square_i^c . The factors $\delta((QA^{(k)''})(b'))$ in the measure constitute an interaction between blocks. It is convenient to treat them directly, so we trade them for a fictitious integration dB , where B is a field on $A_{10}^{(k)c*c}$. We insert $1 = \mathcal{N}^{-1} \int dB \exp(-1/2 \langle B, B \rangle)$ and translate $A^{(k)''}$ by $Q^{s*}B$ to obtain

$$\begin{aligned} & \int dA^{(k)''} |_{A_{10}^{(k)c*c}} \delta_{A_{10}^{(k)c*c}}(QA^{(k)c*c}A) \delta_{A_X, A_{10}^{(k)}}(A^{(k)''}) f(A^{(k)''}) \\ &= \mathcal{N}^{-1} \int dA^{(k)''} dB e^{-1/2 \langle B, B \rangle} \delta(QA^{(k)''} + QQ^{s*}B) \delta_{A_X}(A^{(k)''}) f(A^{(k)''} + Q^{s*}B). \end{aligned} \quad (5.13.1)$$

The translation does not affect δ_{A_X} , and by (I.2.19) we have $QQ^{s*} = I$. Integrating out B yields

$$\mathcal{N}^{-1} \int dA^{(k)''} \exp[-\frac{1}{2} \langle A^{(k)''}, Q^*QA^{(k)''} \rangle] \delta_{A_X}(A^{(k)''}) f((I - Q^{s*}Q)A^{(k)''}).$$

Thus we have a new quadratic form for $A_{10}^{(k)c*c}A^{(k)''}$, namely

$$Q^*Q + (I - Q^*Q^s)\partial^* \sigma_{k, \text{loc}} \partial (I - Q^{s*}Q). \quad (5.13.2)$$

This is still bounded below on the subspace determined by $\delta_{A_X}(A^{(k)''})$: our lower bound on $\partial^* \sigma_{k, \text{loc}} \partial$ implies a lower bound

$$\|QA^{(k)''}\|^2 + O(1)\|A^{(k)''} - Q^{s*}QA^{(k)''}\|^2 \geq O(1)\|A^{(k)''}\|^2.$$

Applying (5.13.1) to numerator and denominator of the $d\mu_{A_{10}^{(k)}}$ -integral in (5.12.8), the \mathcal{N} 's cancel, and we obtain

$$\begin{aligned} & \int d\mu_{A_{10}^{(k)}}(A^{(k)''}, \phi^{(k)''}) \chi_{A_7^{(k)}} \prod_{\sigma_1 \in \tilde{\sigma}_1} F_{k, \text{loc}}^{\tilde{m}}(X_{\sigma_1}) \prod_{Y \in S_Y} (e^{-V^{(k)}(Y)} - 1) \\ & \times \prod_{X \in S_S} (e^{-W_S^{(k)}(X)} - 1) = \left\langle \prod_{i \in I} f(\square_i) \right\rangle_{\underline{1}}, \end{aligned}$$

where $f(\square_i)$ is the product of all the factors under the $d\mu_{A_{10}^{(k)}}$ integral above that are localized in \square_i . (Factors localized in $A_{10}^{(k)c}$ are assigned to the \square_i intersecting the corresponding component of $A_{11}^{(k)c}$.) Our construction of the \square_i ensures no overlap of factors between different \square_i 's. Everywhere $A^{(k)''}$ appears as $(I - Q^{s*}Q)A^{(k)''}$. The expectation $\langle \cdot \rangle_{\underline{1}}$ is in the measure

$$\frac{1}{\mathcal{N}'} d\Phi |_{A_{10}^{(k)}} \delta_{A_X, A_{10}^{(k)}}(A^{(k)''}) \exp[\frac{1}{2} \langle \Phi, \Delta \Phi \rangle + \langle \Phi, \mathcal{F} \rangle].$$

We have simplified the notation by writing $\Phi = (A^{(k)''}, \phi^{(k)''})$, $d\Phi |_{A_{10}^{(k)''}} = dA^{(k)''} |_{A_{10}^{(k)c*c}} d\phi^{(k)''} |_{A_{10}^{(k)}}$, $A_{10}^{(k)c} \Phi = (A_{10}^{(k)c} A^{(k)''}, A_{10}^{(k)c} \phi^{(k)''})$, and so on. The quadratic

and linear forms Δ and \mathcal{F} are obtained in the obvious fashion from (5.12.7), replacing $A_{10}^{(k)c*}A^{(k)''}$ with $(1-Q^{s*}Q)A_{10}^{(k)c*}A^{(k)''}$. The linear form is localized near the boundary of $A_{10}^{(k)}$.

To preserve positivity and boundedness properties of the inverse covariance, we define our s -dependent inverse covariance by taking convex combinations of inverse covariances with Dirichlet boundary conditions. For an arbitrary subset Γ of I we define Dirichlet forms:

$$\Delta_\Gamma = \sum_{i \in \Lambda} \square_i \Delta \square_i + \square^c \Delta \square^c,$$

where $\square^c = \bigcup_{i \notin \Gamma} \square_i$, and all operators are restricted to the subspace $A^{(k)''}(\Gamma_{y,x})=0$.

Next we define an operation

$$a_\Gamma \Delta_\Gamma = \Delta_{\Gamma \cup \Gamma'},$$

and we define a quadratic form for $\underline{s} = \{s_i\}_{i \in I}$:

$$\Delta_{\underline{s}} = \prod_{i \in I} [(1-s_i)a_i + s_i] \Delta = \sum_{\Gamma \subset I} \prod_{i \in \Gamma} (1-s_i) \prod_{i \in I \setminus \Gamma} s_i \Delta_\Gamma.$$

Note that by resumming the expansion above and using the fact that for $i'' \neq i$ or i' , or for $i=i'$, $\square_i(a_{\square_{i''}} \Delta) \square_{i'} = \square_i \Delta \square_{i'}$, we obtain that

$$\begin{aligned} \square_i \Delta_{\underline{s}} \square_{i'} &= s_i s_{i'} \square_i \Delta \square_{i'}, \quad i' \neq i \\ \square_i \Delta_{\underline{s}} \square_i &= \square_i \Delta \square_i. \end{aligned}$$

Using the theorem on unit lattice operators in [6], we can invert this operator to yield an exponentially decaying covariance $C_{\underline{s}} = (-\Delta_{\underline{s}})^{-1}$.

To give our expansion, we use the fundamental theorem of calculus to write

$$\left\langle \prod_{i \in I} f(\square_i) \right\rangle_{\underline{s}} = \sum_{\Gamma \subset I} \int d\underline{s}_\Gamma \frac{\partial}{\partial \underline{s}_\Gamma} \left\langle \prod_{i \in I} f(\square_i) \right\rangle_{\underline{s}_\Gamma}.$$

Here \underline{s}_Γ specifies $s_i=0$ for $i \notin \Gamma$, $d\underline{s}_\Gamma = \prod_{i \in \Gamma} ds_i$, $\partial/\partial \underline{s}_\Gamma = \prod_{i \in \Gamma} d/ds_i$, and $\langle \cdot \rangle_{\underline{s}_\Gamma}$ is the expectation with quadratic form $\Delta_{\underline{s}_\Gamma}$ instead of Δ . To calculate the s -derivatives, note that the first derivative produces a term

$$\left\langle \sum_{j \neq i} s_j \langle \square_i \Phi, \Delta \square_j \Phi \rangle; \prod_{i \in I} f(\square_i) \right\rangle_{\underline{s}_\Gamma}.$$

Subsequent derivatives either hit factors s_j already pulled down or bring new terms down with new truncations. After all derivatives are performed, we set the remaining s_j to zero, so only terms with no s_j multiplying them survive. The result is

$$\left\langle \prod_{i \in I} f(\square_i) \right\rangle_{\underline{s}} = \sum_{\Gamma \subset I} \int d\underline{s}_\Gamma \sum_{\substack{\text{pairings } p = \{p_\gamma\} \text{ of } \Gamma \\ p_\gamma = \{i_\gamma, j_\gamma\}}} \left\langle \prod_{\gamma=1}^{|\Gamma|/2} [\langle \square_{i_\gamma} \Phi, \Delta \square_{j_\gamma} \Phi \rangle]; \prod_{i \in I} f(\square_i) \right\rangle_{\underline{s}_\Gamma}.$$

Recall that we have a linear term in the measure, $e^{\langle \Phi, \mathcal{F} \rangle}$. With this term, integration by parts replaces Φ by $C_{\underline{s}}(\delta/\delta\Phi) + C_{\underline{s}}\mathcal{F}$ (see Eqs. (12.2), (12.3) of [3] where a similar expansion is used). We integrate by parts all fields appearing in this formula. Each Φ contracts through a $C_{\underline{s}}$ to another Φ , to an $f(\square_i)$, or to \mathcal{F} . If a

closed loop forms, or if a train of covariances beginning and ending in \mathcal{F} forms, then the term disappears with truncation. Thus we have only trains beginning with a $\delta/\delta\Phi$ and ending in either $\delta/\delta\Phi$ or \mathcal{F} . The sum over pairings and the sum over ways of arranging the contractions combine into a sum over walks $\{\omega_x\}_{x \in \pi}$, $\omega = (i_1, i_2, \dots, i_{|\omega|})$ involving the sites in α , an element of a partition π of Γ . Thus letting $\mathcal{P}(\Gamma)$ denote the partitions of Γ , we have

$$\left\langle \prod_{i \in I} f(\square_i) \right\rangle_{\underline{\Delta I}} = \sum_{\Gamma \subset I} \int dS_{\Gamma} \sum_{\pi \in \mathcal{P}(\Gamma)} \left\langle \prod_{x \in \pi} \left[\sum_{\omega(x)} \left\langle \frac{\delta}{\delta\Phi}, C_{\underline{\Delta}} \square_{i_1} A \square_{i_2} C_{\underline{\Delta}} \square_{i_3} A \square_{i_4} \dots A \square_{i_{|\omega(x)|}} C_{\underline{\Delta}} \left(\frac{1}{2} \frac{\delta}{\delta\Phi} + \mathcal{F} \right) \right\rangle \right] \prod_{i \in I} f(\square_i) \right\rangle_{\underline{\Delta I}}. \tag{5.13.3}$$

The $1/2$ for the $\delta^2/\delta\Phi^2$ term compensates for the fact that we count a walk as being different from its reverse. The combinatoric structure of (5.13.3) is very similar to that of the GJS cluster expansion [9].

Let us examine the factorization properties of this expansion. The form A has a range less than $1/2r(e_k)$. The $f(\square_i)$ do not couple different \square_i . Hence only adjacent \square_i with $s_i \neq 0$ interact in the above formula. Thus our expression for $d/dS_{\Gamma} \left\langle \prod_{i \in I} f(\square_i) \right\rangle_{\underline{\Delta I}}$ factorizes over the connected components of Γ . (Here we say that \square_i is connected to $\square_{i'}$ if they abut on a hypersurface of any dimension.) The expression also factorizes over the $\square_{i'}$, $i \in I \setminus \Gamma$. Call the factorization regions clusters.

It is worth mentioning here that only clusters intersecting $A_{11}^{(k)c}$ have any dependence on $A_{10}^{(k)c} \phi^{(k)}$, $A_{10}^{(k)c*} A^{(k)}$. This is because \mathcal{F} , $A^{(k)} - A^{(k)'}$, $\phi^{(k)} - \phi^{(k)'}$ are nonzero only in $A_{11}^{(k)c}$. Thus we have finally decoupled clusters that do not intersect $A_{11}^{(k)c}$ from the large field regions - at least in so far as the fields $u^{(k)}$, $\phi^{(k)}$ are concerned. There is still dependence on the block fields v , ψ which have yet to be integrated over and decoupled. We denote by $A_{12}^{(k)}$ the set of sites in clusters not intersecting $A_{11}^{(k)c}$.

We give now the expression for the polymer activities of this expansion. Given some region X , a union of \square_i , we sum Γ over all subsets of $\{i \in I : \square_i \subset X\}$, such that X is a single cluster. Writing

$$C_{\omega} = C_{\underline{\Delta}} \square_{i_1} A \square_{i_2} C_{\underline{\Delta}} \square_{i_3} \dots \square_{i_{|\omega|}} C_{\underline{\Delta}},$$

we have

$$g_1(X) = \sum_{\Gamma} \int dS_{\Gamma} \sum_{\pi \in \mathcal{P}(\Gamma)} \left\langle \prod_{x \in \pi} \left[\sum_{\omega(x)} \left\langle \frac{\delta}{\delta\Phi}, C_{\omega(x)} \left(\frac{1}{2} \frac{\delta}{\delta\Phi} + \mathcal{F} \right) \right\rangle \right] \prod_{\square_i \subset X} f(\square_i) \right\rangle_{\underline{\Delta I}, X}.$$

Here $\underline{\Delta} = \{s_i : \square_i \subset X\}$, and $\langle \cdot \rangle_{\underline{\Delta I}, X}$ is defined by integrating over the fields in X only.

We obtain the following expressions for the $d\mu_{A_{10}^{(k)c}}^{(k)}$ -integral in (5.12.8):

$$\sum_{S_Y} \sum_{S_S} e^{-V_{\text{const}}^{(k)}(A_{10}^{(k)c})} \sum_{\{X_x\} \text{ filling } A_{10}^{(k)c}} \prod_x g_1(X_x) = e^{-V_{\text{const}}^{(k)}(A_{10}^{(k)c})} \sum_{\{X_x\}} \prod_x g_2(X_x). \tag{5.13.4}$$

Here $g_2(X_x)$ is obtained by summing over S_Y, S_S compatible with X_x (each Y, X is contained in X_x or the corresponding component of $A_{11}^{(k)c}$):

$$g_2(X_x) = \sum_{S_Y, S_S \text{ compatible with } X_x} g_1(X_x).$$

Let us estimate $g_2(X_x)$ now. Each time some cubes are joined into one \square_i by an $e^{-V^{(k)}(Y)} - 1$ or an $e^{-W_s^{(k)}(X)} - 1$, we get a factor $e^{\beta(L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha}}$ or $e^{-cr(e_k)}$. Each time some \square_i 's are joined, we have s -derivatives, which produce functional derivatives, chains of covariances $C_{\omega(x)}$, and factors $\mathcal{F} = O(e^{-cr(e_k)})$. Functional derivatives hitting χ -factors farther than $1/2r(e_k)$ from $A_{10}^{(k)c}$ produce factors $e^{-cp(e_k)^2}$ after integrating with respect to $A^{(k)'}$, $\phi^{(k)'}$. These derivatives are supported at $|A^{(k)'}| \geq cp(e_k)$ or $|\phi^{(k)'}| \geq cp(e_k)$ (here we use the fact that the translation vanishes). Thus we can use the arguments at the end of Sect. 14 in [3] to extract the factors $e^{-cp(e_k)^2}$ from the Gaussian measure. Functional derivatives hitting $e^{-V^{(k)}(Y)}$ yield factors $e^{\beta(L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha}}$. Functional derivatives hitting χ' -factors within $\frac{1}{2}r(e_k)$ of $A_{10}^{(k)c}$ are connected through $C_{\omega(x)}$ to $A_{11}^{(k)}$, so we get small factors $e^{-cr(e_k)}$ from the exponential decay of the operators C_s and Λ in $C_{\omega(x)}$. Altogether we have small factors at each end of $C_{\omega(x)}$ (except for contractions to $F_{k, \text{loc}}^{\tilde{m}}(X_{\sigma_1})$). If the walk $\omega(x)$ wanders through more than a few cubes, we begin to pickup factors $e^{-cr(e_k)}$. These control the sum over walks and partitions, and the factorials, as in [9]. (Factorials can be produced when many functional derivatives hit the same object, for example a characteristic function.)

Altogether, we typically get at least a small power of $e^{\beta(L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha}}$ in every cube of X_x . The exceptions are when cubes are in a component of $A_{11}^{(k)c}$, when they support some $F_{k, \text{loc}}^{\tilde{m}}(X_{\sigma_1})$, or when X_x is a single cube. We must allow for divergent factors such as $(L^k \varepsilon)^{-\tilde{m}}$ at $F_{k, \text{loc}}^{\tilde{m}}(X_{\sigma_1})$, where \tilde{m} depends on F . Estimating the sums over S_Y, S_S , and the sums in the cluster expansion leads to combinatoric factors

$$\exp((e^{\beta(L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha}})^{\beta'} |X_x|), \quad \beta' > 0.$$

Such factors are easily beaten by the small factors described above for nonexceptional cubes. For the cubes in $A_{11}^{(k)c}$ or for a single cube, we have to include the proper volume factor in our final estimate.

In sum, we have the following bound on $g_2(X_x)$:

$$\begin{aligned} |g_2(X_x)| \leq & \exp[(e^{\beta(L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha}})^{\beta'} (|X_x \cap A_{11}^{(k)c}| + 1)] \\ & \times \prod_{\sigma_1} (L^k \varepsilon)^{-\tilde{m}(\sigma_1)} (e^{\beta(L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha}})^{\beta' |X_x \setminus A_{11}^{(k)c}|}. \end{aligned}$$

The product over σ_1 runs over $\sigma_1 \in \tilde{\sigma}_1$ such that $X_{\sigma_1} \subset X_x$ or X_{σ_1} is in a component of $A_{11}^{(k)c}$ overlapping X_x . If $|X_x| = 1$, with no $F_{k, \text{loc}}^{\tilde{m}}$ -factors, then we have the more precise bound $|g_2(X_x) - 1| \leq e^{\beta(L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha}}$, obtained from the same estimates on the S_Y, S_S sums, and from extremely small factors when a χ' -factor is replaced by 1.

5.14. Resummation and Extraction of the Perturbation Expansion

The estimates in the last section show that the basic volume dependence or pressure for our expansion is naively of the order of $(e^{\beta(L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha}})^{\beta'}$. We need to do better in $A_{12}^{(k)}$, the region that has been decoupled from the large field regions. We improve our expansion in $A_{12}^{(k)}$ by computing the pressure and the expectation of $F_{k, \text{loc}}^{\tilde{m}}$ as perturbation series plus remainders of the order of $(e^{\beta(L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha}})^{\tilde{m} + 1}$, $(e^{\beta(L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha}})^{\tilde{m} + 1}$, respectively. The remainder terms are so small that they can be treated like the large field effects and ignored in the expansion at the next scale.

The perturbative terms exhibit renormalization cancellations, and so obey the bounds we need for the next step.

To extract the perturbative terms, we resum the decoupling and Mayer expansions in $A_{12}^{(k)}$. Note that $W_5^{(k)}(X) \neq 0$ only for X at the boundary of $A_{10}^{(k)}$. Thus $W_5^{(k)}$ -terms will not appear in the resummed expansion. We obtain for the expansion in (5.13.4)

$$\sum_{\{X_z\}} \prod_{\alpha} g_2(X_z) = \sum_{\{X_z\} \text{ overlapping } A_{11}^{(k)c}} \prod_{\alpha} g_2(X_z) z_F(A_{12}^{(k)}),$$

where

$$z_F(A_{12}^{(k)}) = \left\langle \chi'_{A_{12}^{(k)}} \prod_{\sigma_1: X_{\sigma_1} \subset A_{12}^{(k)}} F_{k,\text{loc}}^m(X_{\sigma_1}) e^{-\tilde{V}^{(k)}(A_{12}^{(k)})} \right\rangle_{1, A_{12}^{(k)}}, \tag{5.14.1}$$

$$\tilde{V}^{(k)}(A_{12}^{(k)}) = \sum_{Y \subset A_{12}^{(k)}} V^{(k)}(Y).$$

Recall that $A^{(k)''} = A^{(k)}$, $\phi^{(k)''} = \phi^{(k)}$ in $A_{12}^{(k)}$, and that $A^{(k)''}$ has been replaced by $(I - Q^{**}Q)A^{(k)''}$ everywhere in the integrand.

We treat $z_F(A_{12}^{(k)})$ as follows:

$$z_F(A_{12}^{(k)}) = \frac{z_F(A_{12}^{(k)})}{z(A_{12}^{(k)})} \exp(\log z(A_{12}^{(k)})),$$

where $z(A_{12}^{(k)}) = z_{F=1}(A_{12}^{(k)})$, and we give expansions for z_F/z and $\log z$. The first expansion will give rise to $F_{k+1,\text{loc}}^L$ plus remainders, the second to $\mathcal{P}_{k+1,\text{loc}}^L$ plus remainders. We consider only $\log z$ for the moment.

Define $z_t(A_{12}^{(k)})$ for $t \in [0, 1]$ by replacing $\tilde{V}(A_{12}^{(k)})$ with $t\tilde{V}(A_{12}^{(k)})$, replacing $\chi(cp(e_k), (I - Q^{**}Q)A^{(k)})$ with $\chi(cp(te_k), (I - Q^{**}Q)A^{(k)})$, and similarly for $\chi(cp(e_k), \phi^{(k)})$. Thus the restrictions and the interactions disappear at $t = 0$, at which point we have a purely Gaussian expectation.

Thus we define perturbative terms for the action,

$$\tilde{\mathcal{P}}_{k+1}(A_{12}^{(k)}) = \sum_{\alpha=1}^{\bar{n}} -\frac{1}{\alpha!} \frac{d^\alpha}{dt^\alpha} \log z_t(A_{12}^{(k)})|_{t=0},$$

and a remainder

$$\mathcal{R}_k(A_{12}^{(k)}) = \int_{\sigma}^1 dt -\frac{(1-t)^{\bar{n}}}{(\bar{n}+1)!} \left\langle \frac{d}{dt}; \dots; \frac{d}{dt} \right\rangle_t. \tag{5.14.2}$$

Here $\langle \cdot \rangle_t$ is the interacting expectation

$$\langle \cdot \rangle_t = \frac{1}{z_t(A_{12}^{(k)})} \langle \cdot \chi'_{A_{12}^{(k)},t} e^{-t\tilde{V}_{12}^{(k)}(A^{(k)})} \rangle_{1, A_{12}^{(k)}},$$

with $\chi'_{A_{12}^{(k)},t}$ defined as above replacing $p(e_k)$ with $p(te_k)$.

We express each d/dt as a sum $\sum_{\gamma} (d/dt)_{\gamma}$, where $(d/dt)_{\gamma}$ acts only on the t before a particular term $V^{(k)}(Y)$ in $\tilde{V}^{(k)}$ or in a particular χ -factor. We cluster expand as before each integral making up the truncated expectation values $\langle (d/dt)_{\gamma_1}; \dots; (d/dt)_{\gamma_{\bar{n}+1}} \rangle_t$. Let $H \subset \{1, \dots, \bar{n}+1\}$ specify which observables are included in one of the integrals. If $J \in H$ then we have a factor $(d/dt)_{\gamma_j}$ in the integral. The partition

$\{\square_i\}$ of $A_{12}^{(k)}$ is determined by the sets Y from $e^{-tV^{(k)}(Y)} - 1$ factors, and by sets Y from $V^{(k)}(Y)$ factors differentiated down. The expansion takes the form

$$\left\langle \prod_{j \in H} \left(\frac{d}{dt} \right)_{\gamma_j} \chi'_{A_{12}^{(k)}, t} e^{-t\tilde{V}^{(k)}(A_{12}^{(k)})} \right\rangle_{1, A_{12}^{(k)}} = \sum_{\{X_\alpha\} \text{ filling } A_{12}^{(k)}} \prod_{\beta} g_3(H_\beta, X_\beta). \quad (5.14.3)$$

Here $H_\beta \subset H$ specifies which $(d/dt)_{\gamma_j}$ have supports intersecting X_β .

The polymer activity g_3 is essentially the same as g_2 , but with additional observables, namely the $(d/dt)_{\gamma_j}$ -factors determined by H_β . Also, the interaction and characteristic functions have been partially interpolated away, and there are not $W_5^{(k)}$ -terms.

If $|X_\beta| = 1$, $H_\beta = \emptyset$, we write $g_3(\emptyset, X_\beta) = 1 + g'_3(\emptyset, X_\beta)$ and the above expansion holds again, but without the condition that $\{X_\beta\}$ fill $A_{12}^{(k)}$. The $\{X_\beta\}$ must cover all cubes connected with the $(d/dt)_{\gamma_j}$, $j \in H$. Let us drop the prime, and prove that

$$|g_3(H_\beta, X_\beta)| \leq (e^\beta (L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha})^{||H_\beta| + \beta^+ |X_\beta \cap H_\beta||}. \quad (5.14.4)$$

We use $X_\beta \setminus H_\beta$ to denote the set of cubes with no $(d/dt)_{\gamma_j}$ factors, $j \in H_\beta$.

The proof of this estimate is similar to the one for g_2 . We mention only the new features. Each factor $V^{(k)}(Y)$ in $\prod_{j \in H_\beta} (d/dt)_{\gamma_j}$ produces a factor $e^\beta (L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha}$ in the final estimate. This is obtained in the Gaussian integration estimate, using the fact that $V^{(k)}(Y)$ is a small polynomial in $A^{(k)}$, $\phi^{(k)}$. [The restrictions disappear as $t \rightarrow 0$, so $V^{(k)}(Y)$ cannot be replaced by its supremum.] The factors $e^{-tV^{(k)}(Y)} - 1$ can be bounded as before, because the coefficient t in front of $V^{(k)}(Y)$ plus a small power of e_k easily beat the bounds $A^{(k)}$, $\phi^{(k)} \leq cp(e_k)$. Each t -derivative of a χ -factor in $\chi_{A_{12}^{(k)}, t}$ gives at least a factor $e^\beta (L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha}$. This follows because with $\chi'(1, x) \equiv d/dx \chi(1, x)$, we have

$$\begin{aligned} \left| \frac{d}{dt} \chi(cp(te_k), A^{(k)}) \right| &= \left| \frac{A^{(k)}}{cp(te_k)^2} \left(\frac{d}{dt} p(te_k) \right) \chi'(1, A^{(k)}/cp(te_k)) \right| \\ &\leq ct^{-1} |\chi'(1, A^{(k)}/cp(te_k))|, \end{aligned}$$

and similarly the n -th derivative in t of $\chi(cp(e_k), A^{(k)})$ is bounded by t^{-n} times a function bounded by a constant and supported in $c_1 p(te_k) \leq |A^{(k)}| \leq c_2 p(te_k)$. After integration over $A^{(k)}$, we obtain factors $ct^{-n} e^{-cp(te_k)^2} \leq (e^\beta (L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha})^n$. Similar bounds hold for $\phi^{(k)}$. The bound for $H_\beta = \emptyset$, $|X_\beta| = 1$ was obtained for g_2 , and the same proof applies here.

Returning to our expansion, let us sum first over $\{H_\gamma\}$, the partition of H determined by the $\{X_\beta\}$. Denote the X_β 's with $H_\beta \neq \emptyset$ by X_γ ; the X_β with $H_\beta = \emptyset$ by Y_δ . The expansion (5.14.2) becomes

$$\sum_{\{H_\gamma\} \in \mathcal{P}(H)} \sum_{\{X_\gamma\}, \{Y_\delta\} \text{ nonoverlapping}} \prod_{\gamma} g_3(H_\gamma, X_\gamma) \prod_{\delta} g_3(\emptyset, Y_\delta).$$

Each X_γ must cover and connect all the t -derivatives specified by H_γ . Next we reorganize this expansion in order to extract the truncated expectation values (5.14.2). This involves adding and subtracting terms in a scheme familiar to one in [10]. We insert factors

$$u(X, Y) = \begin{cases} 0 & \text{if } X, Y \text{ overlap} \\ 1 & \text{if } X, Y \text{ do not overlap,} \end{cases}$$

and similarly factors $u(X_1, X_2), u(Y_1, Y_2)$. We extend the sums over $\{X_\gamma\}, \{Y_\delta\}$ to nonoverlapping sets; however the corresponding subsets of H remain the same – no duplication of t -derivatives. We put $u = 1 + a$ and expand in the usual manner. This enables us to factor out the normalization $z_t(A_{12}^{(k)})$ to obtain

$$\left\langle \prod_{j \in H} \left(\frac{d}{dt} \right)_{\gamma_j} \right\rangle_t = \sum_{\{H_\gamma\} \in \mathcal{P}(H)} \sum_{\{X_\gamma\}} \sum_{(Y_1, \dots, Y_B)} \\ \times \frac{1}{B!} \sum_G \prod_{\mathcal{L} \in G} a(\mathcal{L}) \prod_{\gamma} g_3(H_\gamma, X_\gamma) \prod_{\delta=1}^B g_3(\phi, Y_\delta).$$

Here \mathcal{L} denotes pairs of clusters (lines) and G runs over graphs of such lines in which each Y_δ is connected directly or indirectly to some X_γ . The connected components of G define a partition of H which corresponds to the partition in the formula

$$\left\langle \prod_{j \in H} \left(\frac{d}{dt} \right)_{\gamma_j} \right\rangle_t = \sum_{\{H_\tau\} \in \mathcal{P}(H)} \prod_{\tau} \left\langle \prod_{j \in H_\tau} \left[; \left(\frac{d}{dt} \right)_{\gamma_j} \right] \right\rangle_t.$$

Thus we have a formula

$$\left\langle \prod_{j \in H} \left[; \left(\frac{d}{dt} \right)_{\gamma_j} \right] \right\rangle_t = \sum_{\{H_\gamma\} \in \mathcal{P}(H)} \sum_{\{X_\gamma\}} \sum_{(Y_1, \dots, Y_B)} \\ \times \frac{1}{B!} \sum_{G_c} \prod_{\mathcal{L} \in G_c} a(\mathcal{L}) \prod_{\gamma} g_3(H_\gamma, X_\gamma) \prod_{\delta=1}^B g_3(\phi, Y_\delta),$$

where G_c runs over connected graphs involving all clusters X_γ, Y_δ , and hence all of H .

We use this to give an expansion for the remainder from the perturbation expansion of the interaction:

$$\mathcal{R}_k(A_{12}^{(k)}) = \sum_{X \subset A_{12}^{(k)}} W_6^{(k)'}(X).$$

Here $W_6^{(k)'}(X)$ is obtained by summing only over $\{X_{\gamma_j}\}, (Y_1, \dots, Y_B)$ which fill X , summing over $\{\gamma_{jj}\}$ with $\text{suppt}(d/dt)_{\gamma_j} \subset X$, and integrating over t as in (5.14.2). It is now a standard exercise to estimate the expansion, using (5.14.4). The result is

$$|W_6^{(k)'}(X)| \leq (e^\beta (L^k \varepsilon / \varepsilon_0)^{1/4 - \alpha})^{\bar{n} + 1 + \beta' |X|}.$$

(We allow adjustments in β, α, β' , keeping them small.)

We make some modifications in the perturbative terms to achieve the standard form of the interaction, $\mathcal{P}_{k+1, \text{loc}}^L$. We give random walk expansions for the propagators $C_{A_{12}^{(k)}}^{(k)}, C_{A_{12}^{(k)}}^{(k)}(u_{k+1})$ produced in this step. The leading terms, with only propagators $C_{A_{12}^{(k)}, \text{loc}}^{(k)}, C_{A_{12}^{(k)}, \text{loc}}^{(k)}(u_{k+1})$, we transform further. The others, localized in region X , have a factor of $e^{-cr(e_k)|X|}$. We also consider as remainders any terms whose order in λ and e is greater than \bar{n} .

We wish to replace $C_{A_{12}^{(k)}, \text{loc}}^{(k)}$ with $C_{\text{loc}}^{(k)}$. Recall that $C_A^{(k)}$ is the Dirichlet inverse to (5.13.2), and we define $C_{A, \text{loc}}^{(k)}$ by cutting off the kernel when the arguments are separated by $O(r(e_k))$. $C_{\text{loc}}^{(k)}$ was defined in (2.9), starting from the inverse to $\partial^* \sigma_{k, \text{loc}} \hat{c}$ on the appropriate subspace. The replacement of $C_{A_{12}^{(k)}, \text{loc}}^{(k)}$ with $C_{\mathbb{Z}^d, \text{loc}}^{(k)}$ produces

terms localized near $A_{12}^{(k)c}$. These terms are bounded by a small power of coupling constants, $e^\beta(L^k\varepsilon/\varepsilon_0)^{1/4-\alpha}$. To make the replacement of $C_{\mathbb{Z}^d, \text{loc}}^{(k)}$ with $C_{\text{loc}}^{(k)}$, note that the former always appears between two operators as $(I - Q^{S*}Q)C_{\mathbb{Z}^d, \text{loc}}^{(k)}(I - Q^*Q^S)$. This differs from $C_{\text{loc}}^{(k)}$ by a small, local operator, since after replacing $C_{\mathbb{Z}^d, \text{loc}}^{(k)}$ with $C_{\mathbb{Z}^d}^{(k)}$, we have an identity

$$(I - Q^{S*}Q)C_{\mathbb{Z}^d}^{(k)}(I - Q^*Q^S) = C^{(k)} = C_{\text{loc}}^{(k)} + O(e^{-cr(e_k)}).$$

Thus after removing some $O(e^{-cr(e_k)})$ remainders, we have our standard covariance $C_{\text{loc}}^{(k)}$.

We compose propagators, using also terms from $V_{\text{const}}^{(k)}(A_8^{(k)})$. For gauge field propagators, we apply (5.5.11). For scalar field propagators, we use the identity 2.42 from [7]:

$$\begin{aligned} G_k(\square, u_{k+1}) + \mathbf{1}^2 L^{-2} G_k(\square, u_{k+1}) Q_k^*(u_{k+1}) C^{(k)}(\square, u_{k+1}) \\ \times Q_k(u_{k+1}) G_k(\square, u_{k+1}) = G_{k+1}^\eta(\square, u_{k+1}). \end{aligned}$$

There are also small ($O(e^{-cr(e_k)})$) terms involving $G_{k, \text{loc}}(u_{k+1}) - G_k(\square, u_{k+1})$, $C_{\text{loc}}^{(k)}(u_{k+1}) - C^{(k)}(\square, u_{k+1})$, and $G_{k+1, \text{loc}}(u_{k+1}) - G_{k+1}(\square, u_{k+1})$, and boundary terms as above involving $C_{A_{12}^{(k)}, \text{loc}}^{(k)}(u_{k+1}) - C_{\text{loc}}^{(k)}(u_{k+1})$. We end up with scalar field propagators $G_{k+1, \text{loc}}^\eta(u_{k+1})$. For simplicity we extend the localizations of vertices in all diagrams back to $\bar{A}_8^{(k)}$ (for gauge fields we use a smooth localization function). This produces more boundary terms. Then the terms produced in this step combine with the old terms $V_{\text{const}}^{(k)}(A_8^{(k)})$ to produce the full interaction $\mathcal{P}_{k+1, \text{loc}}^L(A_8^{(k)})$. Altogether we have written

$$V_{\text{const}}^{(k)}(A_8^{(k)}) + \tilde{\mathcal{P}}_{k+1}(A_{12}^{(k)}) = \mathcal{P}_{k+1, \text{loc}}^L(A_8^{(k)}) + \sum_X W_6^{(k)'}(X).$$

If we put $W_6^{(k)}(X) = W_6^{(k)'}(X) + W_6^{(k)''}(X)$, then $W_6^{(k)}(X)$ obeys

$$|W_6^{(k)}(X)| \leq \begin{cases} (e^\beta(L^k\varepsilon/\varepsilon_0)^{1/4-\alpha})^{\bar{m}+1+\beta'|X|}, & \text{dist}(X, A_{12}^{(k),c}) \geq r(e_k) \\ (e^\beta(L^k\varepsilon/\varepsilon_0)^{1/4-\alpha})^{\beta'|X|}, & \text{otherwise.} \end{cases}$$

We apply a somewhat different procedure to extract the proper perturbative terms from the observable. We integrate by parts in the Gaussian expectation (5.14.1). Each $F_{k, \text{loc}}^{\bar{m}}(X_{\sigma_1})$ is a polynomial in $A^{(k)}$, $\phi^{(k)}$; those fields can be contracted via covariances $C_{A_{12}^{(k)}}^{(k)}$ or $C_{A_{12}^{(k)}, \text{loc}}^{(k)}(u_{k+1})$ to other observables, to $\chi'_{A_{12}^{(k)}}$, or to the interaction. After each integration by parts, we replace the covariance by $C_{\text{loc}}^{(k)}$ or $C_{\text{loc}}^{(k)}(u_{k+1})$ and give a random walk expansion for the difference. For each term, let \bar{X} be the union of the cubes covering the X_{σ_1} and the regions from the random walk expansion. A connected component of \bar{X} is called *complete* if a contraction to $\chi'_{A_{12}^{(k)}}$ occurs, if a term from the random walk expansion occurs, if at least $\bar{m}+1$ interactions have been differentiated down, or if the term is constant (all legs contracted). We stop integrating by parts fields in complete components of \bar{X} . After sufficiently many integrations by parts, all components of \bar{X} will be complete.

We break up the observable according to the connected components of \bar{X} . The components containing contractions to $\chi'_{A_{12}^{(k)}}$, terms from the random walk expansions, or at least $\bar{m}+1$ interactions are called remainder components $\{X_r\}$. The other components are called constant components $\{X_c\}$, since the observable

there is independent of $A^{(k)}, \phi^{(k)}$. We can arrange the construction so that the $\{X_c\}$ are determined once the remainder components are specified. Summing all possible diagrams in X_c gives the observable for the next step there, $F_{k+1,loc}^L(X_c)$. Summing all terms in $X_{r'}$ gives an observable $F_{k,rem}(X_{r'})$. Then the result of the integration by parts is

$$\frac{z_F}{z} = \left\langle \prod_{\sigma_1} F_{k,loc}^m(X_{\sigma_1}) \right\rangle_1 = \sum_{\{X_{r'}\}} \prod_c F_{k+1,loc}^L(X_c) \left\langle \prod_{r'} F_{k,rem}(X_{r'}) \right\rangle_1,$$

where $\langle \cdot \rangle_1$ is the interacting expectation at $t=1$.

Having extracted the desired perturbative terms $F_{k+1,loc}^L(X_c)$, we need to finish the calculation of the remainders by giving a cluster expansion for $\left\langle \prod_{r'} F_{k,rem}(X_{r'}) \right\rangle_1$, with appropriate bounds. We use essentially the same expansion as before, Mayer-expanding $V^{(k)}(Y)$'s and interpolating the Gaussian measure. Finally the polymer expansion $u=1+a$ permits us to factor out the normalization. Without going into details, it is clear that the result can be written in the following form:

$$\left\langle \prod_{\sigma_1} F_{k,loc}^m(X_{\sigma_1}) \right\rangle_1 = \sum_{\{X_{r'}\}} \prod_{r'} G_k(X_{r'}) \prod_{c: X_c \not\supseteq X_{r'}} F_{k+1,loc}^L(X_c).$$

The $X_{r'}$ are disjoint, and each one covers at least one X_{σ_1} , the support of one of the observables $F_{k,loc}^m$.

The main source of concern in estimating $G_k(X_{r'})$ is that we only have bounds $|F_{k,loc}(X_{\sigma_1})| \leq c(L^{k\varepsilon})^{-m(c)} e^{-m'(c)}$, coming from our estimates on perturbation expansions of observables; similarly for $F_{k+1,loc}^L(X_c)$. Here $m(c), m'(c)$ depend on the terms in F in X_{σ_1} or X_c . By performing sufficiently many integrations by parts, we have arranged for enough small factors to beat these large factors in the remainder terms (at least if $X_{r'}$ is not at the boundary of $A_{12}^{(k)}$). Near the boundary we have potentially large covariances $C_{A_{12}^{(k)},loc}^{(k)} - C_{loc}^{(k)}$ or $C_{A_{12}^{(k)},loc}(u_{k+1}) - C_{loc}(u_{k+1})$, so we make use of the proximity to $A_{12}^{(k)c}$ to provide the necessary convergence. These considerations lead to the following estimate:

$$|G_k(X)| \leq c(F(X)) (e^{\beta(L^k\varepsilon/\varepsilon_0)^{1/4-z}} \beta^{|X \setminus \cup X_c|} \times \prod_{X_{\sigma_1} \subset X: \text{dist}(X_{\sigma_1}, A_{12}^{(k)c}) < r(e_k)} [c(L^k\varepsilon)^{-m(c)} e^{-m'(c)}]).$$

To summarize the results of this section, we have

$$\begin{aligned} & e^{-V_c^{\text{const}}(A_8^{(k)})} \sum_{\{X_z\}} \prod_{\alpha} g_2(X_z) \\ &= \sum_{\{X_z\} \text{ overlapping } A_{12}^{(k)c}} \prod_{\alpha} g_2(X_z) \sum_{\{X_{r'}\}} \prod_{r'} G_k(X_{r'}) \\ & \times \prod_{c: X_c \not\supseteq X_{r'}} F_{k+1,loc}^L(X_c) \exp\left(-\mathcal{P}_{k+1,loc}^L(A_8^{(k)}) - \sum_X W_6^{(k)}(X)\right). \end{aligned} \tag{5.14.5}$$

5.15. Second Mayer Expansion and Scaling

In this section we recover the induction hypothesis for $k+1$ instead of k , and write a formula for the hole functional $g_{k+1}(X_\omega)$. First we Mayer-expand the irrelevant $W_6^{(k)}$ terms:

$$\exp\left(-\sum_X W_6^{(k)}(X)\right) = \sum_{S_6} \prod_{X \in S_6} (e^{-W_6^{(k)}(X)} - 1). \tag{5.15.1}$$

We define

$$A_{13}^{(k)} = A_{12}^{(k)} \setminus \bigcup_{X \in S_6} X \setminus \bigcup_{r'} X_{r'} \setminus \bigcup_{c: X_c \cap X \neq \emptyset, X \in S_6} X_c;$$

it is a region now completely free of irrelevant terms. We write

$$\prod_{c: X_c \not\subseteq \bigcup X_r} F_{k+1, \text{loc}}^L(X_c) = \prod_{c'} F_{k+1, \text{loc}}^L(X_{c'}) \prod_{\sigma'} F_{k+1, \text{loc}}^L(X_{\sigma'}), \tag{5.15.2}$$

where $\{\sigma'\} = \{c: X_c \subset A_{13}^{(k)}\}$, and $\{c'\}$ are the rest.

By inserting (5.14.5), (5.15.1), (5.15.2) in (5.12.8), we obtain the final form of the density $q_{k+1}^L(v, \psi)$.

We now scale this density from $T_L^{(k+1)}$ to $T_1^{(k+1)}$, putting $\psi^L(y) = L^{-(d-2)/2} \psi^1(L^{-1}y)$. If we define

$$q(v, \psi^1) = \exp \left[- \frac{d-2}{2} (\log L) |T_1^{(k+1)}| \right] q^L(v, L^{-(d-2)/2} \psi^1),$$

then the integral of $q(v, \psi^1)$ is equal to the integral of $q^L(v, \psi^L)$. Thus we define the $(k+1)$ th normalizing energy to be

$$\mathcal{E}_{k+1} = \mathcal{E}_k + E^{(k)} + \frac{d-2}{2} (\log L) |T_1^{(k+1)}|. \tag{5.15.3}$$

Let us describe how the scaling affects a few of the objects that will be needed in the next step. Defining $f^{(k+1)}(p) = (ie_{k+1})^{-1} \log v(p)$, we have that

$$e_k \eta L^{-2} \mathcal{D}_{k+1, \text{loc}}^\eta \hat{\partial}^{\eta*} Q_{k+1}^{e*} f = e_{k+1} L^{-1} \eta \mathcal{D}_{k+1, \text{loc}} \hat{\partial}^{L^{-1}\eta*} Q_{k+1}^{e*} f^{(k+1)},$$

and thus in $A_6^{(k)*}$ we have

$$u_{k+1, b} = (Q^{s*} v) \exp \left[-ie_k L^{-1} \eta \mathcal{D}_{k+1, \text{loc}} \hat{\partial}^{L^{-1}\eta*} Q_{k+1}^{e*} f^{(k+1)} \right]$$

as in the induction hypothesis (4.2). The quadratic forms become

$$\frac{1}{2} \langle A_5^{(k)'} ** f^{(k+1)}, \sigma_{k+1, \text{loc}} A_5^{(k)'} ** f^{(k+1)} \rangle + \frac{1}{2} \langle A_8^{(k)'} \psi, \Delta_{k+1, \text{loc}}(u_{k+1}) A_8^{(k)'} \psi \rangle.$$

The interaction and observables are scaled and written as $\mathcal{P}_{k+1, \text{loc}}(A_8^{(k)})$ and $F_{k+1, \text{loc}}(X_{\sigma'})$, respectively. Propagators and vertices appear scaled to the $L^{-1}\eta$ lattice. The scaled form of the normalization factors is given in (4.6), (4.9).

Let $\{X_{\omega'}\}$ be the components of $A_{13}^{(k)c}$, and let $X_{\omega'}$ also specify $A_x^{(k)c} \cap X_{\omega'}$ and a collection $\{X_{\omega'}\}$ of sets from the previous step. We exhibit the factorization of most of the terms in $q_{k+1}(v, \psi)$ by writing

$$\begin{aligned} q_{k+1}(v, \psi) &= \sum_{\{X_{\omega'}\}} \int \prod_{j=0}^k du^{(j)} \Big|_{A_{10}^{(j)c} * Q'_{k+1}(v, \psi, \{X_{\omega'}\}, \{u^{(j)}\})} \\ & q'_{k+1}(v, \psi, \{X_{\omega'}\}, \{u^{(j)}\}) \\ &= Z_{k+1, A_0^{(k)'}} \prod_{\omega'} g_{k+1}(X_{\omega'}) \prod_{\sigma'} F_{k+1, \text{loc}}(X_{\sigma'}) \prod_{j=0}^k [Z_{A_{10}^{(j)c} * c}^{(j)} Z_{A_{10}^{(j)}}^{(j)}(u_{k+1})] \\ & \times \exp \left[-\frac{1}{2} \langle A_5^{(k)'} ** f^{(k+1)}, \sigma_{k+1, \text{loc}} A_5^{(k)'} ** f^{(k+1)} \rangle \right. \\ & \left. - \frac{1}{2} \langle A_8^{(k)'} \psi, \Delta_{k+1, \text{loc}}(u_{k+1}) A_8^{(k)'} \psi \rangle - \mathcal{P}_{k+1, \text{loc}}(A_8^{(k)}) - \mathcal{E}_{k+1} \right], \end{aligned}$$

which is in the form of our original induction hypothesis, (4.1). The hole functional has the expression

$$\begin{aligned}
 g_{k+1}(X_{\omega'}) = & \sum_{S_4, \tilde{\sigma}_1, \tilde{A}_9^{(k)} \cap X_{\omega'}, \{X_{\alpha_2}, \{X, \nu\}, S_6 \text{ compatible with } X_{\omega'}} \\
 & \times \int d\phi^{(k)}|_{A_{10}^{(k)c} \cap X_{\omega'}} \delta_{A_X, A_{10}^{(k)c} \cap X_{\omega'}}(u^{(k)}) \delta_{A_1^{(k)'} * c \cap X_{\omega'}}(v/Q u^{(k)}) \\
 & \times \delta_{A_1^{(k)'} * c \cap A_{10}^{(k)c} * c \cap X_{\omega'}} \left(\frac{e_k}{2\pi} Q A^{(k)} \right) \zeta_{A_0^{(k)c} \cap X_{\omega'}} \zeta_{A_0^{(k)} \cap A_9^{(k)c} \cap X_{\omega'}} \zeta'_{\tilde{A}_9^{(k)c} \cap X_{\omega'}} \\
 & \times \prod_{\omega} g_k(X_{\omega}) \prod_{\sigma \neq \tilde{\sigma}, \tilde{X}_{\sigma} \subset X_{\omega'}} F'_{k, \text{loc}}(X_{\sigma}) \prod_{(j, X_j, X) \in S_4} (e^{-W_{4,j}^{(k)}(X_j, X)} - 1) \\
 & \times \prod_{\alpha} g_2(X_{\alpha}) \prod_{r'} G_k(X_{r'}) \prod_{X \in S_6} (e^{-W_6^{(k)}(X)} - 1) \prod_{c': X_{c'} \subset X_{\omega}} F_{k+1, \text{loc}}(X_{c'}) \\
 & \times \exp \left[- \sum_{i=1}^8 \mathcal{Q}_i(X_{\omega'}) - \mathcal{P}_{k, \text{loc}}(A_8^{(k-1)}, A_8^{(k)c} \cap X_{\omega'}, \tilde{u}_{k+1}) \right. \\
 & \left. - R^{(k)}(A_8^{(k)c} \cap X_{\omega'}, u_{k+1}, \theta_k H_{k, \text{loc}} A^{(k)}) \right. \\
 & \left. - Q^{(k)}(A_8^{(k)c} \cap X_{\omega'}, u_{k+1}, \theta_k H_{k, \text{loc}} A^{(k)}) \right]. \tag{5.15.4}
 \end{aligned}$$

Compatibility means that the summations run over sets associated only with $X_{\omega'}$ and that the sets would have given us $X_{\omega'}$ in the course of our constructions. Specifically, this implies a certain “density” of terms leading to convergence factors, and compatibility of the sets with the layered structure imposed by the $A_{\alpha}^{(k)}$.

We have discussed the estimates on many of the elements of the expansion in g_{k+1} . However, we cannot complete the estimates until after extracting convergence from the large field conditions. This is accomplished only after integrating over the final v, ψ in the last step. These problems, and the problem of decoupling of the final fields, will be considered in a subsequent paper.

References

1. Bałaban, T., Imbrie, J., Jaffe, A.: Exact renormalization group for gauge theories. In: Progress in gauge field theory. Cargèse 1983. Lehmann, G., 't Hooft, G., Jaffe, A., Mitter, P., Singer, I., Stora, R. (eds.) New York: Plenum 1984
2. Bałaban, T., Imbrie, J., Jaffe, A.: Renormalization of the Higgs model: Minimizers, propagators, and the stability of mean field theory. Commun. Math. Phys. **97**, 299–329 (1985)
3. Bałaban, T., Brydges, D., Imbrie, J., Jaffe, A.: The mass gap for Higgs models on a unit lattice. Ann. Phys. **158**, 281–319 (1984)
4. Imbrie, J.: Renormalization group methods in gauge field theories. In: Critical phenomena, random systems, gauge theories. Les Houches 1984. Osterwalder, K., Stora, R. (eds.) Amsterdam: North-Holland 1986
5. Bałaban, T.: (Higgs)_{2,3} quantum fields in a finite volume. III. Renormalization. Commun. Math. Phys. **88**, 411–445 (1983)
6. Bałaban, T.: Regularity and decay of lattice Green’s functions. Commun. Math. Phys. **89**, 571–597 (1983)
7. Bałaban, T.: (Higgs)_{2,3} quantum fields in a finite volume. I. A lower bound. Commun. Math. Phys. **85**, 603–636 (1983)
8. Bałaban, T.: (Higgs)_{2,3} quantum fields in a finite volume. II. An upper bound. Commun. Math. Phys. **86**, 555–594 (1983)

9. Glimm, J., Jaffe, A., Spencer, T.: The particle structure of the weakly coupled $\mathcal{P}(\phi)_2$ model and other applications of high temperature expansions. In: Constructive quantum field theory. Lecture Notes in Physics, Vol. 25. Velo, G., Wightman, A. (eds.). Berlin, Heidelberg, New York: Springer 1973
10. Gawedzki, K., Kupiainen, A.: Renormalization group for a critical lattice model. Effective interactions beyond the perturbation expansion in bounded spins approximation. *Commun. Math. Phys.* **88**, 77–94 (1983)

Communicated by A. Jaffe

Received March 24, 1987; in revised form May 22, 1987

