# THE RIEMANN HYPOTHESIS FOR PERIOD POLYNOMIALS OF MODULAR FORMS

#### SEOKHO JIN, WENJUN MA, KEN ONO, AND KANNAN SOUNDARARAJAN

For Don Zagier in celebration of his 65th birthday

ABSTRACT. The period polynomial  $r_f(z)$  for an even weight  $k \geq 4$  newform  $f \in S_k(\Gamma_0(N))$  is the generating function for the critical values of L(f,s). It has a functional equation relating  $r_f(z)$  to  $r_f\left(-\frac{1}{Nz}\right)$ . We prove the Riemann Hypothesis for these polynomials: that the zeros of  $r_f(z)$  lie on the circle  $|z| = 1/\sqrt{N}$ . We prove that these zeros are equidistributed when either k or N is large.

#### 1. Introduction

Let  $f \in S_k(\Gamma_0(N))$  be a newform [1, 2] of even weight k and level N. Associated to f is its L-function L(f, s), which has been normalized so that the completed L-function,

$$\Lambda(f,s) := \left(\frac{\sqrt{N}}{2\pi}\right)^s \Gamma(s) L(f,s),$$

satisfies the functional equation  $\Lambda(f,s) = \epsilon(f)\Lambda(f,k-s)$ , with  $\epsilon(f) = \pm 1$ . Recall that the completed L-function arises as a period integral of the newform f:

(1.1) 
$$\Lambda(f,s) = N^{s/2} \int_0^\infty f(iy) y^s \frac{dy}{y}.$$

The focus of this paper is the period polynomial associated to f, the degree k-2 polynomial

(1.2) 
$$r_f(z) := \int_0^{i\infty} f(\tau)(\tau - z)^{k-2} d\tau.$$

Expanding  $(\tau - z)^{k-2}$ , and using (1.1), we may also express the period polynomial by

(1.3) 
$$r_f(z) = i^{k-1} N^{-\frac{k-1}{2}} \sum_{n=0}^{k-2} {k-2 \choose n} (\sqrt{N}iz)^n \Lambda(f, k-1-n),$$

or equivalently as

(1.4) 
$$r_f(z) = -\frac{(k-2)!}{(2\pi i)^{k-1}} \sum_{n=0}^{k-2} \frac{(2\pi i z)^n}{n!} L(f, k-n-1).$$

Key words and phrases. Period polynomials, Modular forms.

The first author thanks KIAS for its generous support. The second author thanks the China Scholarship Council for its generous support. The third author thanks the support of the Asa Griggs Candler Fund and the NSF. The fourth author thanks the support of the NSF, and the Simons Foundation for a Simons Investigator Grant. The authors thank YoungJu Choie, Yuri Manin, Ram Murty, Ken Ribet, Drew Sutherland, and Don Zagier for useful comments and discussions.

<sup>2010</sup> Mathematics Subject Classification: 11F11, 11F67.

In other words,  $r_f(z)$  is a generating function for the critical values  $L(f, 1), L(f, 2), \ldots, L(f, k-1)$ . For general facts on period polynomials, the reader is encouraged to see [3, 4, 5, 6, 7]; other papers broadly related to the themes of this paper are [8, 9].

Using the functional equation  $\Lambda(f,s) = \epsilon(f)\Lambda(f,k-s)$  in (1.3), we find that

$$r_f(z) = -i^k \epsilon(f) (\sqrt{N}z)^{\frac{k-2}{2}} r_f \left(-\frac{1}{Nz}\right),$$

so that if  $\rho$  is a zero of  $r_f(z)$  then so is  $-1/(N\rho)$ . In analogy with the Riemann hypothesis, we may ask whether all the zeros of  $r_f(z)$  lie on the circle  $|\rho| = 1/\sqrt{N}$ . For Hecke eigenforms on  $\mathrm{SL}_2(\mathbb{Z})$ , this was recently established by El-Guindy and Raji [10], who showed that the zeros of  $r_f(z)$  (for N=1) are all on the unit circle |z|=1. Their work was inspired by the previous work by Conrey, Farmer and Imamoglu [11], who proved an analogous result for odd period polynomials again for full level. We show that this "Riemann hypothesis" holds in general for all newforms of weight at least 4 and any level.

**Theorem 1.1.** For any even integer k at least 4, and any level N, all of the zeros of the period polynomial  $r_f(z)$  are on the circle  $|z| = 1/\sqrt{N}$ .

**Remark.** Period polynomials for weight 2 newforms f are constant multiples of L(f,1).

**Example 1.** The period polynomial for the normalized Hecke eigenform  $\Delta(z) \in S_{12}(\Gamma_0(1))$  is

$$r_{\Delta}(z) = \omega_{\Delta}^{+} r_{\Delta}^{+}(z) + \omega_{\Delta}^{-} r_{\Delta}^{-}(z) \approx 0.114379i \left( \frac{36}{691} z^{10} - z^{8} + 3z^{6} - 3z^{4} + z^{2} - \frac{36}{691} \right) + 0.00926927(4z^{9} - 25z^{7} + 42z^{5} - 25z^{3} + 4z).$$

All ten zeros of  $r_{\Delta}(z)$  are on |z| = 1.

**Example 2.** For the unique weight 4 newform  $f(z) = q - 4q^3 - 2q^5 + \cdots$  on  $\Gamma_0(8)$ , we have

$$L(f,1) \approx 0.3545006..., \quad L(f,2) \approx 0.6900311..., \quad L(f,3) \approx 0.8746953...,$$

which in turn implies that  $r_f(z) \approx 0.0564205361iz^2 + 0.0349573870z - 0.00705256701815496i$ . The roots are  $\approx \pm 0.17037672 + 0.30979311i$ , and their norms are  $\approx 1/(2\sqrt{2})$ .

**Remark.** Manin [12] has used the work of Conrey, Farmer and Imamoglu [11] to construct zeta functions which satisfy the Riemann Hypothesis. He suggests that these polynomials arise from non-Tate motives and geometric objects lying below Spec  $\mathbf{Z}$  but not over  $\mathbf{F}_1$ . Using the  $P_f(z)$  defined below, one obtains further such polynomials mutatis mutandis.

If the weight or level is large enough, then the zeros of  $r_f$  are regularly spaced on the circle  $|z| = 1/\sqrt{N}$ . To state this conveniently, and for our later work, we shall put m := (k-2)/2 throughout and define

(1.5) 
$$P_f(z) = \frac{1}{2} {2m \choose m} \Lambda(f, \frac{k}{2}) + \sum_{j=1}^m {2m \choose m+j} \Lambda(f, \frac{k}{2}+j) z^j.$$

Then, using the functional equation, we see that

(1.6) 
$$r_f\left(\frac{z}{i\sqrt{N}}\right) = i^{k-1}N^{-\frac{k-1}{2}}\epsilon(f)z^m\left(P_f(z) + \epsilon(f)P_f\left(\frac{1}{z}\right)\right).$$

Therefore, to understand the zeros of  $r_f$ , it is enough to understand the zeros of  $P_f(z) + \epsilon(f) P_f(1/z)$ , and Theorem 1.1 states that this function has all its zeros on the unit circle |z| = 1. If we restrict to the unit circle |z| = 1, then  $P_f(z) + \epsilon(f) P_f(1/z)$  is either a trigonometric cosine or a trigonometric sine polynomial (depending on whether  $\epsilon(f)$  equals 1 or -1), and our proof of Theorem 1.1 proceeds by finding the right number of sign changes as z varies over the unit circle. If k or N is large enough, the proof allows us to establish the following result on the location of the roots.

## Theorem 1.2. The following are true.

- (i) Suppose that k = 4. If  $\epsilon(f) = -1$ , then the zeros of  $r_f(z)$  are  $\pm i/\sqrt{N}$ . If  $\epsilon(f) = 1$  and N is sufficiently large, then the zeros of  $r_f(z)$  are located at  $\pm (1 + O(N^{-\frac{1}{4} + \epsilon}))/\sqrt{N}$ .
- (ii) If  $k \geq 6$  and either N or k is large enough, then the roots of  $r_f(z)$  may be written as

$$\frac{1}{i\sqrt{N}}\exp\left(i\theta_{\ell} + O\left(\frac{1}{2^{k}\sqrt{N}}\right)\right),\,$$

where for  $0 \le \ell \le 2m-1$  we denote by  $\theta_{\ell}$  the unique solution in  $[0,2\pi)$  to the equation

$$m\theta_{\ell} - \frac{2\pi}{\sqrt{N}}\sin\theta_{\ell} = \begin{cases} \frac{\pi}{2} + \ell\pi & if \ \epsilon(f) = 1\\ \ell\pi & if \ \epsilon(f) = -1. \end{cases}$$

Our arguments readily allow us to quantify the results in Theorem 1.2. For example, the arguments in Section 6 give that in part (ii) above, the implied O-constant may be taken as  $10^9$ , although this is a gross overestimate. The arguments in Section 5 locate sign changes even if the values of k or N are only moderately large.

Suppose that  $\epsilon(f) = 1$ . By counting sign changes, one consequence of Theorem 1.1 is that  $P_f(-1)$  has sign  $(-1)^m$ . In other words, if  $\epsilon(f) = 1$ , then we must have

(1.7) 
$$\frac{1}{2} {2m \choose m} (-1)^m \Lambda(f, \frac{k}{2}) + \sum_{j=0}^{m-1} (-1)^j {2m \choose 2m-j} \Lambda(f, k-1-j) > 0.$$

For any weight k, this inequality is clear for large enough N since the term j=0 above dominates all other terms. However it is interesting that such an inequality holds for all small weights and small level as well, and we wonder if it has any other significance. In Section 4 we give a proof of this inequality in the weight 6 case based on the Hadamard factorization formula. We also give there a more illuminating proof of this inequality based on the Riemann hypothesis for  $\Lambda(f,s)$ .

# 2. Preliminaries

Here we collect preliminary facts about L-functions which we shall find useful. The completed L-function  $\Lambda(f,s)$  is an entire function of order 1. Its zeros all lie in the strip  $|\operatorname{Re}(s) - \frac{k}{2}| < \frac{1}{2}$ , with the Riemann hypothesis predicting that all zeros lie on the line  $\operatorname{Re}(s) = \frac{k}{2}$ . Recall also that the central value  $\Lambda(f,\frac{k}{2})$  is known to be non-negative by the work of Waldspurger [13].

Hadamard's factorization formula applies to the entire function  $\Lambda(f,s)$ , and we may write

(2.1) 
$$\Lambda(f,s) = e^{A+Bs} \prod_{\rho} \left(1 - \frac{s}{\rho}\right) e^{s/\rho}.$$

Here the product is over all the zeros of  $\Lambda(f,s)$  (that is, the non-trivial zeros of L(f,s)), and A and B are constants. Note that if  $\rho$  is a zero then so too are  $\overline{\rho}$  and  $k-\rho$ . Since  $\Lambda(f,s)$  is real-valued on the real line, and in view of the functional equation, we have that B is real-valued and

$$B = -\sum_{\rho} \operatorname{Re} \frac{1}{\rho} = -\sum_{\rho} \frac{\operatorname{Re} (\rho)}{|\rho|^2}.$$

These considerations also show that for real s

(2.2) 
$$\Lambda(f,s) = e^A \prod_{\rho \in \mathbb{R}} \left( 1 - \frac{s}{\rho} \right) \prod_{\operatorname{Im}(\rho) > 0} \left| 1 - \frac{s}{\rho} \right|^2,$$

where we have paired the complex conjugate roots together so that the product is convergent.

**Lemma 2.1.** The function  $\Lambda(f,s)$  is monotone increasing for  $s \geq \frac{k}{2} + \frac{1}{2}$ . Moreover, we have

$$0 \le \Lambda(f, \frac{k}{2}) \le \Lambda(f, \frac{k}{2} + 1) \le \Lambda(f, \frac{k}{2} + 2) \le \dots$$

If  $\epsilon(f)$  is -1, then  $\Lambda(f, \frac{k}{2}) = 0$  and

$$0 \le \Lambda(f, \frac{k}{2} + 1) \le \frac{1}{2}\Lambda(f, \frac{k}{2} + 2) \le \frac{1}{3}\Lambda(f, \frac{k}{2} + 3) \le \dots$$

Monotonicity results such as Lemma 2.1 are familiar in the literature; for example, the Riemann hypothesis for L-functions is equivalent to the monotonicity of the absolute value of the completed L-function along horizontal lines starting from the critical line. In a different context Stark and Zagier observed a similar result in [14].

*Proof.* Since all the zeros lie in  $|\text{Re}(s) - \frac{k}{2}| < \frac{1}{2}$ , we see that  $|1 - s/\rho|$  is increasing for  $s \ge \frac{k}{2} + \frac{1}{2}$ . So by (2.2) it follows that  $\Lambda(f, s)$  is increasing in  $\text{Re}(s) \ge \frac{k}{2} + \frac{1}{2}$ . Further, we have

$$\left|1 - \frac{k/2}{\rho}\right| \le \left|1 - \frac{k/2 + 1}{\rho}\right|,$$

and so  $\Lambda(f, k/2) \leq \Lambda(f, k/2 + 1)$ . When  $\epsilon(f) = -1$ , we apply the same reasoning and now take into account that there must be a zero of odd order at  $\frac{k}{2}$ .

We record a useful inequality for L-values in the range of absolute convergence.

**Lemma 2.2.** If 0 < a < b and k is the weight of f, then we have

$$\frac{L(f, \frac{k+1}{2} + a)}{L(f, \frac{k+1}{2} + b)} \le \frac{\zeta(1+a)^2}{\zeta(1+b)^2}.$$

*Proof.* The Euler product for L(f,s) gives rise to

$$-\frac{L'}{L}(f,s) = \sum_{n=1}^{\infty} \frac{\Lambda_f(n)}{n^s},$$

where  $|\Lambda_f(n)| \leq 2n^{\frac{k-1}{2}}\Lambda(n)$  for all n. Here  $\Lambda(n)$  is the usual von Mangoldt function, and this estimate is an alternative way of encoding the Ramanujan bounds established by Deligne [15] (see also Li [2] for the Euler factors at primes dividing the level). The point is that for prime powers  $n = p^r$  we have  $\Lambda_f(n) = (\alpha_p^r + \beta_p^r) \log p$ , where the pth Fourier coefficient of f satisfies  $a(p) = \alpha_p + \beta_p$ .

Therefore, we have

$$\frac{L(f, \frac{k+1}{2} + a)}{L(f, \frac{k+1}{2} + b)} = \exp\left(\int_a^b -\frac{L'}{L}(f, \frac{k+1}{2} + t)dt\right) \le \exp\left(2\int_a^b -\frac{\zeta'}{\zeta}(1 + t)dt\right) = \frac{\zeta(1 + a)^2}{\zeta(1 + b)^2}.$$

#### 3. The weight 4 case

If f is a form of weight k=4 (so m=(k-2)/2=1), then  $P_f(z)=\Lambda(f,2)+\Lambda(f,3)z$ . If  $\epsilon(f)=-1$ , then the roots of  $P_f(z)-P_f(1/z)=\Lambda(f,3)(z-1/z)$  are at  $z=\pm 1$  and so the period polynomial has roots at  $\pm i/\sqrt{N}$ .

If  $\epsilon(f) = 1$ , then with  $z = e^{i\theta}$  we have  $P_f(z) + P_f(1/z) = 2\Lambda(f,2) + 2\Lambda(f,3)\cos\theta$ . Since  $\Lambda(f,2) < \Lambda(f,3)$  by Lemma 2.1, the above equation has two solutions for  $\theta \in [0,2\pi)$ : namely,  $\theta$  satisfying  $\cos\theta = -\Lambda(f,2)/\Lambda(f,3)$ . This completes the proof of Theorem 1.1 for weight 4.

Note that  $\Lambda(f,3) \gg N^{\frac{3}{2}}$  for large N, while the Phrágmen-Lindelöf principle gives  $\Lambda(f,2) \leq \max_{t \in \mathbb{R}} |\Lambda(f,\frac{5}{2}+\epsilon+it)| \ll N^{\frac{5}{4}+\epsilon}$  (this is the "convexity bound" for L-functions). Therefore the ratio  $\Lambda(f,2)/\Lambda(f,3)$  is small (precisely  $\ll N^{-\frac{1}{4}+\epsilon}$ ), and hence the corresponding values of  $\theta$  tend to  $\pi/2$  and  $3\pi/2$ . Thus for large level, the zeros of the period polynomial (in the  $\epsilon(f)=1$  case) are located at  $\pm (1+O(N^{-\frac{1}{4}+\epsilon}))/\sqrt{N}$ .

## 4. The weight 6 case

If f is a form of weight k=6 (so that m=2) then

$$P_f(z) = 3\Lambda(f,3) + 4\Lambda(f,4)z + \Lambda(f,5)z^2.$$

If  $\epsilon(f) = -1$ , then we are interested in the roots of

$$P_f(z) - P_f(1/z) = \left(z - \frac{1}{z}\right) \left(4\Lambda(f, 4) + \Lambda(f, 5)\left(z + \frac{1}{z}\right)\right).$$

Clearly there are two solutions  $z=\pm 1$ . Since  $\epsilon(f)=-1$ , we know that  $2\Lambda(f,4)<\Lambda(f,5)$  by Lemma 2.1, and so there are two solutions in  $[0,2\pi)$  to  $\cos\theta=-2\Lambda(f,4)/\Lambda(f,5)$ . Thus we have shown Theorem 1.1 in this case. Moreover if N is large, then  $\Lambda(f,4)/\Lambda(f,5)$  is small and  $\theta$  tends to  $\pi/2$  or  $3\pi/2$ . So for large N (and odd sign) the period polynomial has two zeros exactly at  $\pm i/\sqrt{N}$  and the other two zeros are very near  $\pm 1/\sqrt{N}$ .

It remains now to consider when  $\epsilon(f) = 1$ . With  $z = e^{i\theta}$  we must show that

(4.1) 
$$P_f(z) + P_f(1/z) = 2\cos(2\theta)\Lambda(f,5) + 8\cos\theta\Lambda(f,4) + 6\Lambda(f,3)$$

has two zeros in  $[0, \pi]$  (and therefore four zeros in  $[0, 2\pi)$ ). Differentiating the above with respect to  $\theta$  gives  $-8\sin\theta(\Lambda(f, 4) + \cos\theta\Lambda(5))$  so that there are critical points at  $\theta = 0$ ,  $\pi$ , and at the solution  $\theta_0 \in (0, \pi)$  to  $\cos\theta = -\Lambda(f, 4)/\Lambda(f, 5)$ . We would like the quantity in (4.1) to be positive at  $\theta = 0$ , negative at  $\theta_0$  and positive again at  $\theta = \pi$ , which would ensure two zeros in  $(0, \pi)$  (and note that these conditions are also necessary for the period polynomial to have all zeros on a circle).

The value at  $\theta = 0$  is clearly positive. That the value should be positive at  $\pi$  is equivalent to

$$\Lambda(f,5) + 3\Lambda(f,3) > 4\Lambda(f,4).$$

The condition that the value should be negative at  $\theta_0$  is equivalent to

(4.3) 
$$\Lambda(f,5)^2 + 2\Lambda(f,4)^2 \ge 3\Lambda(f,3)\Lambda(f,5).$$

**Lemma 4.1.** Suppose  $a_1$ ,  $a_2$ ,  $b_1$ ,  $b_2$ , and  $c_1$ ,  $c_2$  are all positive with  $a_i \ge \max(b_i, c_i)$ . Suppose that  $a_i + \gamma c_i \ge (1 + \gamma)b_i$ , where  $\gamma$  is positive. Then  $a_1a_2 + \gamma c_1c_2 \ge (1 + \gamma)b_1b_2$ .

*Proof.* Multiply the relation  $a_1 + \gamma c_1 \ge (1 + \gamma)b_1$  by  $b_2$ . It suffices to show that

$$a_1a_2 + \gamma c_1c_2 \ge a_1b_2 + \gamma c_1b_2$$
;

or, rearranging that  $a_1(a_2 - b_2) \ge \gamma c_1(b_2 - c_2)$ . Since  $(a_2 - b_2) \ge 0$ , and  $a_1 \ge c_1$ , the LHS above is at least  $c_1(a_2 - b_2)$  which is  $\ge \gamma c_1(b_2 - c_2)$ .

Proof of (4.2). We use Lemma 4.1 suitably, together with the Hadamard factorization formula ((2.1) and (2.2)), proceeding zero by zero. We use the Hadamard formula for  $\Lambda(f,3)$ ,  $\Lambda(f,4)$  and  $\Lambda(f,5)$ ; note that at all these values  $\Lambda$  is known to be non-negative (This is clear for 4 and 5, and work of Waldspurger for 3.), so we can also assume that the products are taken with absolute values.

Suppose first that  $\rho = 3 + z$  is a real zero, and then  $6 - \rho = 3 - z$  is also a real zero. (Note that even if  $\rho = 3$ , we get zeros of even multiplicity at the center, which may be paired.) Then note that this pair of zeros contributes to  $\Lambda(f,5)$  the amount  $a = (4-z^2)/(9-z^2)$ , to  $\Lambda(f,4)$  the amount  $b = (1-z^2)/(9-z^2)$ , and to  $\Lambda(f,3)$  the amount  $c = z^2/(9-z^2)$  (using here the absolute value remark). Note that with  $\gamma = 3$  we have the inequality  $a + 3c \ge 4b$ .

Now consider a zero  $\rho=3+iy$  on the critical line, and pair it with its conjugate 3-iy. These contribute to  $\Lambda(f,5)$  the amount  $a=(4+y^2)/(9+y^2)$ , to  $\Lambda(f,4)$  the amount  $b=(1+y^2)/(9+y^2)$  and to  $\Lambda(f,3)$  the amount  $c=y^2/(9+y^2)$ , and we check again that  $a+3c\geq 4b$  (and indeed equality holds).

Finally consider a zero  $\rho=3+z$  not on the critical line with z=x+iy. This comes in a set of four zeros  $3\pm x\pm iy$ . Note that these four zeros contribute (multiply through by  $|\rho|^2|6-\rho|^2$ ) to  $\Lambda(f,5)$  an amount  $a=|4-z^2|^2$ , to  $\Lambda(f,4)$  an amount  $b=|1-z^2|^2$ , and to  $\Lambda(f,3)$  the amount  $c=|z^2|^2$ . We can check again that  $a+3c\geq 4b$ .

Thus, when grouped as above, each group of zeros appearing in the Hadamard formula satisfies a version of (4.2). By Lemma 4.1, taking products of these groups of zeros we again obtain a version of (4.2). Letting these products run over all zeros and taking the limit, we obtain (4.2).

*Proof of* (4.3). This proof is similar, appealing to Lemma 4.1 with  $\gamma = 2$  and using Hadamard's formula and grouping zeros as before.

The inequality (4.2) is implied by the usual Riemann Hypothesis for  $\Lambda(f,s)$ . Note that RH for  $\Lambda(f,s)$  implies also that the derivatives  $\Lambda^{(j)}(f,s)$  satisfy RH. Moreover, at the central point one sees that  $\Lambda^{(j)}(f,3)=0$  for all odd j, and that  $\Lambda^{(j)}(f,3)\geq 0$  for all even j. Therefore, taking Taylor expansions around 3, we see that

$$\Lambda(f,5) + 3\Lambda(f,3) = 4\Lambda(f,3) + \sum_{j=1}^{\infty} \frac{\Lambda^{(2j)}(f,3)}{(2j)!} 2^{2j} \ge 4\Lambda(f,3) + 4\sum_{j=1}^{\infty} \frac{\Lambda^{(2j)}(f,3)}{(2j)!} = 4\Lambda(f,4).$$

This reasoning in general explains why the period polynomial has the right sign at  $\pi$  (see (1.7)).

5. Weights between 8 and 14: Applications of results of Pólya and Szegö

Classical work of Pólya [16] and Szegö [17] considers trigonometric polynomials

$$u(\theta) = a_0 + a_1 \cos \theta + a_2 \cos(2\theta) + \dots + a_n \cos(n\theta),$$
  
$$v(\theta) = a_1 \sin \theta + a_2 \sin(2\theta) + \dots + a_n \sin(n\theta).$$

If  $0 \le a_0 \le a_1 \le a_2 \dots \le a_{n-1} < a_n$ , then Szegö showed that u and v both have exactly n zeros in  $[0,\pi)$  and that these zeros are simple. Each interval  $(\frac{\ell-\frac{1}{2}}{n+\frac{1}{2}}\pi,\frac{\ell+\frac{1}{2}}{n+\frac{1}{2}}\pi)$  for  $\ell=1,\ldots,n$  has precisely one zero of u, and apart from  $\theta=0$ , each interval  $(\frac{\ell}{n+\frac{1}{2}}\pi,\frac{\ell+1}{n+\frac{1}{2}}\pi)$  for  $1 \le \ell \le n-1$  has exactly one zero of v. His proof is a simple sign change argument using the positivity of the Fejér kernel.

When the level is suitably large, these results apply and provide a quick proof of Theorem 1.1. For weight k, for Szegö's theorem to apply we must verify the criteria

(5.1) 
$${2m \choose m} \Lambda(f, \frac{k}{2}) \le 2 {2m \choose m+1} \Lambda(f, \frac{k}{2}+1),$$

and for all  $1 \le j \le m-1$  that

(5.2) 
$${2m \choose m+j} \Lambda(f, \frac{k}{2}+j) \le {2m \choose m+j+1} \Lambda(f, \frac{k}{2}+j+1).$$

Since  $\Lambda(f, \frac{k}{2}) \leq \Lambda(f, \frac{k}{2} + 1)$ , the condition (5.1) is immediate for all  $k \geq 4$ . Now suppose  $k \geq 6$ . Using the definition of  $\Lambda$ , and simplifying a little, the condition (5.2) becomes (for  $1 \leq j \leq m-1$ )

$$\sqrt{N} \ge \frac{2\pi}{(k/2 - j - 1)} \frac{L(f, \frac{k}{2} + j)}{L(f, \frac{k}{2} + j + 1)},$$

and by Lemma 2.2 we conclude that our criterion (5.2) is met if

(5.3) 
$$N \ge \max_{1 \le j \le k/2 - 2} \left(\frac{2\pi}{k/2 - j - 1}\right)^2 \frac{\zeta(j + 1/2)^4}{\zeta(j + 3/2)^4}.$$

For any given k, we can compute the bound (5.3). Thus, for k = 8, it suffices to take  $N \ge 142$ ; for k = 10 it suffices to have  $N \ge 64$ ; for k = 12 it suffices to have  $N \ge 45$ ; for k = 14 it suffices to have  $N \ge 42$ . We have used sage to check (5.2) for those newforms not covered by (5.3) for weights  $8 \le k \le 14$ . The zeros of those newforms which do not satisfy (5.2) still lie on  $|z| = 1/\sqrt{N}$ .

**Remark.** Eventually, this cannot furnish a bound better than  $4\pi^2$  for N, and so we must turn to another approach for large k and small N, which we carry out in the next section.

## 6. Larger Weights: A second approach

Here we consider larger weights by reformulating the previous approach of [11] and [10]. Recast the definition (1.5) of  $P_f(z)$  as

$$P_f(z) = (2m)! \left(\frac{\sqrt{N}}{2\pi}\right)^{2m+1} L(f, 2m+1)Q_f(z),$$

where

(6.1) 
$$Q_f(z) = z^m \sum_{j=0}^{m-1} \frac{1}{j!} \left(\frac{2\pi}{z\sqrt{N}}\right)^j \frac{L(f, 2m+1-j)}{L(f, 2m+1)} + \frac{1}{2(m!)^2} \left(\frac{2\pi}{\sqrt{N}}\right)^{2m+1} \frac{\Lambda(f, \frac{k}{2})}{L(f, 2m+1)}.$$

We wish to show that on the unit circle |z| = 1, the real and imaginary parts of  $Q_f(z)$  (which correspond to the even and odd signs of the functional equation) have exactly 2m zeros.

Now let us write

$$Q_f(z) = z^m \exp\left(\frac{2\pi}{z\sqrt{N}}\right) + S_1(z) + S_2(z) + S_3(z),$$

with

$$S_1(z) = z^m \sum_{j=1}^{m-1} \frac{1}{j!} \left( \frac{2\pi}{z\sqrt{N}} \right)^j \left( \frac{L(f, 2m+1-j)}{L(f, 2m+1)} - 1 \right),$$

$$S_2(z) = -z^m \sum_{i=m}^{\infty} \frac{1}{j!} \left(\frac{2\pi}{z\sqrt{N}}\right)^j$$
, and  $S_3(z) = \frac{1}{2(m!)^2} \left(\frac{2\pi}{\sqrt{N}}\right)^{2m+1} \frac{\Lambda(f, \frac{k}{2})}{L(f, 2m+1)}$ .

For  $z=e^{i\theta}$  on the unit circle, the argument of  $z^m \exp(2\pi/(z\sqrt{N}))$  is  $m\theta - 2\pi(\sin\theta)/\sqrt{N}$ , which is monotone increasing as  $\theta$  varies from 0 to  $2\pi$ , and changes by  $2\pi m$  overall. Therefore the real and imaginary parts of  $z^m \exp(2\pi/(z\sqrt{N}))$  both have exactly 2m zeros. More precisely, consider first the real part of  $z^m \exp(2\pi/(z\sqrt{N})) = \cos(m\theta - 2\pi(\sin\theta)/\sqrt{N}) \exp(2\pi(\cos\theta)/\sqrt{N})$ , and clearly we can find m values of  $\theta$  with  $\cos(m\theta - 2\pi(\sin\theta)/\sqrt{N}) = 1$  and m interlacing values where it is -1. Between two such interlacing values there must be a zero of the real part. Further, since  $\exp(2\pi(\cos\theta)/\sqrt{N}) \ge \exp(-2\pi/\sqrt{N})$  for all  $\theta$ , if

(6.2) 
$$|S_1(z) + S_2(z) + S_3(z)| < \exp\left(-\frac{2\pi}{\sqrt{N}}\right),$$

then the real part of  $Q_f(z)$  will also have sign changes and thus a zero in these intervals. A similar argument applies to the imaginary part of  $Q_f(z)$ , and so it suffices to check the criterion (6.2).

Now by Lemma 2.2 we see that  $L(f, 2m+1-j)/L(f, 2m+1)-1 \le \zeta(\frac{1}{2}+m-j)^2-1$  so that

$$|S_1(z) + S_2(z)| \le \sum_{j=1}^{m-1} \frac{1}{j!} \left(\frac{2\pi}{\sqrt{N}}\right)^j \left(\zeta(\frac{1}{2} + m - j)^2 - 1\right) + \sum_{j=m}^{\infty} \frac{1}{j!} \left(\frac{2\pi}{\sqrt{N}}\right)^j.$$

For the term j=m-1, note that  $\zeta(\frac{3}{2})^2-1\leq \frac{35}{6}$  by direct computation. Note that for  $2^x(\zeta(\frac{1}{2}+x)^2-1)$  is decreasing in  $x\geq 2$  and so may be bounded by  $4(\zeta(5/2)^2-1)\leq \frac{16}{5}$ . Using this observation for smaller values of j, we obtain

$$|S_1(z) + S_2(z)| \le \frac{16}{5} \sum_{j=1}^{m-1} \frac{1}{j!} \left(\frac{2\pi}{\sqrt{N}}\right)^j \frac{2^j}{2^m} + \frac{17}{4} \frac{1}{(m-1)!} \left(\frac{2\pi}{\sqrt{N}}\right)^{m-1} + \sum_{j=m}^{\infty} \frac{1}{j!} \left(\frac{2\pi}{\sqrt{N}}\right)^j \frac{2^j}{2^m}.$$

Combining the first and third terms, we conclude that

(6.3) 
$$|S_1(z) + S_2(z)| \le \frac{16}{5} 2^{-m} \left( \exp\left(\frac{4\pi}{\sqrt{N}}\right) - 1 \right) + \frac{17}{4} \frac{1}{(m-1)!} \left(\frac{2\pi}{\sqrt{N}}\right)^{m-1}.$$

To bound  $S_3(z)$ , note that  $\Lambda(f, \frac{k}{2}) \leq \Lambda(f, \frac{k}{2} + 1) \leq (\frac{\sqrt{N}}{2\pi})^{m+2} (m+1)! \zeta(\frac{3}{2})^2$ , so for  $m \geq 7$  we have

$$|S_3(z)| \le \frac{m+1}{2(m!)} \left(\frac{2\pi}{\sqrt{N}}\right)^{m-1} \zeta(\frac{3}{2})^2 \le \frac{4}{(m-1)!} \left(\frac{2\pi}{\sqrt{N}}\right)^{m-1}.$$

Combining this with (6.3), we conclude that

$$(6.4) |S_1(z)| + |S_2(z)| + |S_3(z)| \le \frac{16}{5} \frac{1}{2^m} \left( \exp\left(\frac{4\pi}{\sqrt{N}}\right) - 1 \right) + \frac{33}{4} \frac{1}{(m-1)!} \left(\frac{2\pi}{\sqrt{N}}\right)^{m-1}.$$

Thus, to verify the condition (6.2), we need only ensure that

(6.5) 
$$\frac{16}{5} \frac{1}{2^m} \left( \exp\left(\frac{4\pi}{\sqrt{N}}\right) - 1 \right) + \frac{33}{4} \frac{1}{(m-1)!} \left(\frac{2\pi}{\sqrt{N}}\right)^{m-1} < \exp\left(-\frac{2\pi}{\sqrt{N}}\right).$$

For values of m at least as large as the figure in the first row, the table below gives a bound N(m) such that estimate (6.5) holds for all  $N \ge N(m)$ :

m	29	21	18	16	14	13	12	11	10	9	8	7
N(m)	1	2	3	4	5	6	7	9	11	14	20	28

We used sage to confirm Theorem 1.1 for the finitely many newforms missed by (6.5).

## 7. Proof of Theorem 1.2.

The weight 4 case was already treated in Section 3. For  $m \geq 2$  (that is weights  $k \geq 6$ ), the argument in Section 6 shows that for  $z = e^{i\theta}$  on the unit circle we have

$$Q_f(z) = \exp\left(im\theta + \frac{2\pi}{\sqrt{N}}e^{-i\theta}\right) + O\left(\frac{1}{2^m\sqrt{N}}\right).$$

Thus we have that

$$\operatorname{Re}(Q_f(z)) = \exp\left(\frac{2\pi}{\sqrt{N}}\cos\theta\right)\cos\left(m\theta - \frac{2\pi}{\sqrt{N}}\sin\theta\right) + O\left(\frac{1}{2^m\sqrt{N}}\right).$$

For  $\theta \in [0, 2\pi)$  the first term above vanishes when  $m\theta - 2\pi(\sin\theta)/\sqrt{N} = \frac{\pi}{2} + \ell\pi$  with  $0 \le \ell \le 2m-1$ . For such a point  $\theta_{\ell}$ , if we consider the values at  $\theta_{\ell} - C/(2^m\sqrt{N})$  and  $\theta_{\ell} + C/(2^m\sqrt{N})$  for a suitable constant C > 0 (and if  $2^m\sqrt{N}$  is large enough) then  $\text{Re}(Q_f(z))$  has differing signs at these points, and hence a zero in between. When  $\epsilon(f) = 1$ , the zeros of the period polynomial  $r_f(z)$  are located at  $1/(i\sqrt{N})$  times the zeros of  $\text{Re}(Q_f(z))$ , and this proves Theorem 1.2 in this case. The case when  $\epsilon(f) = -1$  corresponds to  $\text{Im}(Q_f(z))$ , and a similar argument applies here.

## 8. Remarks on the calculations

In the previous sections we proved Theorem 1.1 for k=4,6 and  $k\geq 42$ . For  $8\leq k\leq 40$  finitely many newforms remain to complete the proof (see the discussions after (5.3) and (6.5)). We used inequality (5.3) for  $8\leq k\leq 14$ . The most levels remain for weight k=8; we are left to consider those newforms with  $N\leq 141$ . For weights  $16\leq k\leq 40$  we employed (6.5). The table after (6.5) gives the remaining levels. The most levels remain for weight k=16; we are left with  $N\leq 27$ .

Using sage we confirmed Theorem 1.1 for these remaining newforms. Running the commands CuspForms and newforms on a laptop, we had no difficulty computing these newforms. We then used Dokchitser's sage L-functions calculator to compute the values  $\Lambda(f,1), \ldots, \Lambda(f,k-1)$  to very high precision. We tested inequality (5.2), and found that it held for many of the remaining newforms. However, (5.2) fails for some newforms with low weight and level. For example, (5.2) fails for some weight k=8 newforms with  $N \in \{2,3,5-17,19\}$ .

For the forms which do not satisfy (5.2), we computed the trigonometric polynomials and checked that on the unit disk that they have the required number of sign changes for the truth of Theorem 1.1. As an example, consider the unique newform  $f \in S_{10}(\Gamma_0(12))$ . We have that

$$L(f,1) \approx 343.041936898889, L(f,2) \approx 140.422365373567, L(f,3) \approx 32.9164131544840,$$
  
 $L(f,4) \approx 6.41626479306637, L(f,5) \approx 1.71889934464323, \dots,$ 

which in turn implies for  $z = e^{i\theta}$  that

$$(P_f(z) + \epsilon(f)P_f(1/z))/2 \approx 189.128932153817\cos(4\theta) + 341.466246468159\cos(3\theta) + 308.910589184567\cos(2\theta) + 199.188643773093\cos(\theta) + 73.5501402820398.$$

This has four zeros for  $\theta \in [0, \pi)$  as required, and they are in the intervals

$$\left(\frac{4\pi}{20}, \frac{5\pi}{20}\right), \left(\frac{10\pi}{20}, \frac{11\pi}{20}\right), \left(\frac{14\pi}{20}, \frac{15\pi}{20}\right), \left(\frac{18\pi}{20}, \frac{19\pi}{20}\right).$$

#### References

- [1] A. O. L. Atkin and J. Lehner, Hecke operators on  $\Gamma_0(m)$ , Math. Ann. 185 (1970), 134-160.
- [2] W. Li, Newforms and functional equations, Math. Ann. 212 (1975), 285–315.
- [3] Y. Choie, Y. K. Park, and D. Zagier, Periods of modular forms on  $\Gamma_0(N)$  and products of Jacobi theta functions, in press.
- [4] M. Knopp, Some new results on the Eichler cohomology of automorphic forms, Bull. Amer. Math. Soc. 80 (1974), 607–632.
- [5] W. Kohnen and D. Zagier, *Modular forms with rational periods*, Modular forms (Durham, 1983), Ellis Horwood Ser. Math. Appl.: Statst. Oper. Res., Horwood, Chichester, 1984, 197-249.
- [6] V. Paşol, A. Popa, Modular forms and period polynomials, Proc. Lond. Math. Soc. (3) 107 (2013), no. 4, 713–743.
- [7] D. Zagier, Periods of modular forms and Jacobi theta functions, Invent. Math. 104 (1991), 449-465.
- [8] S. Gun, M. R. Murty, and P. Rath, Transcendental values of certain Eichler integrals, Bull. London Math. Society 43 (2011), no. 5, 939-952.
- [9] M. R. Murty, C. Smyth and R.J. Wang, Zeros of Ramanujan polynomials, J. Ramanujan Math. Society, 26 (2011), 107-125.
- [10] A. El-Guindy, W. Raji, Unimodularity of zeros of period polynomials of Hecke eigenforms, Bull. Lond. Math. Soc. 46 (2014), no. 3, 528–536.
- [11] J.B. Conrey, D.W. Farmer, O. Imamoğlu, The nontrivial zeros of period polynomials of modular forms lie on the unit circle, Int. Math. Res. Not. (2013), no. 20, 4758–4771.
- [12] Y. I. Manin, Local zeta factors and geometries under Spec Z, Izv. Russian Acad. Sci. (Volume dedicated to J.-P. Serre), accepted for publication, arXiv:1407.4969.
- [13] J.-L. Waldspurger, Sur les valeurs de certaines fonctions L-automorphes en leur centre de symétrie [On the values of certain automorphic L-functions at the center of symmetry], Compositio Math. 54 (1985), 173-242. French
- [14] H. Stark and D. Zagier, A property of L-functions on the real line, J. Number Th. 12 (1980), 49-52.
- [15] P. Deligne, La conjecture de Weil. I, Inst. Hautes Études Sci. Publ. Math. No. 43 (1974), 273–307.
- [16] G. Pólya, Über die Nullstellen gewisser ganzer Funktionen [On the zeros of certain entire functions], Math. Zeit., vol. 2 (1918), 353–383. German
- [17] G. Szegö, Inequalities for the zeros of Legendre polynomials and related functions, Trans. Amer. Math. Soc. vol. 39 (1), (1936), 1–17.

School of Mathematics, Korea Institute for Advanced Study, Hoegiro 85, Dongdaemun-gu, Seoul 130-722, Korea

 $E ext{-}mail\ address: seokhojin@kias.re.kr}$ 

SCHOOL OF MATHEMATICS, SHANDONG UNIVERSITY, JINAN, SHANDONG, 250100, CHINA

 $E ext{-}mail\ address: wenjunma.sdu@hotmail.com}$ 

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, EMORY UNIVERSITY, ATLANTA, GEORGIA 30322

E-mail address: ono@mathcs.emory.edu

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, STANFORD, CALIFORNIA 94305

E-mail address: ksound@math.stanford.edu